

Foot of a. Endon Profile
of same.

E.A.:

87-14

7L
A m 2, N14
N25-8A-14
196115


JOURNAL
OF THE
Indian Institute of Science.

CONTENTS.

CONSTITUTION OF MANASSE'S HYDROXYCAMPHOR.

BY

M. O. Forster and P. P. Shukla.



DR. M. O. FORSTER, F.R.S., CHAIRMAN OF EDITORIAL BOARD.

JOURNAL
OF THE
INDIAN INSTITUTE OF SCIENCE.

VOLUME I.—(1914-18).

Parts I-XXI.

VOLUME II.—(1918-19).

Part I.—The Training of Students in Applied Chemistry. II. Indian Essential Oils. III. The Oleo-resin from *Hardwickia pinnata*. IV. The Oleo-resin from *Dipterocarpus indicus*. V. i. The Vapour Pressures of Acetone. II. The Vapour Pressures of concentrated Solutions. VI. Note on the Oil from Compressed Oil Gas. VII. Wood Distillation. Parts i. and ii. VIII. i. Studies on Alcoholysis. II. Alcoholysis in the absence of a catalyst. IX. Tests for Ganja and other products from *Cannabis indica*. X. Some Experiments on the Cathode Fall. Part i. in Neon and Helium. Part ii. in Argon, Hydrogen and Nitrogen. XI. The Hydrolysis of Napthol Ethers. XII. The Enolisation of Carbonyl Compounds under the Influence of Grignard's Reagents. XIII. Some Enzymes of the Leaf of *Lantana camara*. XIV. A Laboratory Vacuum Film Evaporator. XV. Oil Splitting by Castor Seed Lipase. Parts i. to iv.

VOLUME III.—(1920).

Part I.—Studies on Alcoholysis III. The Alcoholysis of Aromatic Esters and the inhibiting influence of Ortho-substituents. II.—An Automatic Separator for use in Essential Oil Distillation. III.—The Starting Conditions in Synchronous Machines. IV.—Studies in the Fermentation of Cellulose. V.—Tartaric Acid from Tamarinds. VI.—Studies in the Bio-Chemistry of the Mahua Flower. VII.—The Perishing of Paper in Indian Libraries. VIII.—The Conservation of Nitrogen with special reference to Activated Sludge. IX.—Wood Distillation. Part iii. The Distillation of Some Mysore and Baroda Woods and of Certain Waste Products: Part iv. The Tar from Coconut Shells.

VOLUME IV.—(1921).

Part I.—Research Notes on the Acetone Fermentation Process in India. II.—The Specific Gravities of Mixtures of Butyl Alcohol and Water. III.—A Suggested Method for the Extraction of Turpentine, Resin and Gum from the Gum-oleo-resin of *Boswellia Serrata* without the use of Solvents. IV.—Additive Compounds of Arylamines with Nitro-derivatives of Naphthalene. V.—The Manufacture of Thymol from Ajowan. VI.—Phase Transformation and Phase Balancing. VII.—Some experiments in the Use of Antiseptics in the Manufacture of Glue and Gelatine. VIII.—Studies relating to the Bacteria associated with Rice and other cereals. IX.—Preliminary Note on the Purification of Water by Activated Silt. X.—Additive Compounds of *m*-Dinitrobenzene. XI. The Action of Methyl and Ethyl Alcohols on Esters of 2:6 Dinitro- and 2:4:6-Trinitro-benzole acids. XII.—Studies in Alcoholysis IV. The Alcoholysis of esters of a β -unsaturated acids and of the corresponding saturated esters. XIII.—The Bio-Chemistry of the Indigenous Indigo Dye Vat. XIV.—The Development of Chemical Industries in India. XV.—Some experiments on the production of Power Alcohol and Paper Pulp from Megasse. XVI.—Stress and Tensions of Solid Copper Overhead Conductors under Indian Conditions.

Price, As. 8.]

[Vol. 8A, Part XIV, pp. 267 to 273.

JOURNAL

OF THE

Indian Institute of Science.


CONTENTS.

CONSTITUTION OF MANASSE'S HYDROXYCAMPHOR.

BY

M. O. Forster and P. P. Shukla.

DR M. O. FORSTER, F.R.S., CHAIRMAN OF EDITORIAL BOARD.



2

A 2. N 14

N 23. 8A. 14

CONSTITUTION OF MANASSE'S HYDROXYCAMPHOR.¹

By M. O. Forster and P. P. Shukla.

At various times the name hydroxycamphor, or *oxycampher* in German, has been applied to numerous compounds having the empirical formula, $C_{10}H_{16}O_2$. Beilstein classifies campholenic acid under that heading, and an early paper by R. Schiff² mentions without reference one substance prepared by Wheeler melting at 137° , and another by Schmiedeberg and Meyer with the alternative name camphol. Schiff's own contribution to the series was produced in association with much camphoric acid by the action of nitrous acid on α -aminocamphor, and melted at $154-155^\circ$, whilst Kachler and Spitzer³ oxidised the hydrocarbon, $C_{10}H_{16}$ from camphor dichloride with chromic acid, and obtained a small quantity of a substance melting at $59-61^\circ$. Schrötter⁴ oxidised bornyl acetate with chromic acid and obtained an acetyl derivative, $C_{12}H_{18}O_3$ hydrolysable to a hydroxycamphor which melted at $248-249^\circ$.

No further additions were made to the group until Manasse⁵ reduced camphorquinone with zinc dust and acetic acid, and subsequently⁶ claimed to recognise two isomeric substances, indistinguishable in appearance and optical activity; the ' α ' modification melted at $203-205^\circ$ with $[\alpha]_D 12.8^\circ$, whilst the ' β ' modification, produced by hydrolysing the solid methyl ether, melted at $212-213^\circ$ with $[\alpha]_D 12.3^\circ$.

It is unfortunate that Manassé denominated his materials in this perplexing manner, because it was already known that replacement of hydrogen in the camphor molecule might occur elsewhere than at the α -position; $\alpha\beta$ -dibromocamphor, originally called β -dibromocamphor, was first prepared in 1866 by Swarts and reinvestigated many times after 1881, whilst β -bromocamphor was isolated in 1901. Hence, at the time of publishing his experiments, Manasse might well have been aware that a substance called β -hydroxycamphor should correspond in orientation with β -bromocamphor; confusion.

¹ Excepting the opening paragraphs, this paper was published under the title 'Studies in the Camphane Series. Part XL. Constitution of Manasse's Hydroxycamphor,' in the *Journal of the Chemical Society*, 1925, 127, 1855.

² *Ber.*, 1880, 13, 1404.

³ *Annalen*, 1880, 200, 358.

⁴ *Montash.*, 1881, 2, 224.

⁵ *Ber.*, 1897, 30, 659.

⁶ *Ber.*, 1902, 35, 3811.


with true β -hydroxycamphor, obtained by Forster and Howard¹ on hydrolysing with oxalic acid the semicarbazone, $C_{11}H_{19}O_2N_3$ produced from β -aminocamphor and semicarbazide, might thus have been avoided.

Finally, Bredt and Goeb² re-examined Schrötter's acetoxycamphor and found that his hydroxycamphor is oxidisable to a *paradiketo*-camphane which, being optically active, must have the definite constitution implied by that name. They were further led to regard Schrötter's hydroxycamphor as a mixture of the *exo* and *endo* forms, of which the former, corresponding to borneol, melts at $237.5-238^\circ$.

Thus, among all these forms of hydroxycamphor, the only compounds which appear to arise by replacement of an α -hydrogen atom in the camphor molecule are those of Schiff and of Manasse, and our object has been to explain the isomerism of the two modifications which the latter investigator claimed to recognise.

Manasse supported his belief in the existence of two distinct modifications by the production of two oximes, m.p. $86-87^\circ$ (α) and $83-84^\circ$ (β), two phenylhydrazones, m.p. 137.5° (α) and $111-113^\circ$ (β), two semicarbazones, m.p. $182-183^\circ$ (α) and $202-204^\circ$ (β) and two benzenesulphonyl derivatives, m.p. $95-96^\circ$ (α) and $111-113^\circ$ (β). Excepting the last-named, however, these pairs might have arisen from a single hydroxycamphor, and in view of the slender distinction between the two parents, one of us (M. O. F.) with Dr. A. Zimmerli began in 1911 an inquiry which was interrupted and could not be resumed until eighteen months ago.

Simultaneous light and shadow have now been cast on the subject by Karrer and Takashima,³ who suggest that the solid methyl ether is the *cycloacetal* (I) and not the true α -methoxycamphor formulated by Manasse (II):




Thereupon they propose for Manasse's α - and β -hydroxycamphors the representation (III), ascribing to configuration-change in one of the asterisked carbon atoms the respective divergence in properties.

¹ *J. Chem. Soc.*, 1913, 103, 63.


² *J. pr. Chem.*, 1920 [ii], 101, 273.

³ *Helv. Chim. Acta*, 1925, 8, 242.

Our experiments amply confirm the above suggestion that the solid methyl ether is a *cycloacetal*, but we cannot support the complete constitutional formula given by Karrer and Takashima. We prefer the alternative representation (IV) with the consequence that Manasse's ' β '-hydroxycamphor is the individual (V), whilst his ' α '-hydroxycamphor is probably that substance associated with a variable amount of the stereoisomeride in which the hydroxyl group and hydrogen atom both occupy the alternate plane.



With Karrer and Takashima, we agree that the solid methyl ether does not reveal a carbonyl group; we failed to produce an oxime and recovered the substance unchanged by treatment with magnesium methyl iodide. Moreover, on treating separately (1) the crude product of reducing camphorquinone, (2) ' α '-hydroxycamphor, and (3) ' β '-hydroxycamphor with excess of the Grignard agent, the quantitative amount of methane required by one hydroxyl group was liberated in each case, and the original product was recovered unchanged. If Manasse's hydroxycamphor had the conventional constitution (VI), it could scarcely escape at least partial conversion into the secondary-tertiary diol (VII), in view of the readiness with which camphorquinone is transformed into the dimethyldiol by magnesium methyl iodide.¹




Our reason for questioning the transference of the hydroxyl group from the true α -position (in V) to the adjacent carbon atom (in III), as represented by Karrer and Takashima, is the one familiar to all chemists who have closely studied the members of this family, namely, the greatly superior activity of the α -carbon atom to that of its ketonic neighbour. The comparative behaviour of camphor and of camphorquinone towards aniline, phenylhydrazine, hydroxylamine, semicarbazide, nitromethane, ethyl cyanoacetate, and phenylacetonitrile may be recalled in this connexion: and finding that Manasse's hydroxycamphor is readily formed by reducing a cold ethereal solution of camphorquinone with aluminium amalgam, we

¹ Forster, *J. Chem. Soc.*, 1905, 87, 241.

cannot believe that in this reaction alone the diketone has belied itself.

Aside from the question of orientation, however, the above diagnosis by Karrer and Takashima has, in our judgment, clarified a situation which has been hitherto most obscure, and harmonises with the occurrence of Manasse's oximes, phenylhydrazones and semicarbazones, these doubtless arising from cleavage of the ethylene oxide-ring. The two benzenesulphonyl derivatives, to which we have added two hydrogen phthalic esters, are reasonably attributable to the stereoisomeric forms represented by formula V, because, although tertiary alcohols do not generally incur the formation of hydrogen phthalic esters, even limited occurrence of the equilibrium,



would account for the result observed. Moreover, the irregular course of ester-hydrolysis revealed by the polarimeter indicates the operation of concurrent mutarotation in the hydroxycamphor regenerated.

EXPERIMENTAL.


Reduction of Camphorquinone.—(a) *Zinc and acetic acid.* Camphorquinone (40 g.) dissolved in hot glacial acetic acid (70 c.c.) was precipitated by hot water (500 c.c.), and zinc dust (40 g.) added in small quantities to the continuously shaken suspension. The clear, pale green liquid was heated to coagulate undissolved zinc, the filtrate from which was saturated with salt and allowed to cool. Hydroxycamphor was filtered; after treating the filtrate with just sufficient sodium hydroxide (20%) to redissolve the zinc hydroxide, more was extracted by ether (three times), in which the filtered portion was then dissolved. Dried with calcium chloride, the ether was distilled and left a very pale yellow, camphoraceous residue (32.5 g.) (yield 80%).

(b) *Aluminium amalgam.* Camphorquinone dissolved in undried ether was left in contact with freshly amalgamated aluminium during 12 hours. The colour faded to very pale green, and the decanted liquid with ether-washings having been dried and distilled, deposited a colourless residue (yield 85%).

After crystallisation from petroleum the material prepared by each of the above methods melts at 203–205°; the less readily fusible product described by Manasse has been obtained only by hydrolysing

the solid methyl ether. This compound crystallises in colourless plates from a solution of the initial product in methyl alcohol containing hydrogen chloride (6%); it melts at 149–150° (Found: MeO, 15.9. Calc., MeO, 17.0%) and has $[\alpha]_D^{20}$ 182.8° (1% in chloroform). The methyl ether is not affected by boiling alcoholic potassium hydroxide (10%), but is rapidly hydrolysed by cold hydrochloric acid, from which sodium carbonate precipitates the less readily fusible form of hydroxycamphor; this melts at 215°, and in all other respects is indistinguishable from the material arising directly on reduction of camphorquinone.

The Hydroxyl Value of Hydroxycamphor.—At one time it seemed possible that the peculiar behaviour of hydroxycamphor might be due to its representing the equilibrium mixture,



but this is now precluded by several indications. In the first place, hydroxycamphor in chloroform does not decolorise bromine, and in ether does not develop a coloration with ferric chloride; moreover, the aqueous solution does not give a precipitate with copper acetate, and the methyl ether is equally indifferent to all these agents. When the action of bromine is pressed by heating hydroxycamphor with the halogen in acetic acid which has dissolved sodium acetate, a very slow action occurs; but this leads to camphoric anhydride without any indication of addition or substitution. The same treatment converts the solid methyl ether also into camphoric anhydride, and in both cases a considerable proportion of unchanged hydroxycamphor is recovered, having survived this attack during several hours.

Conclusive evidence against the above possibility, however, was given by the behaviour of hydroxycamphor towards magnesium methyl iodide, from which the quantitative amount of methane for one hydroxyl group was liberated equally by the crude product of reducing camphorquinone and by specimens of hydroxycamphor melting at 203–205° and at 215°, whilst the methyl ether also remains unaltered by the Grignard agent without liberating methane. These experiments were conducted in phenetole, and appeared remarkable in view of the observations by Bredt-Savelsberg and others¹ relating to enolisation of camphor by the Grignard agent,

¹ J. pr. Chem., 1924, 107, 65.

which may exceed 40%; but they now receive their explanation in the *cycloacetal* representation of Karrer and Takashima.¹

The Camphoryl Hydrogen Phthalates.—A mixture of phthalic anhydride and hydroxycamphor (m. p. 215°) in equal weights, and therefore containing a slight molar excess of the anhydride, was heated at 110–120° during 8 hours and kneaded in a solution of sodium carbonate. On extracting the filtered liquid with ether, about 15% of the hydroxycamphor was recovered unchanged whilst the hydrogen phthalate was precipitated along with phthalic acid by neutralising the solution of sodium salts; it was separated by dissolution in chloroform, which deposited crystals on evaporation. Recrystallisation from benzene-petroleum gave *camphoryl hydrogen phthalate* in rectangular plates, m. p. 164–165° (Found: C, 68.2; H, 6.6. $C_{18}H_{20}O_5$ requires C, 68.3; H, 6.3%). The $[\alpha]_D$ in aqueous alcohol (1 : 1) was 53.7°.

On repeating this operation with hydroxycamphor melting at 203–205°, or with a specimen recovered unchanged by the methylation process, the proportion of unconverted hydroxycamphor was about the same, and the first crop of crystals melted at 152–156°; this was ultimately resolved into a mixture of the foregoing substance with an isomeric *camphoryl hydrogen phthalate* crystallising in slender needles, m. p. 147° (Found: C, 68.1; H, 6.4%). The $[\alpha]_D$ in aqueous alcohol (1 : 1) was 42.2°.

Thus there appear to be two definite hydrogen phthalates, but it has not been possible to decide whether hydrolysis leads to the respective parents. In fact, on following the hydrolysis polarimetrically, there is an indication of the result being an equilibrium system.


A clear solution prepared by mixing equal volumes of alcohol (97.5%) and aqueous potassium hydroxide (20%) was used. When dissolved in this, the higher-melting hydrogen phthalate passed from the initial $[M]_D$ 102.7° to 61.6° during 200 hours, while the isomeride changed from $[M]_D$ 55.0° to 64.4°. Intermediate stages were various, however, the higher-melting form having sunk to $[M]_D$ 60.3° when the isomeride was at 98.6° (after 50 hours), and rising again to $[M]_D$ 73.0° when the isomeride had fallen to 78.3° (after 145 hours). These changes are probably due to varying rates of hydrolysis overlapping unequal mutarotation of the resulting hydroxycamphor, because, in the same alcoholic alkali, higher-melting hydroxycamphor with initial $[M]_D$ 34.2° reaches a maximum of 82.6° in 26 hours and

¹ *Loc. cit.*

begins to fall after 48 hours, whilst the lower-melting form has initial $[M]_D$ 50.9°, reaches its maximum of 69.0° at about the same time as the other, and then remains constant. The later observations are only approximate because the liquids, although initially colorless, develop a brown tint which increases in depth with passage of time: moreover, the hydroxycamphor recovered from such solutions melts lower and more indefinitely than the original materials.

Department of Organic Chemistry,
Indian Institute of Science,
Bangalore.

[Accepted, 29-9-25.]



72
A-2, 214
225-8A-147

PRINTED AND PUBLISHED
BY GEORGE KENNETH AT THE DIOCESAN PRESS
POST BOX 155, MADRAS--1925. C10309

VOLUME V.—(1922).

Part I.—Studies in Acidolysis: I. The Equilibrium between Acetic Acid, Trichloroacetic Acid and their Ethyl Esters; II. The Equilibrium between Acetic Acid, Trichloroacetic Acid and their Methyl Esters. II.—An Attempted Synthesis of Norpinic Acid. III.—Experiments on the influence of *s*-Trinitrobenzene on Reactions in which Arylamines are involved. IV.—Induction Motors used as Synchronous Machines. V.—The Relation Between the Iodine Values and Refractive Indices of some Hardened Vegetable Oils, Part I. VI.—Performance of Three-phase Induction Motors on Unbalanced Voltages. VII.—The Production of Barium Salts from Barytes. VIII.—The Addition of Bromine to the α - and β -Chloro- and bromocinnamic Acids and their Esters. IX.—Oil Splitting by Castor-seed Lipase. Parts V—VII. X.—Some West Coast Vegetable Oils. XI.—Utilisation of Tamarind. XII.—West Australian 'Sandalwood Oil.' XIII.—The Extraction of Caffeine. XIV.—Mechanical Tests of Some Mysore Timbers.

VOLUME VI.—(1923).

Part I.—Mohna Oil. II.—Malabar Sardine Oil. III.—Studies in Esterification. IV.—The Action of Bases on $\alpha\alpha\beta$ - and $\alpha\beta\beta$ -Tribromo- β -phenylpropionic Acids and their Esters. V.—Hongay Oil. VI.—Cashew Kernel Oil. VII.—The Fruit of *Bassia Longifolia*. VIII.—Studies in Intensive Bacterial Oxidation, I. The Oxidation of Alcohol to Acetic Acid. IX.—Contributions to the Study of the Prickly Pear Problem. X.—A Chemical Examination of Samples of Water and of Shells from the Inle Valley. XI.—The Refining of Crude Saltpetre. XII.—Enolising Action of Hydrogen Chloride on Aldehydes and Ketones. XIII.—The Conversion of Eugenol into Isoeugenol.

VOLUME VII.—(1924).

Part I.—Studies in Alcoholysis: V. The Alcoholysis of Esters of $\alpha\beta$ -unsaturated Acids and of the corresponding saturated Esters, by B. Dasannacharya. Price, Re. 1.
Part II.—Chemical Factors in Denitrification, by Gilbert J. Fowler and V. N. Kotwal. Price, As. 12.
Part III.—The Retting of Coconut Husk for the Production of Coir, by Gilbert J. Fowler and F. Marsden. Price, Re. 1.
Part IV.—Reactions of Chromates at High Temperatures. Part I. The Synthesis and Decomposition of Calcium, Sodium and Magnesium Chromates in Air, by M. Raman Nayar, H. E. Watson and J. J. Sudborough. Price, Re. 1.
Part V.—Oil from *Mimusops Hexandra*: Rayan Oil, by C. K. Patel. Price, As. 12.
Part VI.—The Relation Between the Iodine Values and Refractive Indices of Hardened Oils, Part II, by H. E. Watson and J. J. Sudborough. Price, As. 12.
Part VII.—Contributions to the Scientific Study of the Lac Industry. Parts I—VIII, by Gilbert J. Fowler, S. Mahdihassan and M. Sreenivasaya. Price, Rs. 3.
Part VIII.—I. The Systematic Nomenclature of Polycyclic Carbon Systems, by J. J. Sudborough. II. Polycyclic and Cage Systems of Carbon Compounds, by J. J. Sudborough and P. Ramaswami Ayyar. III. The Systematic Nomenclature of Heterocyclic Compounds including Polycyclic Structures, by J. J. Sudborough. Price, Rs. 4.
Part IX.—The Use of Mixed Catalysts in the Hydrogenation of Oils, by C. K. Patel. Price, As. 12.

- Part X. Condensation of Aromatic Amines with Chloroform or Carbon Tetrachloride in the Presence of Finely Divided Copper, by R. C. Shah. Price, Re. 1.
- Part XI.—I. Notes on Bixin, by K. Venkataraman. II. An Alkaloid from Anona Squamosa Leaves, by N. Trimurti. Price, As. 12.
- Part XII.—Iodine as a Catalyst in Reactions Involving Elimination of Hydrogen Halides, by R. D. Desai. Price, Re. 1.
- Part XIII.—Studies Relating to the Symbiosis of Seeds and Bacteria, by Gilbert J. Fowler and Miss R. K. Christie. Price, Rs. 1-8-0.
- Part XIV.—The Bio-Genesis of Mahua Oil, by Gilbert J. Fowler and Talwar Dinanath. Price, Re. 1.
- Part XV.—Contributions to the Scientific Study of the Lac Industry. Parts IX—X, by Gilbert J. Fowler, S. Mahdihassan and M. Sreenivasaya. Price, As. 12.

VOLUME VIII (A)—(1925).

- Part I.—The Photochemical Oxidation of Aromatic Hydrocarbons. I, by J. J. Sudborough, H. E. Watson and B. T. Narayanan. Price, As. 8.
- Part II.—Kachi-Grass Oil, by Bijoor Sanjiva Rao and J. J. Sudborough. Price, Re. 1.
- Part III.—Argemone Oil, by S. Narayana Iyer, J. J. Sudborough and P. Ramaswami Ayyar. Price, As. 12.
- Part IV.—Essential Oil of Cyperus Rotundus, by B. Sanjiva Rao, P. B. Panicker and J. J. Sudborough. Price, As. 8.
- Part V.—Dehydration of Rectified Spirit by means of Anhydrous Calcium Chloride, by J. J. Sudborough and P. Ramaswami Ayyar. Price, As. 8.
- Part VI.—The Hydrolysis of the Amides of $\alpha\beta$ -Unsaturated Acids and of their Saturated Analogues, by A. R. Yathiraja and J. J. Sudborough. Price, Re. 1.
- Part VII.—I. Studies Relating to the Acetone-Producing Organisms, by Gilbert J. Fowler and V. Subramanyan. II. Mahua Flowers as Raw Material for the Acetone-Fermentation Process, by A. G. Gokhale. Price, Re. 1.
- Part VIII.—Studies in Esterification, by B. V. Bhide and J. J. Sudborough. Price, Rs. 1-8-0.
- Part IX.—Constituents of the Marking-Nut: 'Semecarpus Anacardium' Linn., by D. Satyanarayana Naidu. Price, As. 8.
- Part X.—Notes on Some Indian Essential Oils, by Bijoor Sanjiva Rao, J. J. Sudborough, and H. E. Watson. Price Rs. 2.
- Part XI.—Bromo-Derivatives of Para-Methoxycinnamic Acid, by K. V. Hariharan and J. J. Sudborough. Price, Rs. 1-8-0.
- Part XII.—I. An Examination of Some Gum-Enzymes, by Gilbert J. Fowler and M. A. Malandkar. II. Chemical Constitution of the Gum from 'Ecswellia Serrata,' by M. A. Malandkar. Price, Re. 1.
- Part XIII.—Industrial Wastes as Manure. I. Ajowan and Mohua Cakes as Fertilisers. II. Utilisation of Refuse, by R. D. Rege. Price, Re. 1.
- Part XIV.—Constitution of Manasse's Hydroxycamphor, by M. O. Forster and P. P. Shukla. Price, As. 8.