

Verified
pr

Journal of the Indian

Institute of Science

Vol A Part 2, 1927

Pp 15-31

5L

A.d.m.2, N1413

N27.10.2

196355

46'

Price, As. 12.]


[Vol. 10A, Part II, pp. 15 to 31.

JOURNAL

OF THE

Indian Institute of Science.

CONTENTS.



OILS AND FATS FROM THE SEEDS OF INDIAN FOREST PLANTS

PART VIII. THE OIL FROM THE SEEDS OF THEVETIA NERIIFOLIA
(JUSS.).

BY

Ramkanta Bhattacharya and P. Ramaswami Ayyar.

PART IX. THE OIL FROM THE SEEDS OF CERBERA ODOLLAM
(GAERTN.).

PART X. THE OIL FROM THE SEEDS OF NOLARRHENA
ANTIDYSENTERICA.

PART XI. THE OIL FROM THE SEEDS OF ANONA SQUAMOSA (LINN.).

BY

Ramachandra Vishnu Ghanekar and P. Ramaswami Ayyar.

A, d, m 2, N1415

22.10.2

196-55

STER. P.R.S. CHAIRMAN OF EDITORIAL BOARD.

JOURNAL
OF THE
INDIAN INSTITUTE OF SCIENCE

Parts I.-XXI. VOLUME I.—(1914-18).
Parts I.-XV. VOLUME II.—(1918-19).
Parts I.-IX. VOLUME III.—(1920).
Parts I.-IX. VOLUME IV.—(1921).

Part I.—Research Notes on the Acetone Fermentation Process in India. II.—The Specific Gravities of Mixtures of Butyl Alcohol and Water. III.—A Suggested Method for the Extraction of Turpentine, Resin and Gum from the Gum-oleo-resin of *Boswellia Serrata* without the use of Solvents. IV.—Additive Compounds of Arylamines with Nitro-derivatives of Naphthalene. V.—The Manufacture of Thymol from Ajowan. VI.—Phase Transformation and Phase Balancing. VII.—Some experiments in the Use of Antiseptics in the Manufacture of Glue and Gelatine. VIII.—Studies relating to the Bacteria associated with Rice and other cereals. IX.—Preliminary Note on the Purification of Water by Activated Silt. X.—Additive Compounds of *m*-Dinitrobenzene. XI.—The Action of Methyl and Ethyl Alcohols on Esters of 2:6 Dinitro- and 2:4:6-Trinitro-benzoic acids. XII.—Studies in Alcoholysis. IV. The Alcoholysis of esters of α - β -unsaturated acids and of the corresponding saturated esters. XIII.—The Bio-Chemistry of the Indigenous Indigo Dye Vat. XIV.—The Development of Chemical Industries in India. XV.—Some experiments on the production of Power Alcohol and Paper Pulp from Megasse. XVI.—Sags and Tensions of Solid Copper Overhead Conductors under Indian Conditions.

VOLUME V.—(1922).

Part I.—Studies in Acidolysis: I. The Equilibrium between Acetic Acid, Trichloroacetic Acid and their Ethyl Esters; II. The Equilibrium between Acetic Acid, Trichloroacetic Acid and their Methyl Esters. II.—An Attempted Synthesis of Norpinic Acid. III.—Experiments on the influence of *s*-Trinitrobenzene on Reactions in which Arylamines are involved. IV.—Induction Motors used as Synchronous Machines. V.—The Relation Between the Iodine Values and Refractive Indices of some Hardened Vegetable Oils, Part I. VI.—Performance of Three-phase Induction Motors on Unbalanced Voltages. VII.—The Production of Barium Salts from Barytes. VIII.—The Addition of Bromine to the α - and β -Chloro- and bromocinnamic Acids and their Esters. IX.—Oil Splitting by Castor-seed Lipase, Parts V-VII. X.—Some West Coast Vegetable Oils. XI.—Utilisation of Tamarind. XII.—West Australian 'Sandalwood Oil.' XIII.—The Extraction of Caffeine. XIV.—Mechanical Tests of Some Mysore Timbers.

VOLUME VI.—(1923).

Part I.—Mohua Oil. II.—Malabar Sardine Oil. III.—Studies in Esterification. IV.—The Action of Bases on $\alpha\alpha\beta$ - and $\alpha\beta\beta$ -Tribromo- β -phenylpropionic Acids and their Esters. V.—Hongay Oil. VI.—Cashew Kernel Oil. VII.—The Fruit of *Bassia Longifolia*. VIII.—Studies in Intensive Bacterial Oxidation, I The Oxidation of Alcohol to Acetic Acid. IX.—Contributions to the Study of the Prickly Pear Problem. X.—A Chemical Examination of Samples of Water and of Shells from the Inle Valley. XI.—The Refining of Crude Saltpetre. XII.—Enolising Action of Hydrogen Chloride on Aldehydes and Ketones. XIII.—The Conversion of Eugenol into Isoeugenol.

Price, As. 12.]

[Vol. 10A, Part II, pp. 15 to 31.

JOURNAL

OF THE



Indian Institute of Science.

CONTENTS.

OILS AND FATS FROM THE SEEDS OF INDIAN FOREST PLANTS.

PART VIII. THE OIL FROM THE SEEDS OF THEVETIA NERIFOLIA
(JUSS.).

BY

Ramkanta Bhattacharya and P. Ramaswami Ayyar.

PART IX. THE OIL FROM THE SEEDS OF CERBERA ODOLLAM
(GAERTN.).

PART X. THE OIL FROM THE SEEDS OF HOLARRHENA
ANTIDYSENTERICA.

PART XI. THE OIL FROM THE SEEDS OF ANONA SQUAMOSA (LINN.).

BY

Ramachandra Vishnu Ghanekar and P. Ramaswami Ayyar.

DR. M. O. FORSTER, F.R.S., CHAIRMAN OF EDITORIAL BOARD.



7L

A, d, m 2, N 14 15

N 27.10.12

PART VIII. THE OIL FROM THE SEEDS OF THEVETIA NERIIFOLIA (JUSS.).

By Ramkanta Bhattacharya and P. Ramaswami Ayyar.

The shrub *Thevetia Nerifolia* (Juss.), the yellow oleander, is commonly cultivated throughout India as an ornamental plant. It is well known that the fruits contain a highly poisonous glucoside which finds considerable use as a cattle poison. During the course of an investigation directed to preparing the pure glucoside and determining its constitution, a considerable quantity of the oil present in the seeds was obtained, and it appeared of interest to subject it to a detailed examination (cf. *Pharm. Indica*, Vol. II, 409).

The fruit of *T. Nerifolia* is globular, slightly fleshy, green, between 1.5 and 2 inches in diameter. It contains a hard nut which is light brown in colour and triangular in shape with a deep groove corresponding to the base of its triangle. Each nut contains two pale yellow seed kernels.

The nuts used in the present investigation were obtained from Madras. The average weight of a nut is 5 grams, the seed kernels forming approximately 25 per cent. of the nuts. The seeds, after decortication by hand, were extracted with light petroleum when a pale yellow oil was obtained, the oil content being about 57 per cent. An examination of the oil in the usual manner has shown it to contain the glycerides of palmitic, stearic, oleic, linolic and arachidic acids. In view of the low iodine value (76) the oil may be classed as a non-drying oil.

EXPERIMENTAL.

The oil remaining after the extraction of the seeds with light petroleum (b.p. 60–80°) and removal of the solvent had the constants given in Table I.

TABLE I.

n_{D}^{20}	...	...	...	...	...	0.903
n_{D}^{40}	...	...	...	...	...	1.4599
Acid value	...	...	...	...	...	4.3
Saponification value	...	...	...	...	...	194.1
Unsaponifiable matter	...	...	...	...	...	1.4 per cent.
Acetyl value	...	...	...	...	...	0.0
Iodine value	...	...	...	...	...	76.0
Polenske value	...	...	...	...	...	0.5
Reichert-Polenske value	...	...	...	...	...	0.4
Hehner value	...	...	...	...	...	95.6

¹ Part VII, *Ind. For. Rec.*, 1924, 10, 23.

For the determination of the fatty acids present the oil was saponified with an alcoholic solution of sodium hydroxide, the soap extracted with ether to remove unsaponifiable matter and the fatty acids regenerated from the salt by acidification with hydrochloric acid. The mixed fatty acids had the constants given in Table II.

TABLE II.

n_D^{40}	...	...	...	...	...	...	1.4531
Iodine value	...	...	...	...	...	...	77
Titre test	...	...	...	...	...	...	34°
M.W.	...	...	...	...	...	...	277.5

The mixture of fatty acids was separated into the saturated and unsaturated acids by Twitchell's process (*Journ. Ind. Eng. Chem.*, 1921, 13, 806) the operation being repeated twice. In Table III the result of the separation is summarised.

TABLE III.

	Saturated Acids (29.3 per cent.)	Unsaturated Acids (70.6 per cent.)
Iodine value	0.4	100.0
M.W.	272	279.6
n_D^{60}	1.4361	n_D^{40} 1.4464

Identification of the Saturated Acids.—The crude mixture of saturated acids was converted into the methyl esters in the usual manner and the esters repeatedly fractionated under diminished pressure with the aid of a column, when ultimately nine fractions were obtained. The results of examining these fractions are given in Table IV. From the data it is possible, assuming that only palmitic and stearic acids are present, to calculate the percentages of these acids in each fraction from the titres of the acids and esters by the methods described elsewhere (Lewkowitsch, 1921, I, 118; *this Journ.*, 1923, 6, 126). These values are given in columns VII, VIII and IX. It will be observed that the results are in fairly good agreement with one another except in the case of fraction 2.

The final fraction (9) was purified by repeated crystallisation from methyl alcohol when pure methyl arachidate m.p. 54° was ultimately isolated. A portion of the crude ester was hydrolysed and the resulting acid after crystallisation from alcohol melted at 75–76°, and was thus identified as arachidic acid. A determination by Renard's method (*Compt. rend.*, 1871, 73, 1330) of the arachidic acid in this

TABLE IV.

No. of Fraction	I B. P.	II Yield per cent.	III M.W. of ester	IV Titre of ester	V M.P. of acid	VI Titre of acid	Percentage of methyl stearate		
							VII from III	VIII IV	IX VI
1	176-178°/7 mm.	27.9	269.9	26.6°	62-63°	62.5°	0.0	0.0	0.0
2	176-180°/7 mm.	15.2	272.8*	26.2°	61.5°	61.0°	10	5	5
3	180°/11.5 mm.	9.9	282.5	26.7°	57.0°	56.3°	45	42	42
4	182-183°/11.5 mm.	10.0	286.0	28.5°	60.0°	57.5°	59	54	57
5	183-185°/11.5 mm.	6.9	287.5	28.8°	60.0°	58.8°	62	60	60
6	186-190°/11.5 mm.	5.2	282.4	31.0°	63.0°	63.0°	80	78	78
7	188-191°/11.5 mm.	8.8	285.5	34.5°	67.2°	67.2°	91	89	90
8	195-200°/7 mm.	7.5	286.8	36.5°	68.8°	68.9°	97	99	99
9	Residue and loss	8.6	310.1	40.4°	64.0°	...	57*	...	...

* This fraction was assumed to be a mixture of methyl stearate and methyl arachidate since fraction 8 was found to be nearly pure methyl stearate.

fraction showed it to be 43.3 per cent., in good agreement with the value calculated from the molecular weight of the ester.

From the data given above the composition of the mixture of saturated fatty acids is—

Palmitic acid				58.5 per cent.
Stearic acid				40.3 „ „
Arachidic acid				1.2 „ „

Identification of the Unsaturated Acids.—Prior to the separation of the liquid unsaturated acids by the ester method the additive bromo-compounds were prepared by Jamieson and Baughman's method (*Journ. Am. Chem. Soc.*, 1920, 42, 2398). It was found that 1.49 grams of the mixed acids gave 0.28 gram of tetrabromo-acid (m.p. 112°) whilst the non-crystalline residue was 2.09 grams (Br = 36.0. Calc. for dibromo-oleic acid, Br = 36.2 per cent.). A hexabromo-acid was not formed. From this it appeared probable that oleic and linolic acids alone were present, and this was confirmed by a fractional distillation of the methyl esters when the following results were obtained.

TABLE V.

No. of Fraction	B.P. (10 mm.)	Yield per cent.	M.W.	Iodine Value
1	185–190°	17.3	293	87
2	190–195°	24.9	295	87.5
3	195–200°	37.9	296	99
4	200–202°	11.8	296	100
5	Residue and loss.	8.1	296	100

(Iodine value of methyl oleate 86.0, methyl linolate 172.8).

From these results the conclusion may be drawn that the unsaturated acids consist of—

Oleic acid				91.0 per cent.
Linolic acid				9.0 „ „

Unsaponifiable Matter.—The unsaponifiable matter (1.4 per cent.) which had been extracted by ether from the sodium salts of the fatty acids was found by the digitonin method to contain 15.3 per cent. of a sterol. The sterol was purified by crystallisation from ether, when it

separated in radiating needles m.p. 137°. Its identity with sitosterol was confirmed by the preparation of the acetyl derivative, m.p. 129–130°.

SUMMARY.

The seeds of *Thevetia Neriifolia* (Juss.) yield 57 per cent. of a non-drying oil. The oil consists of the glycerides of the following fatty acids:—

Palmitic acid				17.1 per cent.
Stearic acid				11.8 „ „
Arachidic acid				0.4 „ „
Oleic acid				64.3 „ „
Linolic acid				6.3 „ „

Sitosterol was separated from the unsaponifiable matter which comprises 1.4 per cent. of the oil.

In conclusion we wish to express our indebtedness to Prof. J. L. Simonsen under whose direction this investigation was carried out.

TABLE V.

No. of Fraction	I B. P. (7 mm.)	II Yield per cent.	III M.W. of ester	IV Titre of ester	V M.P. of acid	VI Titre of acid	Percentage of methyl stearate calculated from			
							III	IV	V	VI
							VII			
1	172-174°	7.8	266.7	26.7°	60.5-61.5°	...	...	...	...	...
2	174°	11.0	271	25.4°	62-62.5°	61.6°	3.6	3.0	0.0	3.0
3	177-179°	36.7	272.5	24°	61-61.5°	59.1°	9.0	8.0	0.5	11.0
4	180-183°	22.3	276.2	23.5-24°	57-57.5°	55.6°	22.2	18.0	20.0	25.0
5	183-186°	5.9	283.4	26.5-27°	55.5-56°	56.3°	47.9	45.0	40.0	45.0
6	187-190°	4.8	289.2	28.5°	60.5-61°	59.5°	67.3	60.0	65.0	63.0
7	195-200°	7.6	298.4	36.2°	67.5-68.5°	...	99.9	97.0	95.0	...
8	Residue	3.9	343.4	...	64.5-65°	...†	44.8	...	...	...

* Mixture of myristic acid (11 per cent.) and palmitic acid (88.2 per cent.)

† Mixture of stearic and lignoceric acids.

The examination of the saturated acids thus shows them to have the following composition:—myristic acid? (0.9 per cent.), palmitic acid (72.9 per cent.), stearic acid (24.1 per cent.) and lignoceric acid (2.1 per cent. by difference).

Unsaponifiable matter.—The unsaponifiable matter (0.8 per cent. was purified by the digitonin method and the crude sterol (32.2 per cent.) was converted into the acetyl derivative which after crystallisation from alcohol melted at 125°. It consisted apparently mainly of acetyl sitosterol (m.p. 129°), but was not obtained in sufficient quantity for further purification.

SUMMARY.

The seeds of *Cerbera odollam* contain 43.1 per cent. of a non-drying oil. The oil contains the glycerides of linolic acid (16.4 per cent.), oleic acid (4.2 per cent.), myristic acid? (0.4 per cent.), palmitic acid (30 per cent.), stearic acid (9.9 per cent.) and lignoceric acid (0.9 per cent.).

The unsaponifiable matter forms 0.8 per cent. of the oil and contains 32.2 per cent. of a sterol, probably sitosterol.

PART X. THE OIL FROM THE SEEDS OF HOLARRHENA ANTIDYSENTERICA.

By Ramachandra Vishnu Ghanekar and P. Ramaswami Ayyar.

During the extraction of the alkaloid, conessine, from the seeds of *H. antidysenterica* (Journ. Chem. Soc., 1926, 2123) it was observed that the seeds contained an oil, and a considerable quantity having been obtained it appeared a desirable subject for detailed examination. The oil was present in the seeds to the extent of 19 per cent. and was a drying oil.

EXPERIMENTAL.

The oil was extracted from the finely ground seeds with light petroleum and was quite free from alkaloidal impurities. It had the following constants:—

TABLE I.

d_{15}^{15}	...	...	...	...	...	...	0.9354
n_D^{60}	...	...	...	...	...	...	1.4666
Acid value...	...	...	...	...	...	...	36.1
Saponification value	...	...	...	...	...	...	180.5
Unsaponifiable matter	...	...	...	...	...	...	3.5 per cent.
Acetyl value	...	...	...	...	...	...	22.9
Iodine value	...	...	...	...	...	...	149.1
Reichert-Meisel value	...	...	...	...	...	...	1.7
Reichert-Polenske value	...	...	...	...	...	...	0.4
Hehner value	...	...	...	...	...	...	94.3

After saponification the oil gave a mixture of fatty acids with the properties given in Table II.

TABLE II.

n_D^{60}	...	...	...	...	...	...	1.4597
Iodine value	...	...	...	...	...	...	151.3
Titre test	...	...	...	...	...	...	24.7°
M. W.	...	...	...	...	...	...	283.8

The mixed fatty acids were separated by the Twitchell process (*loc. cit.*) and the results are summarised in Table III.

TABLE III.

—	Saturated Acids (14.4 per cent.)	Unsaturated Acids (85.3 per cent.)
n_D^{60}	1.4484	1.4616
Iodine value	2.8	180.0
M. W.	290.0	280.0

The Unsaturated Acids.—The acids present in the mixture of liquid acids were identified by Jamieson and Baughman's process (*loc. cit.*) and were found to consist of a mixture of linolic acid (63.9 per cent.), linolenic acid (11.6 per cent.) and oleic acid (24.5 per cent. by difference). The bromination results from which these percentages were calculated are given in Table IV, and are in good agreement with the iodine value (180) since a mixture of acids in the proportions given above would have an iodine value of 179.

TABLE IV.

Bromo-derivatives of the Unsaturated Acids.

Quantity brominated	...	3.58 g	4.93 g
Yield of di- and tetrabromo-acids	...	6.11 g	8.45 g
Yield of hexabromo-acid	...	1.14 g	1.54 g
Br ₂ in crude bromo-acid	...	48.2 per cent.	48.3 per cent.
M. P. of tetrabromo-acid	...	112-113°	112-113°
M. P. of hexabromo-acid	...	179-180°	179-180°

The Saturated Acids.—The separation of the saturated acids was effected by distillation of the methyl esters, composition of the various fractions being determined in the usual manner.

The ester fractions 1-6 were hydrolysed and the acids examined. No evidence was found of the presence of any acids other than palmitic and stearic. Assuming the presence of these two acids the percentage composition of the mixed esters as determined from the titre of the ester, etc. and shown in columns VII, VIII, IX and X of the table are in fair agreement. Fraction 7 was nearly pure methyl stearate, but from fraction 8 an acid of m.p. 65°, M.W. 303 was separated. This acid was not homogeneous and since from fraction 9 it was found possible to separate lignoceric acid, m.p. 80°,

TABLE V.

No. of Fraction	I B.P. (8 mm.)	II Yield per cent.	III M. W. of ester	IV Titre of ester	V M. P. of acid	VI Titre of acid	Percentage of stearic acid calculated from			
							III	IV	V	VI
1	below 180°	13.9	273.3	23.8°	57-57.5°	..	12.0	16.0	19.0	..
2	180-182°	14.0	278.2	24.8°	55-55.5°	54.8°	29.0	28.0	30.0	30.0
3	182-185°	8.4	280.5	25.9°	56-56.5°	55.4°	37.5	35.0	36.0	35.0
4	185-188°	11.4	283.3	26.8°	56.5-56.8°	56°	47.5	45.0	50.0	40.0
5	189-193°	8.8	284	27.2°	56.6-57.2°	56.4°	50.0	46.5	52.0	50.0
6	195-198°	13.8	295.0	31.6°	67.3-67.8°	65.8°	90.0	80.0	89.0	87.0
7	203-205°	7.5	301.6	32.0°	63.5-64°	..	96.0†	..	..	..
8	208-215°	2.8	305.5	33.8°	62-62.5°	..	91.0†	..	..	..
9	Residue	19.4	340.2*	..	63.5-64°	..	33.0†	..	..	..

* This is the M. W. of the acid and not of the methyl ester.

† These are assumed to be mixtures of stearic and lignoceric acids.

M.W. 366.6, we have assumed it to be a mixture of stearic and lignoceric acids.

From the above results the mixed saturated acids have the following composition:—palmitic acid (39.1 per cent.), stearic acid (47.4 per cent.) and lignoceric acid (13.5 per cent. by difference).

Unsaponifiable Matter.—The unsaponifiable matter (3.5 per cent.) yielded on treatment with digitonin a sterol (17.4 per cent.). This was identified as phytosterol by the preparation of the acetyl derivative, m.p. 119-120°.

SUMMARY.

The seeds of *H. Antidysenterica* contain 19 per cent. of a drying oil composed of the glycerides of the following acids:—linolenic acid (10 per cent.), linolic acid (54.7 per cent.), oleic acid (21 per cent.), palmitic acid (5.6 per cent.), stearic acid (6.8 per cent.) and lignoceric acid (1.9 per cent.).

The unsaponifiable matter (3.5 per cent.) contains phytosterol (17.4 per cent.).

PART XI. THE OIL FROM THE SEEDS OF ANONA SQUAMOSA (Linn.).

By Ramachandra Vishnu Ghanekar and P. Ramaswami Ayyar.

Anona squamosa (Linn.), the custard apple tree, is a native of tropical America but has long been naturalised in India and occurs commonly in Southern India, being cultivated for its fruit. The seeds, leaves and immature fruits are stated to contain an acid resinous principle which is destructive to insect life (*Pharm. Indica*, I, 44). The oil in the seeds has not been investigated and a detailed study of the fatty acids obtained on its hydrolysis forms the subject of this communication.

EXPERIMENTAL.

The seeds were obtained from the fresh, ripe fruits and after air drying consisted of husks and shells 30 per cent., kernels 70 per cent. The seeds dried at 110° (loss in weight 33.3 per cent.) yield on extraction with light petroleum (b.p. 40–60°) 30 per cent. of an oil. The hot petrol extract on standing deposited a small quantity of a resin (1.2 per cent.); this was separated and, after removal of the solvent, an oil remained which had the constants given in Table I. It may be classed as a non-drying oil.

TABLE I.

$n_D^{15^\circ}$	...	...	...	...	...	...	0.9126
$n_D^{60^\circ}$	...	...	...	...	...	...	1.4558
Acid value	...	...	...	...	...	...	0.8
Saponification value	...	...	...	...	...	...	188.3
Unsaponifiable matter	...	...	...	...	...	...	0.2 per cent.
Acetyl value	...	...	...	...	...	...	18.5
Iodine value	...	...	...	...	...	...	85.6
Polenske value	...	...	...	...	...	...	0.1
Reichert-Polenske value	...	...	...	...	...	...	0.6
Höner value	...	...	...	...	...	...	93.8

The mixed fatty acids obtained on hydrolysis of the oil had, after removal of the unsaponifiable matter, the constants given in Table II.

TABLE II.

$n_D^{60^\circ}$	...	...	...	...	...	...	1.4470
Iodine value	...	...	...	...	...	...	84.8
Titre test	...	...	...	...	...	...	31.8°
M.W.	...	...	...	...	...	...	276.3

The properties of the saturated and unsaturated acids obtained after separation by the Twitchell process are summarised in Table III.

TABLE III.

	Saturated acids (26.6 per cent.)	Unsaturated acids (73.3 per cent.)
$n_D^{60^\circ}$	1.4382	1.4511
Iodine value	3.0	112.7
M.W.	271.9	281.6

The Unsaturated Acids.—From an examination of the bromo-acids only linolic acid (24.7 per cent.) and oleic acid (75.3 per cent.) appear to be present. This percentage composition agrees well with the observed iodine value (found, 112.7; calc., 112.5).

TABLE IV.

Bromo-derivatives of Unsaturated Acids.

Quantity brominated	...	4.406 g.	3.739 g.
Yield of di- and tetrabromo-acids	...	7.550 g.	6.400 g.
Br ₂ in crude bromo-acid	...	41.5 per cent.	41.4 per cent.
M.P. of tetrabromo-acid	...	112–113°	112–113°

The Saturated Acids.—The saturated acids were separated by fractional distillation of the methyl esters, composition of the various fractions being determined in the usual manner.

By the hydrolysis of the individual fractions it was found possible to separate palmitic and stearic acids in a pure state. From fraction 1 a liquid acid was isolated with an equivalent of 126.5. Owing to the limited quantity of material available it could not be identified. From fraction 8 a small amount of an acid was obtained which after repeated crystallisation had m.p. 78–79°, M.W. 392. This acid is possibly cerotic acid which is stated to melt at 78° and has M.W. 396. The percentage composition of the saturated acids is (see cols. VII to X) lower fatty acids (1.2), palmitic acid (55.4), stearic acid (40.1) and cerotic acid (3.3).

TABLE V.

No. of Fraction	I B.P.	II Yield per cent.	III M. W. of ester	IV Titre of ester	V M.P. of acid	VI Titre of acid	Percentage of Palmitic acid from				
							VII	VIII	IX	X	
							III	IV	V	VI	
1	161-166°/3mm.	9.7	255.7	18.5-19.2°	55-56°	...	89.0	...	...	...	
2	166-169°/3mm.	7.2	267.8	22.5-23°	58.5-59.5°	...	98.3	...	...	...	
3	170-173°/3mm.	23.8	275.4	24.2-24.5°	56.5-57.5°	...	80.7	75.0	80.0	...	
4	175-178°/3mm.	3.8	278.4	26-26.6°	55.5-56.5°	54.9°	70.0	65.0	65.0	70.0	
5	185-187°/4mm.	7.0	280.5	26.4-26.6°	56-56.5°	-55.3°	62.5	58.0	60.0	65.0	
6	190-192°/4mm.	36.8	287.9	29.5-29.8°	62.5-63°	60.7°	36.1	30.0	30.0	33.0	
7	193-195°/4mm.	5.3	297.5	35-35.5°	66-66.5°	...	1.8	5.0	15.0	...	
8	Residue ...	6.4	341.3	...	65-68°	...	48.9°	...	...	...	

* Assumed to be a mixture of methyl stearate and methyl cerotate.


Unsaponifiable Matter.—The oil contained 0.2 per cent. of unsaponifiable matter which on treatment with digitonin gave 15.9 per cent. of a sterol. This was identified as sitosterol by the preparation of the acetyl derivative, m.p. 128-129°.

SUMMARY.

The seeds of *Anona squamosa* contain 14 per cent. of a non-drying oil and 0.56 per cent. of a neutral resin. The oil is composed of the glycerides of oleic acid (18.1 per cent.), linolic acid (55.2 per cent.), palmitic acid (14.7 per cent.), stearic acid (10.7 per cent.) and cerotic acid? (0.9 per cent.). An unidentified liquid saturated acid (0.3 per cent.) was also present. The unsaponifiable matter (0.2 per cent.) contains sitosterol (15 per cent.).

Department of Organic Chemistry,
Indian Institute of Science,
Bangalore.

[Accepted, 2-2-27.]



A, d, m2, N1423

N27.10A.2

VOLUME VII.—(1924).

Part I.—Studies in Alcoholysis V. The Alcoholysis of Esters of $\alpha\beta$ -unsaturated Acids and of the corresponding saturated Esters. II.—Chemical Factors in Denitrification. III.—The Retting of Coconut Husk for the Production of Coir. IV.—Reactions of Chromates at High Temperatures. Part I. The Synthesis and Decomposition of Calcium, Sodium and Magnesium Chromates in Air. V.—Oil from *Mimusops Hexandra*: Rayan Oil. VI.—The Relation Between the Iodine Values and Refractive Indices of Hardened Oils. Part II. VII.—Contributions to the Scientific Study of the Lac Industry. Parts I—VIII. VIII.—I. The Systematic Nomenclature of Polycyclic Carbon Systems. II. Polycyclic and Cage Systems of Carbon Compounds. III. The Systematic Nomenclature of Heterocyclic Compounds including Polycyclic Structures. IX.—The Use of Mixed Catalysts in the Hydrogenation of Oils. X.—Condensation of Aromatic Amines with Chloroform or Carbon Tetrachloride in the Presence of Finely Divided Copper. XI.—I. Notes on Bixin, by K. Venkataraman. II. An Alkaloid from *Anona Squamosa* Leaves. XII.—Iodine as a Catalyst in Reactions Involving Elimination of Hydrogen Halides. XIII.—Studies Relating to the Symbiosis of Seeds and Bacteria. XIV.—The Bio-Genesis of Mahua Oil. XV.—Contributions to the Scientific Study of the Lac Industry. Parts IX—X.

VOLUME VIII (A)—(1925).

Part I.—The Photochemical Oxidation of Aromatic Hydrocarbons. I. II.—Kachi-Grass Oil. III.—Argemone Oil. IV.—Essential Oil of *Cyperus Rotundus*. V.—Dehydration of Rectified Spirit by means of Anhydrous Calcium Chloride. VI.—The Hydrolysis of the Amides of $\alpha\beta$ -Unsaturated Acids and of their Saturated Analogues. VII.—I. Studies Relating to the Acetone-Producing Organisms. II. Mahua Flowers as Raw Material for the Acetone-Fermentation Process. VIII.—Studies in Esterification. IX.—Constituents of the Marking-Nut: '*Semecarpus Anacardium*' Linn. X.—Notes on Some Indian Essential Oils. XI.—Bromo-Derivatives of Para-Methoxycinnamic Acid. XII.—I. An Examination of Some Gum-Enzymes. II. Chemical Constitution of the Gum from '*Boswellia Serrata*'. XIII.—Industrial Wastes as Manure. I. Ajowan and Mohua Cakes as Fertilisers. II. Utilisation of Refuse. XIV.—Constitution of Manasse's Hydroxycamphor. XV.—The Reaction Between Sodium Sulphite and Sulphur. XVI.—The Occurrence of Sylvestrene.

VOLUME VIII (B)—1925.

Part I.—Lecher Wire Measurements. II.—The Aluminium Anode Film Dielectric.

VOLUME IX (A)—1926.

Part I.—Contributions to the Scientific Study of the Lac Industry. Part XI. Early Recognition of Sex among Lac Insects, by S. Mahdihassan. Price, Rs. 4.
 Part II.—Vegetable Oils Containing Glycerides of Erucic Acid, by J. J. Sudborough, H. E. Watson and P. Ramaswami Ayyar. Price, Rs. 2-8.
 Part III.—The Production of Ether by Solid Catalysts, by S. K. Kulkarni Jatkar and H. E. Watson. Price Rs. 3.
 Part IV.—The Constituents of Some Indian Essential Oils. Part XVIII. Derivatives of Abietic Acid, by Madyar Gopal Rau and John Lionel Simonsen. Price, As. 8.

PRINTED AND PUBLISHED
 BY GEORGE KENNETH AT THE DIOCESAN PRESS,
 POST BOX 455, MADRAS 1927. C2507

- Part. V.—The Fat from 'Salvadora Oleoides': Khakan Fat, by C. K. Patel, S. Narayana Iyer, J. J. Sudborough and H. E. Watson. Price, As. 12.
- Part VI.—The Constituents of Some Indian Essential Oils, Part XIX. The Essential Oil from the Rhizomes of *Kaempferia galanga* by Puthan Madhathil Bhaskara Panicker, B. Sanjiva Rao and John Lionel Simonsen. Part XX. The Essential Oil from the Rhizomes of *Curcuma Aromatica* Salisb., by B. Sanjiva Rao, Vishnu Purushottam Shintre and John Lionel Simonsen. Part XXI. The Essential Oil from the Wood of *Erythroxylon monogynum* Roxb., by B. Sanjiva Rao, Vishnu Purushottam Shintre and John Lionel Simonsen. Price, As. 12.
- Part VII.—Reactions of Chromates at High Temperatures, Part II. The System $\text{CaO} - \text{Cr}_2\text{O}_3 - \text{O}_2$, by K. S. Nargund and H. E. Watson. Price, Re. 1.
- Part VIII.—I. The Catalytic Hydrogenation of Carone, by Subramania Narayana Iyer and John Lionel Simonsen. II. Conessine, by Darab Dinsha Kanga, Panchandana Ramaswami Ayyar and John Lionel Simonsen. Price, As. 8.
- Part IX.—Derivatives of Naphthaquinolines and Naphthaisoquinolines. By Charles Stanley Gibson, Kalvoi Venkatakrishna Hariharan, Kottiazath Narayana Menon and John Lionel Simonsen. Price As. 12.
- Part X.—I. Isomeric Phenylserines. II. The Unstable Modification of *iso*Nitrosocamphor. III. *d*-Mannitol from *Gardenia Turgida*, by M. O. Forster and Keshaviah Aswath Narain Rao. Price, As. 12.

VOLUME IX (B)—1926.

- Part I.—Double Synchronous Speed Alternators, by K. P. Roy. Price, Re. 1.
- Part II.—The Characteristics of Beam Transmitting Aerials, by J. K. Catterson-Smith. Price, Re. 1.
- Part III.—Aerial Testing, by J. K. Catterson-Smith. Price, Rs. 1-8.
- Part IV.—Circulating Currents in Wave-Wound Armatures, by F. N. Mowdawalla and G. K. Pradhan. Price Rs. 1-8.
- Part V.—Madras (Fort) Radio Field Intensity Measurements at Bangalore, by K. Sreenivasan. Price, Rs. 3.

VOLUME X (A)—1927.

- Part I.—I. Some Reactions of Carone, by Kottiazath Narayana Menon and John Lionel Simonsen. II. A Synthesis of Morindone, by Ramkanta Bhattacharya and John Lionel Simonsen. III. Derivatives of Acenaphthpyridine. Part I, by Srikumaran Unni Nair and John Lionel Simonsen. Price, As. 12.
- Part II.—Oils and Fats from the Seeds of Indian Forest Plants. Part VIII. The Oil from the Seeds of *Thevetia Neriifolia* (Juss.). By Ramkanta Bhattacharya and P. Ramaswami Ayyar. Part IX. The Oil from the Seeds of *Cerbera Odollam* (Gaertn.). Part X. The Oil from the Seeds of *Holarrhena Antidysenterica*. Part XI. The Oil from the Seeds of *Anona Squamosa* (Linn.). By Ramachandra Vishnu Ghanekar and P. Ramaswami Ayyar. Price As. 12.