

Verified
R

Journal of the Indian Institute of

Science.

V. 10 A. Part 4. 1981/7.

Pp 43-55

SL

A, d, m 2, N 14 15

N 27. W 10. 4


196357

JOURNAL

OF THE

Indian Institute of Science.


CONTENTS.



THE CONSTITUTION OF THE ACID FORMED BY THE ACTION OF
SULPHURIC ACID ON CAMPHORQUINONE.

BY

Madhay Balaji Bhagvat and John Lionel Simonsen.



FORSTER, F.R.S., CHAIRMAN OF EDITORIAL BOARD.

JOURNAL
OF THE
INDIAN INSTITUTE OF SCIENCE

VOLUME I.—(1914-18).

Parts I.-XXI.

VOLUME II.—(1918-19).

Part I.-XV.

VOLUME III.—(1920).

Parts I.-IX.

VOLUME IV.—(1921).

Part I.—Research Notes on the Acetone Fermentation Process in India. II.—The Specific Gravities of Mixtures of Butyl Alcohol and Water. III.—A Suggested Method for the Extraction of Turpentine, Resin and Gum from the Gum-oleo-resin of *Boswellia Serrata* without the use of Solvents. IV.—Additive Compounds of Arylamines with Nitro-derivatives of Naphthalene. V.—The Manufacture of Thymol from Ajowan. VI.—Phase Transformation and Phase Balancing. VII.—Some experiments in the Use of Antiseptics in the Manufacture of Glue and Gelatine. VIII.—Studies relating to the Bacteria associated with Rice and other cereals. IX.—Preliminary Note on the Purification of Water by Activated Silt. X.—Additive Compounds of *m*-Dinitrobenzene. XI.—The Action of Methyl and Ethyl Alcohols on Esters of 2:6 Dinitro- and 2:4:6-Trinitro-benzoic acids. XII.—Studies in Alcoholysis IV. The Alcoholysis of esters of $\alpha\beta$ -unsaturated acids and of the corresponding saturated esters. XIII.—The Bio-Chemistry of the Indigenous Indigo Dye Vat. XIV.—The Development of Chemical Industries in India. XV.—Some experiments on the production of Power Alcohol and Paper Pulp from Megasse. XVI.—Sags and Tensions of Solid Copper Overhead Conductors under Indian Conditions.

VOLUME V.—(1922).

Part I.—Studies in Acidolysis: I. The Equilibrium between Acetic Acid, Trichloroacetic Acid and their Ethyl Esters; II. The Equilibrium between Acetic Acid, Trichloroacetic Acid and their Methyl Esters. II.—An Attempted Synthesis of Norpinic Acid. III.—Experiments on the influence of *s*-Trinitrobenzene on Reactions in which Arylamines are involved. IV.—Induction Motors used as Synchronous Machines. V.—The Relation Between the Iodine Values and Refractive Indices of some Hardened Vegetable Oils, Part I. VI.—Performance of Three-phase Induction Motors on Unbalanced Voltages. VII.—The Production of Barium Salts from Barytes. VIII.—The Addition of Bromine to the α - and β -Chloro- and bromocinnamic Acids and their Esters. IX.—Oil Splitting by Castor-seed Lipase, Parts V-VII. X.—Some West Coast Vegetable Oils. XI.—Utilisation of Tamarind. XII.—West Australian 'Sandalwood Oil.' XIII.—The Extraction of Caffeine. XIV.—Mechanical Tests of Some Mysore Timbers.

VOLUME VI.—(1923).

Part I.—Mohua Oil. II.—Malabar Sardine Oil. III.—Studies in Esterification. IV.—The Action of Bases on $\alpha\alpha\beta$ - and $\alpha\beta\beta$ -Tribromo- β -phenylpropionic Acids and their Esters. V.—Hongay Oil. VI.—Cashew Kernel Oil. VII.—The Fruit of *Bassia Longifolia*. VIII.—Studies in Intensive Bacterial Oxidation, I. The Oxidation of Alcohol to Acetic Acid. IX.—Contributions to the Study of the Prickly Pear Problem. X.—A Chemical Examination of Samples of Water and of Shells from the Inle Valley. XI.—The Refining of Crude Saltpetre. XII.—Enolising Action of Hydrogen Chloride on Aldehydes and Ketones. XIII.—The Conversion of Eugenol into Isoeugenol.

Price, As. 8.]

[Vol. 10A, Part IV, pp. 43 to 55.]

JOURNAL

OF THE

Indian Institute of Science.


CONTENTS.

THE CONSTITUTION OF THE ACID FORMED BY THE ACTION OF
SULPHURIC ACID ON CAMPHORQUINONE.

BY

Madhay Balaji Bhagvat and John Lionel Simonsen.

DR. M. O. FORSTER, F.R.S., CHAIRMAN OF EDITORIAL BOARD.




TL
 A. J. M. S. 1912
 1912-1914


THE CONSTITUTION OF THE ACID FORMED BY THE ACTION OF SULPHURIC ACID ON CAMPHORQUINONE¹

By Madhay Balaji Bhagvat and John Lionel Simonsen.

In a recent communication (J.C.S., 1925, 127, 1294), evidence was advanced which appeared to prove that the keto-acid, $C_{10}H_{16}O_3$, first prepared by Manasse and Samuel (*Ber.*, 1897, 30, 3157; 1902, 35, 3831) by the action of sulphuric acid on camphorquinone, was *d*-2 : 2 : 4-trimethylcyclohexan-3-one-1-carboxylic acid (I). The assignment of this constitution to the acid was based essentially on two facts: (i) on oxidation with nitric acid it gave β -methylpentane- $\beta\gamma$ -tricarboxylic acid (III), and (ii) on treatment with amyl nitrite and hydrochloric acid it did not yield an oximino-derivative, the formation of which might have been expected if the keto-acid had the alternative constitution (II) (compare footnote, *loc. cit.*, p. 1295), since in this case there is a methylene group adjacent to the carbonyl group.



In deciding between these two formulæ, owing to the symmetry of the molecules, simple degradation products were unlikely to be of much assistance, but it seemed probable that, on bromination, a dibromo-acid would result of either formula (IV) or (V). It was further anticipated that this acid would, on removal of two molecules of hydrogen bromide, tend to lose simultaneously a methyl group and so pass into a phenolic acid (VI or VII) the identification of which would have offered little difficulty.





On bromination in acetic acid, the crystalline and the liquid methyl ester of the keto-acid each gave a *dibromo-ester*, $C_{11}H_{16}O_3Br_2$.

¹ Reprinted from the *Journal of the Chemical Society*, 1927, p. 77.

which reacted in an identical manner with alkali. In some experiments, under conditions which could not be exactly defined, the dibromo-ester was obtained crystalline. Since both the pure crystalline ester and the crude liquid ester yielded the same products, they were apparently stereoisomerides.


On treatment with a solution of barium or potassium hydroxide the dibromo-ester gave a complex mixture of acids, from which, with considerable difficulty, (A) a *hydroxyketo-acid*, $C_{10}H_{14}O_4$, m. p. 151–152°, (B) a dibasic *hydroxy-acid*, $C_{10}H_{16}O_5$, m. p. 191–192°, (C) and (D) unsaturated dibasic *acids*, $C_{10}H_{14}O_4$, m. p. 257° and 215–216°, respectively, were isolated.

The formation of the hydroxyketo-acid, which also reacted as a diketone, yielding a *dioxime* and a *disemicarbazone*, would appear to indicate that the dibromo-ester has two bromine atoms attached to the same carbon atom—a fact most simply explained if the original keto-ester has formula (II), the dibromo-ester then being represented by (VIII). An ester of this formula should yield on treatment with alkali a diketone (IX); the acid obtained, however, and also its *methyl* ester, were colourless substances which gave intense colorations with ferric chloride, and the methyl ester was readily soluble in dilute potassium hydroxide solution and reprecipitated by carbon dioxide. Moreover, in alkaline solution, the acid was unstable to potassium permanganate and on treatment with acetyl chloride it yielded an *acetyl* derivative. There can be no doubt, therefore, that the hydroxyketo-acid is represented by (X) or (XI), the latter being the more probable, since the acid shows no tendency to yield a lactone. Owing to the small quantity of material available, the decision between these two formulæ could not be made.




In its general properties this acid resembles the diketones or, more correctly, the hydroxy-ketones obtained by Wallach (*Annalen*, 1918, 414, 296) in an analogous manner by the action of alkali on the dibromo-derivatives of various cyclic ketones. Wallach considered that the two bromine atoms were not attached to the same carbon atom

and that the formation of the diketone proceeded in accordance with the scheme:





Our experiments do not furnish any evidence in support of this somewhat complicated series of reactions and we prefer for the present to assume that the dibromo-ester has formula (VIII).

The hydroxyketo-ester (or acid), on digestion with a strong solution of potassium hydroxide (50 per cent. KOH), behaves in a similar manner to Wallach's hydroxy-ketones, undergoing molecular rearrangement with addition of water and becoming converted into a dibasic hydroxy-acid identical with the acid (B) that forms the main product of the action of alkali on the dibromo-ester. There can be no doubt that this acid is 1-hydroxy-4:4:5-trimethylcyclopentane-1:3-dicarboxylic acid (XII). It yields an unsaturated acid when heated above its melting point and gives an *acetyl* derivative (XIII) when digested with acetic anhydride. It is somewhat remarkable that the



acid yields neither a lactone nor an anhydride, since either the carboxyl groups or the hydroxyl and one carboxyl group must be in the *cis*-position. We are of opinion that the carboxyl groups are in the *trans*-position, because the acetyl derivative of the hydroxy-acid, when heated to its melting point, loses acetic acid and is converted quantitatively into an unsaturated acid, $C_{10}H_{14}O_4$, m. p. 257°. This acid is identical with the unsaturated acid (C) referred to on p. 44 and must be represented by formula (XIV) or (XV); there can be no doubt that it is 4:4:5-trimethyl- Δ^5 -cyclopentene-1:3-dicarboxylic acid (XIV). On oxidation with potassium permanganate, it yields a dibasic *keto-acid*, $C_9H_{14}O_5$, m. p. 125–126°, which, on further oxidation with sodium hypobromite, gives γ -methylbutane- $\alpha\beta\gamma$ -tricarboxylic acid (XVIII). The keto-acid must therefore be γ -acetyl- γ -methylbutane- $\alpha\beta$ -dicarboxylic acid (XVII), γ -Methylbutane $\alpha\beta\gamma$ -tricarboxylic acid is also formed


when 4:4:5-trimethyl- Δ^5 -cyclopentene-1:3-dicarboxylic acid is oxidised with dilute nitric acid.



When 4:4:5-trimethyl- Δ^5 -cyclopentene-1:3-dicarboxylic acid is oxidised with potassium permanganate at 0° , in addition to γ -acetyl- γ -methylbutane- $\alpha\beta$ -dicarboxylic acid a second *keto-dicarboxylic* acid, $\text{C}_{10}\text{H}_{12}\text{O}_5$, m. p. $183-185^\circ$, is formed. This acid, which was characterised by the preparation of the *oxime* and *semicarbazone*, was unstable to potassium permanganate in alkaline solution. It was unfortunately not obtained in quantity and we are unable to suggest a satisfactory formula or account for its formation. The reaction would appear to be abnormal and recalls the conversion of Δ^1 -cyclohexeneacetic acid into 1-acetyl- Δ^1 -cyclopentene by an alkaline solution of potassium permanganate (Perkin and Wallach, *Ber.*, 1909, 42, 145). On oxidation with dilute nitric acid, the keto-acid yielded γ -methylbutane- $\alpha\beta\gamma$ -tricarboxylic acid, whilst on treatment with sodium hypobromite solution, although carbon tetrabromide and bromoform separated, the product was a resin which could not be crystallised and contained an appreciable quantity of bromine.

4:4:5-Trimethyl- Δ^5 -cyclopentene-1:3-dicarboxylic acid was somewhat resistant to reducing agents, but it was converted into the saturated cyclopentane acid (XVI) by an excess of sodium and boiling amyl alcohol.


Unlike all the substances so far described, the unsaturated acid (D), m. p. $215-216^\circ$ (p. 44) was optically active, the *sodium* salt being laevorotatory. Since the acid was racemised, with formation of the saturated cyclic acid (XVI), on reduction with sodium and amyl alcohol and yielded γ -methylbutane- $\alpha\beta\gamma$ -tricarboxylic acid on oxidation with nitric acid, there can be no doubt that it is the *laevo*-form of (XIV).



The foregoing results appear to us to establish definitely that the keto-acid first described by Manasse and Samuel must be *d*-2:2:3-trimethylcyclohexan-4-one-1-carboxylic acid (II), and it became necessary


once more to attempt the preparation of an oximinoderivate. Although neither the acid nor the methyl ester could be induced to react with benzaldehyde or piperonal, the methyl ester reacted with amyl nitrite in the presence of sodium to yield an oil which was soluble in alkali and was evidently the crude *oximino-ester* (XIX). This ester could not be obtained crystalline, but on hydrolysis with formaldehyde and hydrochloric acid, followed by treatment with a dilute solution of potassium hydroxide, it gave an acid which partly crystallised on keeping. This acid, which was evidently somewhat impure 4-hydroxy-2:2:3-trimethyl- Δ^3 -cyclohexen-5-one-1-carboxylic acid (XI), was not purified, but was converted into the cyclopentene acid by the reactions outlined above. The acid melted at 257° and was identical with that prepared from the dibromo-ester. The formation of 4:4:5-trimethyl- Δ^5 -cyclopentene-1:3-dicarboxylic acid in this manner would appear to eliminate definitely and possibility of molecular rearrangement during its formation from the bromo-ester.

The suggestion (*loc. cit.*, p. 1296) that the keto-acid can exist in keto- and enol-forms has been confirmed by Mr. Aswath Narain Rao, who has prepared the *acetyl* derivative of the enolic form by treating camphorquinone with acetic anhydride and zinc chloride. This acetyl derivative is laevorotatory in chloroform solution and does not show mutarotation. It may be represented by formula (XX) or (XXI), and from analogy with the other unsaturated acids here referred to we are inclined to consider (XX) the more probable.



On hydrolysis, the acetyl derivative yields the ordinary ketonic form of the acid and all attempts to prepare it by direct acetylation have proved unsuccessful.

Since the acid previously considered to be *d*-2:2:4-trimethylcyclohexan-3-one-1-carboxylic acid has now been shown to be *d*-2:2:3-trimethylcyclohexan-4-one-1-carboxylic acid, the constitution of some of the compounds prepared from it require revision. *trans-d*-3-Hydroxy-2:2:4-trimethylcyclohexane-1-carboxylic acid should be *trans-d*-4-hydroxy-2:2:3-trimethylcyclohexane-1-carboxylic acid (XXII), and the unsaturated acid derived therefrom must be represented by formula (XXIII) or (XXIV.)



EXPERIMENTAL.

Bromination of Methyl 2 : 2 : 3-Trimethylcyclohexan-4-one-1-carboxylate. Methyl 5 : 5-Dibromo-2 : 2 : 3-trimethylcyclohexan-4-one-1-carboxylate (VIII).—To a well-cooled solution of the crystalline methyl ester (55 g.) in acetic acid (280 c.c.), bromine (91.4 g.), dissolved in acetic acid (91.4 c.c.), was added. The reaction mixture was poured on to ice and the viscid oil which separated was washed with dilute aqueous sodium carbonate solution and dried in ethereal solution. On evaporation of the ether, the *dibromo-ester* was usually obtained as a brown oil, but in one experiment this gradually crystallised in needles: these were collected, drained on porous porcelain, and recrystallised from methyl alcohol, the dibromoester thus being obtained in glistening prisms, m. p. 57–58° (Found: Br, 45.1. $C_{11}H_{16}O_3Br_2$ requires Br, 44.9).

Action of Barium Hydroxide on Methyl 5 : 5-Dibromo-2 : 2 : 3-trimethylcyclohexan-4-one-1-carboxylate.—A mixture of the crystalline (or liquid) dibromo-ester (26 g.) with a hot solution of barium hydroxide (52 g. in 170 c. c. of water) was boiled with constant agitation during one hour. The hot solution was filtered from a little insoluble matter (from which a further quantity of the mixed acids was obtained after treatment with a boiling solution of potassium hydroxide) and acidified. The liberated acids, which were isolated as a viscid brown oil (15 g.) by means of ether and partly crystallised on keeping, were esterified with methyl alcohol in the usual manner, and the esters formed were dissolved in ether. The ethereal solution was washed with dilute aqueous sodium carbonate (A), and then with 5 per cent sodium hydroxide solution (B) until a test portion no longer gave a colour with ferric chloride. The ethereal solution was then dried and the solvent evaporated (C).

The sodium carbonate extract (A) on acidification yielded a small quantity of a resinous acid which was not further examined.

Methyl 4-Hydroxy-2 : 2 : 3-trimethyl- Δ^3 -cyclohexen-5-one-1-carboxylate (formula as XI).—The deep red sodium hydroxide extract (B) was saturated with carbon dioxide and the colourless crystalline solid that separated was collected (1.4 g.); the alkaline solution was then acidified and the viscid red oil produced was added to the mixture of acids obtained in later experiments and re-esterified.

The crystalline solid separated from light petroleum in rosettes of prismatic needles, m. p. 73–74° (Found: C, 61.9; H, 7.6. $C_{11}H_{16}O_4$ requires C, 62.3; H, 7.5 per cent.). The ester was appreciably soluble in hot water and dissolved readily in all the ordinary organic solvents except light petroleum, in which it was somewhat sparingly

soluble. It gave with alcoholic ferric chloride a dirty green coloration, which was discharged by a trace of alkali.

The *dioxime*, which was obtained by heating the ester with hydroxylamine hydrochloride and sodium acetate in methyl-alcoholic solution, crystallised from benzene in fine needles, m. p. 138–139° (Found: C, 54.1; H, 7.5; N, 11.9. $C_{11}H_{16}O_4N_2$ requires C, 54.5; H, 7.4; N, 11.6 per cent.). In alcoholic solution it gave with ferric chloride a deep brown coloration. The *disemicarbazone* separated from hot water, in which it was somewhat readily soluble, in prisms, m. p. 206–237° (Found: N, 25.9. $C_{13}H_{22}O_4N_6$ requires N, 25.8 per cent.).

4-Hydroxy-2 : 2 : 3-trimethyl- Δ^3 -cyclohexen-5-one-1-carboxylic acid (XI), obtained by hydrolysing the methyl ester on the water-bath with a slight excess of dilute potassium hydroxide solution, crystallised from hot water or toluene in prismatic needle, m. p. 151–152° (Found: C, 60.8, 60.7; H, 7.4, 7.2. $C_{10}H_{14}O_4$ requires C, 60.6; H, 7.1 per cent. It was readily soluble in hot water, alcohol, acetone, or ether, but more sparingly soluble in chloroform, benzene, or toluene, and gave a dirty bluish-green coloration with aqueous or alcoholic ferric chloride. In alkaline solution it immediately decolorised potassium permanganate, but it did not react with bromine. The acid was readily oxidised by hydrogen peroxide, but the crystalline products were too small in quantity for identification. It condensed with *o*-phenylenediamine, and on treatment with phenylhydrazine it gave a mixture of the *mono*- and the *di*-phenylhydrazones as a yellow, amorphous solid, which was purified by solution in alkali and precipitation by acid (Found: N, 11.5. $C_{16}H_{20}O_3N_2$ requires N, 9.7 per cent. $C_{22}H_{26}O_2N_4$ requires N, 14.8 per cent.).

The *acetyl* derivative, prepared by digesting the acid with acetyl chloride, crystallised from water in irregular plates; these melted at 95° when air-dried and at 128° after being dried over sulphuric acid. They retained water of crystallisation ($\frac{1}{2}$ mol.) even after being kept for some months in a vacuum over phosphoric oxide (Found: C, 58.0; H, 6.7. $C_{12}H_{16}O_5 \cdot \frac{1}{2}H_2O$ requires C, 57.8; H, 6.8 per cent.).

Conversion of 4-Hydroxy-2 : 2 : 3-trimethyl- Δ^3 -cyclohexen-5-one-1-carboxylic acid into 1-Hydroxy-4 : 4 : 5-trimethylcyclopentan-1 : 3-dicarboxylic Acid (XII).—A mixture of the hydroxyketo-acid (1 mol.) and 50 per cent. potassium hydroxide solution (3 mols.) was boiled for one hour. The product, obtained in quantitative yield on cooling and acidification, melted at 191–192° after crystallisation from hot water and was identified with the following acid by the mixed melting-point method (Found: C, 55.1; H, 7.9 per cent.).

1-Hydroxy-4:4:5-trimethylcyclopentane-1:3-dicarboxylic Acid (XII).—The ester fraction (C) (p. 48) distilled almost entirely at 150–160°/10 mm. as a colourless oil, which was a mixture. The ester from one experiment (55 g.) was heated with an excess of methyl-alcoholic potassium hydroxide on the water-bath for 1 hour, and the solution kept at 0° for 12 hours. A somewhat sparingly soluble potassium salt separated in needles. This was well washed with ice-cold methyl alcohol (the filtrate, C₁, was reserved for later examination), dissolved in the minimum quantity of water, and acidified; a precipitate (16 g.) separated, and a little more was extracted from the filtrate with ether, *1-Hydroxy-4:4:5-trimethylcyclopentane-1:3-dicarboxylic acid* thus obtained crystallised from hot water in glistening needles, m. p. 191–192° after softening at 184°, which contained $\frac{1}{2}$ H₂O (Found: C, 53.2; H, 7.8; H₂O, 4.0. C₁₀H₁₆O₅, $\frac{1}{2}$ H₂O requires C, 53.3; H, 7.6; H₂O, 4.0 per cent. Found in material dried at 110°: C, 55.5; H, 7.5. C₁₀H₁₆O₅ requires C, 55.5; H, 7.4 per cent. It was sparingly soluble in benzene, chloroform, or light petroleum, readily soluble in alcohol or acetic acid, gave no coloration with ferric chloride, was stable to potassium permanganate in alkaline solution, and was optically inactive.

The acid was treated with phosphorus pentachloride (3 mols.) and the product, after removal of the phosphorus oxychloride under diminished pressure, poured into formic acid. *4:4:5-Trimethyl-Δ⁵-cyclopentene-1:3-dicarboxylic acid*, m. p. 257°, was thus obtained in poor yield; the other products of the reaction were not examined.

The *acetyl* derivative (XIII) formed when the hydroxy-acid was digested for 1 hour with an excess of acetic anhydride was obtained in needles on addition of water and concentration of the solution; it was recrystallised from hot water, in which it was somewhat sparingly soluble (Found: C, 55.7; H, 7.5. C₁₂H₁₈O₆ requires C, 55.8; H, 7.0 per cent.) It decomposed at 207–208° and immediately resolidified to a product, m. p. 257°, which was *4:4:5-trimethyl-Δ⁵-cyclopentene-1:3-dicarboxylic acid*; the elimination of acetic acid is quantitative and this is the simplest method for the preparation of the pure unsaturated acid.

4:4:5-Trimethyl-Δ⁵-cyclopentene-1:3-dicarboxylic Acid (XIV).—The methyl-alcoholic solution (C₁, above) from which the potassium salt of the hydroxy-acid had been separated was freed from alcohol on the water-bath and acidified; the crystalline precipitate was collected (7 g.) and the filtrate (C₂) reserved. By recrystallising the former from methyl alcohol, *4:4:5-trimethyl-Δ⁵-cyclopentene-1:3-dicarboxylic acid* was obtained in glistening rhombohedra, m. p. 257–258° (Found: C, 60.5; H, 7.2; M, 197.8. C₁₀H₁₄O₄ requires C, 60.6; H, 7.1 per cent.; M, 198). It was very sparingly soluble in water, ether, benzene,

or chloroform, and more readily soluble in hot methyl alcohol, acetic acid, or formic acid. It immediately decolorised an alkaline solution of potassium permanganate, but reacted neither with bromine in acetic acid, solution nor with sodium amalgam at the ordinary temperature. When heated above its melting point, it did not yield an anhydride, but sublimed in long needles with very little decomposition. The aqueous solution of the sodium salt was optically inactive. The *dianilide*, prepared from the acid chloride in the usual manner, crystallised from dilute alcohol in fine needles, m. p. 200–201° (Found: C, 75.7; H, 7.4. C₂₂H₂₄O₂N₂ requires C, 75.9; H, 6.9 per cent.)

4:4:5-Trimethylcyclopentane-1:3-dicarboxylic Acid (XVI) was obtained in an excellent yield when the unsaturated acid was reduced with sodium and amyl alcohol. It separated from hot water in soft, glistening prisms, m. p. 182–183°, and showed no tendency to yield an anhydride, small quantities distilling unchanged. It would appear, therefore, to be the *trans*-acid. It was readily soluble in hot water, alcohol, or ethyl acetate, and very sparingly soluble in benzene or chloroform (Found: C, 59.7; H, 8.3. C₁₀H₁₆O₄ requires C, 60.0; H, 8.0 per cent.)

1:4:4:5-Trimethyl-Δ⁵-cyclopentene-1:3-dicarboxylic Acid.—The filtrate (C₂), on extraction with ether, yielded a viscid yellow oil which could not be purified. Since it apparently contained a quantity of the hydroxycyclopentanedicarboxylic acid, it was digested with acetic anhydride and the crude acetyl derivative was heated at 200–210° until all evolution of acetic acid had ceased. The melt was dissolved in sodium carbonate solution, filtered from a little tar, and acidified. The solid which separated was collected, the filtrate (C₃) being reserved, and recrystallised from methyl alcohol, a further quantity of the cyclopentene acid, m. p. 257°, being obtained. The methyl-alcoholic filtrate was evaporated, and the residue repeatedly crystallised from formic acid; *1:4:4:5-trimethyl-Δ⁵-cyclopentene-1:3-dicarboxylic acid* was thus obtained in irregular plates, m. p. 215–216° (Found: C, 60.5; H, 7.5. C₁₀H₁₄O₄ requires C, 60.6; H, 7.1 per cent.) This acid was very much more readily soluble in all the ordinary organic solvents than its inactive isomeride. On reduction with sodium and amyl alcohol, it yielded the inactive cyclopentane acid (XVI). Under the conditions used for the oxidation of the racemic acid with nitric acid (see below), it gave *γ*-methylbutane-*αβγ*-tricarboxylic acid, decomp. 154–155°, thus leaving no doubt as to its constitution. The sodium salt was levorotatory in aqueous solution; $[\alpha]_D^{25} - 62^\circ$.

The *dianilide* crystallised from dilute alcohol in glistening cubes, m. p. 208–209° (Found: N, 8.4. C₂₂H₂₄O₂N₂ requires N, 8.0 per cent.)

51

196357

196357

The filtrate (C_3 , above) gave on extraction with ether a viscid oil which on trituration with formic acid yielded a further small quantity of the *l*-unsaturated acid. The formic acid-mother-liquor contained an oil which was not examined.

Oxidation of 4 : 4 : 5-Trimethyl- Δ^5 -cyclopentene-1 : 3-dicarboxylic Acid.—(i) *With nitric acid.* The unsaturated acid (2 g.) was heated with nitric acid (d 1.2; 35 c.c.) on a water-bath for 8 hours and on a sand-bath for 8 hours; nitric acid (d 1.4; 15 c.c.) was then added and the heating continued for a further 8 hours, a clear solution being obtained. The excess of mineral acid was removed on the water bath; the viscid oily residue, on trituration with a little hydrochloric acid, crystallised almost completely. After draining on porous porcelain, it was recrystallised from a little hydrochloric acid, γ -methylbutane- $\beta\gamma$ -tricarboxylic acid separating in rosettes of needles, decomp. 154–155° (Found: C, 47.2; H, 5.8; M , 205. Calc.: C, 47.1; H, 5.9 per cent; M , 204). The identity of this acid was confirmed by the preparation of the anhydride, m.p. 142°.

(ii) *With potassium permanganate.* To a solution, at 0–5° of the cyclopentene acid (5 g.) in aqueous sodium carbonate through which a current of carbon dioxide was passing, 2.5 per cent. potassium permanganate solution (equiv. to 1 O_2) was added; the colour then remained permanent for 5 minutes. The filtered solution was concentrated in a current of carbon dioxide, acidified, and repeatedly extracted with ether and the extracts were dried and evaporated; a viscid brown oil remained which partly crystallised. This was triturated with concentrated hydrochloric acid, which removed a readily soluble brown oil (Z), and the residue was drained on porous porcelain and recrystallised from hydrochloric acid or from acetone-benzene; a ketodicarboxylic acid was thus obtained in glistening prisms, m.p. 182–183°, softening at 177° (Found: C, 56.4, 56.9; H, 6.0, 5.8; M , 212.6. $C_{10}H_{12}O_5$ requires C, 56.6 H, 5.6, per cent.; M , 212). This acid was readily soluble in water, alcohol, ethyl acetate, or acetone, and more sparingly soluble in benzene or chloroform. It was slowly attacked by hot chromic acid solution, did not absorb bromine in acetic acid solution, and rapidly decolorised an alkaline solution of potassium permanganate at room temperature. It was readily oxidised by sodium hypobromite with formation of bromoform and carbon tetrabromide; the acid resulting from the oxidation was a gum which contained bromine and was not further examined. The salts of the keto-acid are readily soluble in water.

The *oxime* crystallised in small plates, decomp. 214°, containing apparently $\frac{1}{2}H_2O$ which it did not lose at 100° (Found: C, 50.6, 50.7; H, 5.8, 6.2. $C_{10}H_{13}O_3N, \frac{1}{2}H_2O$ requires C, 50.8; H, 5.9 per cent.).

The *semicarbazone* crystallised from dilute alcohol in microscopic prisms, decomp. 241° (Found: C, 48.8; H, 5.9. $C_{11}H_{15}O_3N_3$ requires C, 49.1; H, 5.6 per cent.

The keto-acid, on oxidation with nitric acid (d 1.2), gave an excellent yield of γ -methylbutane- $\alpha\beta\gamma$ -tricarboxylic acid.

γ -Acetyl- γ -methylbutane- $\alpha\beta$ -dicarboxylic Acid (XVII).—The hydrochloric acid solution (Z, above) gave, after the removal of the solvent in a vacuum over potassium hydroxide, a viscid brown oil. This (7 g.) was oxidised in sodium carbonate solution with potassium permanganate at 15–20°. Repeated extraction of the filtered, concentrated, and acidified solution with ether yielded an oil which partly crystallised. The solid was collected and recrystallised from dry ether, in which it was only sparingly soluble, γ -acetyl- γ -methylbutane- $\alpha\beta$ -dicarboxylic acid separating in fine, glistening prisms, m. p. 125–126°. This acid was readily soluble in water or alcohol, but sparingly soluble in benzene or light petroleum. It did not react with phenylhydrazine or semicarbazide (Found: C, 53.1; H, 7.2. $C_9H_{14}O_5$ requires C, 53.5, H, 6.9 per cent.).

When a solution of the sodium salt of the keto-acid was treated with the calculated quantity of a solution of sodium hypobromite, bromoform immediately separated. After 1 hour, this was removed and the alkaline solution was acidified and extracted with ether; the oil obtained on removal of the solvent immediately crystallised and was identified as γ -methylbutane- $\alpha\beta\gamma$ -tricarboxylic acid.

Methyl 5-oximino-2 : 2 : 3-trimethylcyclohexan-4-one-1-carboxylate (XIX).—Methyl *d*-2 : 2 : 3-trimethylcyclohexan-4-one-1-carboxylate (0.5 g.) was dissolved in ether containing finely divided sodium (0.06 g.), and to the well-cooled mixture *isoamyl* nitrite (0.26 g.) in ether was added. After the mixture had been kept at 0° for 12 hours the sodium had completely dissolved and a flocculent, yellow precipitate had separated. Ice was then added and the yellow, aqueous solution was separated and acidified. The brown oil obtained (0.3 g.), which showed no tendency to crystallise, was analysed after being kept for some days in a vacuum over sulphuric acid. The first and the second analysis refer to specimens prepared from the crystalline and the liquid ester, respectively (Found: N, 6.5, 6.7. $C_{11}H_{17}O_4N$ requires N, 6.2 per cent.) The *oximino-ester* was soluble in all the ordinary organic solvents and in a solution of potassium hydroxide, but was insoluble in aqueous sodium carbonate; it gave with alcoholic ferric chloride a deep brown coloration.

The ester¹ was hydrolysed by treatment with formaldehyde and hydrochloric acid, followed by dilute aqueous potassium hydroxide; the viscous oil obtained gave with ferric chloride a dirty bluish-green coloration. On keeping, the oil partly crystallised, but, as the quantity of material available was small, no attempt was made at purification and the oil was converted directly into 4:4:5-trimethyl- Δ^5 -cyclopentene-1:3-dicarboxylic acid by the methods already described. The cyclopentene acid prepared in this manner crystallised from methyl alcohol in the characteristic rhombohedra, m. p. 257–258°, and was identical in every way with the acid prepared from the dibromo-ester.

1-4-Acetyl-2:2:3-trimethyl- Δ^3 -cyclohexene-1-carboxylic Acid (XX [with ASWATH NARAIN RAO].—A mixture of camphorquinone (25 g.) acetic anhydride (75 c.c.), and freshly-fused zinc chloride (5 g.), was heated on a sand-bath, a vigorous reaction taking place with effervescence and evolution of heat. As soon as the reaction started, the source of heat was removed and the dark brown solution was allowed to cool, diluted with water (150 c.c.), and warmed on the water-bath for a short time to decompose the acetic anhydride. After 12 hours, the pale yellow precipitate of needles and tar was collected and washed with water (yield, 25.5 g.). The filtrate, on neutralisation of the excess of acetic acid with solid sodium carbonate, deposited a further quantity of solid (1 g.). The crude product was repeatedly extracted with dilute sodium carbonate solution until a test portion no longer gave a precipitate on acidification. The residue was extracted with dilute sodium hydroxide solution, which left undissolved camphorquinone (5.4 g.) and a little tar. The sodium hydroxide solution, on acidification, deposited a resinous mass which could not be purified. The deep brown sodium carbonate extract was carefully acidified with dilute sulphuric acid; after the addition of a small quantity of acid, the solution was extracted with ether to remove resinous impurities; on further addition of acid, an oil was deposited which, on evaporation of the dissolved ether, solidified. The crude acid (13.2 g.) was crystallised from hot water, dilute alcohol, or, preferably, light petroleum (b.p. 60–80°), 1-4-acetyl-2:2:3-trimethyl- Δ^3 -cyclohexene-1-carboxylic acid separating in colourless, transparent, glistening laminae, m. p. 109–111° (Found: C, 63.2; H, 8.3; *M*, 224. $C_{12}H_{18}O_4$ requires C, 63.7; H, 8.0 per cent.; *M*, 226). In chloroform solution (1.283 g. in 25 c.c.), it showed $[\alpha]_D^{25} -66.3^\circ$, and this value did not alter during 48 hours. In carbon tetrachloride solution the acid absorbed an equivalent weight of bromine, hydrogen bromide being evolved, and on evaporation of the solvent a sharp-smelling oil remained. In

¹ During the hydrolysis with potassium hydroxide amyl alcohol separated, indicating that 'alcoholysis' with partial formation of the amyl ester had taken place during the reaction with amyl nitrite.


alkaline solution, the acid was unstable to potassium permanganate; the products of the reaction could not be identified.

A solution of the acid (1 g.) in 15 per cent sodium hydroxide solution (8 c.c.) was heated on the water-bath for three hours. On acidification of the cooled solution, an oil was precipitated which partly solidified over-night. The solid was separated, drained on porous porcelain, and recrystallised from water; it was then obtained in needles, m. p. 69–70°. The anhydrous acid melted at 97–98°. It yielded a semicarbazone, m. p. 227–228°, and was identified as 2:2:3-trimethylcyclohexan-1-one-1-carboxylic acid by direct comparison with an authentic specimen. The oil extracted from the porous porcelain on which the keto-acid had been drained gave no colour with ferric chloride and evidently contained no hydroxy-acid. The enolic form, therefore, would appear to be incapable of existence. Attempts were made to acetylate the keto-acid directly, but in all cases the acid was recovered unchanged.

The authors wish to thank Dr. M. O. Forster, F. R. S., for much kindly criticism and also for permission to publish the experiments on 1-4-acetyl-2:2:3-trimethyl- Δ^3 -cyclohexene-1-carboxylic acid, which were carried out under his direction.

Department of Organic Chemistry,
Indian Institute of Science,
Bangalore.

[Accepted, 10-3-27.]



RECEIVED
MAY 10 1927

VOLUME VII.—(1924).

Part I.—Studies in Alcoholysis V. The Alcoholysis of Esters of $\alpha\beta$ -unsaturated Acids and of the corresponding saturated Esters. II.—Chemical Factors in Denitrification. III.—The Retting of Coconut Husk for the Production of Coir. IV.—Reactions of Chromates at High Temperatures. Part I. The Synthesis and Decomposition of Calcium, Sodium and Magnesium Chromates in Air. V.—Oil from *Mimusops Hexandra*: Rayan Oil. VI.—The Relation Between the Iodine Values and Refractive Indices of Hardened Oils. Part II. VII.—Contributions to the Scientific Study of the Lac Industry. Parts I—VIII. VIII.—I. The Systematic Nomenclature of Polycyclic Carbon Systems. II. Polycyclic and Cage Systems of Carbon Compounds. III. The Systematic Nomenclature of Heterocyclic Compounds including Polycyclic Structures. IX.—The Use of Mixed Catalysts in the Hydrogenation of Oils. X.—Condensation of Aromatic Amines with Chloroform or Carbon Tetrachloride in the Presence of Finely Divided Copper. XI.—I. Notes on Bixin, II. An Alkaloid from *Anona Squamosa* Leaves. XII.—Iodine as a Catalyst in Reactions Involving Elimination of Hydrogen Halides. XIII.—Studies Relating to the Symbiosis of Seeds and Bacteria. XIV.—The Bio-Genesis of Mahua Oil. XV.—Contributions to the Scientific Study of the Lac Industry. Parts IX—X.

VOLUME VIII (A)—(1925).

Part I.—The Photochemical Oxidation of Aromatic Hydrocarbons. I. II.—Kachi-Grass Oil. III.—Argemone Oil. IV.—Essential Oil of *Cyperus Rotundus*. V.—Dehydration of Rectified Spirit by means of Anhydrous Calcium Chloride. VI.—The Hydrolysis of the Amides of $\alpha\beta$ -Unsaturated Acids and of their Saturated Analogues. VII.—I. Studies Relating to the Acetone-Producing Organisms. II. Mahua Flowers as Raw Material for the Acetone-Fermentation Process. VIII.—Studies in Esterification. IX.—Constituents of the Marking-Nut: '*Semecarpus Anacardium*' Linn. X.—Notes on Some Indian Essential Oils. XI.—Bromo-Derivatives of Para-Methoxycinnamic Acid. XII.—I. An Examination of Some Gum-Enzymes. II. Chemical Constitution of the Gum from '*Boswellia Serrata*'. XIII.—Industrial Wastes as Manure. I. Ajowan and Mohua Cakes as Fertilisers. II. Utilisation of Refuse. XIV.—Constitution of Manasse's Hydroxycamphor. XV.—The Reaction Between Sodium Sulphite and Sulphur. XVI.—The Occurrence of Sylvestrene.

VOLUME VIII (B)—1925.

Part I.—Lecher Wire Measurements. II.—The Aluminium Anode Film Dielectric.

VOLUME IX (A)—1926.

Part I.—Contributions to the Scientific Study of the Lac Industry. Part XI. Early Recognition of Sex among Lac Insects, by S. Mahdibassan. Price, Rs. 4.
Part II.—Vegetable Oils Containing Glycerides of Erucic Acid by J. J. Sudborough, H. E. Watson and P. Ramaswami Ayyar. Price, Rs. 2-8.
Part III.—The Production of Ether by Solid Catalysts, by S. K. Kulkarni Jatkar and H. E. Watson. Price Rs. 3.
Part IV.—The Constituents of Some Indian Essential Oils. Part XVIII. Derivatives of Abietic Acid, by Madyar Gopal Rau and John Lionel Simonsen. Price, As. 8.

PRINTED AND PUBLISHED
BY GEORGE KENNETH AT THE DIOCESAN PRESS,
POST BOX 455, MADRAS—1927. C2807

Part V.—The Fat from 'Salvadora Oleoides': Khakan Fat, by C. K. Patel, S. Narayana Iyer, J. J. Sudborough and H. E. Watson. Price, As. 12.

Part VI.—The Constituents of Some Indian Essential Oils, Part XIX. The Essential Oil from the Rhizomes of *Kaempferia galanga* by Puthan Madhathil Bhaskara Panicker, B. Sanjiva Rao and John Lionel Simonsen. Part XX. The Essential Oil from the Rhizomes of *Curcuma Aromatica* Salisb., by B. Sanjiva Rao, Vishnu Purushottam Shintre and John Lionel Simonsen. Part XXI. The Essential Oil from the Wood of *Erythroxylon monogynum* Roxb., by B. Sanjiva Rao, Vishnu Purushottam Shintre and John Lionel Simonsen. Price, As. 12.

Part VII.—Reactions of Chromates at High Temperatures, Part II. The System $\text{CaO} - \text{Cr}_2\text{O}_3 - \text{O}_2$, by K. S. Nargund and H. E. Watson. Price, Re. 1.

Part VIII.—I. The Catalytic Hydrogenation of Carone, by Subramania Narayana Iyer and John Lionel Simonsen. II. Conessine, by Darab Dinsha Kanga, Panchandana Ramaswami Ayyar and John Lionel Simonsen. Price, As. 8.

Part IX.—Derivatives of Naphthoquinolines and Naphthalisoquinolines. By Charles Stanley Gibson, Kalvo Venkatakrishna Hariharan, Kottiazath Narayana Menon and John Lionel Simonsen. Price As. 12.

Part X.—I. Isomeric Phenylserines. II. The Unstable Modification of *iso*Nitrosocamphor. III. *d*-Mannitol from *Gardenia Turgida*, by M. O. Forster and Keshaviah Aswath Narain Rao. Price, As. 12.

VOLUME IX (B)—1926.

Part I.—Double Synchronous Speed Alternators, by K. P. Roy. Price, Re. 1

Part II.—The Characteristics of Beam Transmitting Aerials, by J. K. Catterson-Smith. Price, Re. 1.

Part III.—Aerial Testing, by J. K. Catterson-Smith. Price, Rs. 1-8.

Part IV.—Circulating Currents in Wave-Wound Armatures, by F. N. Mowdawalla and G. K. Pradhan. Price, Rs. 1-8.

Part V.—Madras (Fort) Radio Field Intensity Measurements at Bangalore, by K. Sreenivasan. Price, Rs. 3.

VOLUME X (A)—1927.

Part I.—I. Some Reactions of Carone, by Kottiazath Narayana Menon and John Lionel Simonsen. II. A Synthesis of Morindone, by Ramkanta Bhattacharya and John Lionel Simonsen. III. Derivatives of Acenaphthpyridine. Part I, by Srikumaran Unni Nair and John Lionel Simonsen. Price, As. 12.

Part II.—Oils and Fats from the Seeds of Indian Forest Plants. Part VIII. The Oil from the Seeds of *Thevetia Neriifolia* (Juss.) By Ramkanta Bhattacharya and P. Ramaswami Ayyar. Part IX. The Oil from the Seeds of *Cerbera Odollam* (Gaertn.). Part X. The Oil from the Seeds of *Holarrhena Antidysenterica*. Part XI. The Oil from the Seeds of *Anona Squamosa* (Linn.). By Ramachandra Vishnu Gharekar and P. Ramaswami Ayyar. Price As. 12.

Part III.—Influence of the Sulphur Atom on Optical Rotatory Power. By P. P. Shukla. Price, As. 8.

Part IV.—The Constitution of the Acid formed by the Action of Sulphuric Acid on Camphorquinone. By Madhay Balaji Bhagvat and John Lionel Simonsen. Price, As. 8.