

verified
m

Journal of the Indian Institute of
Science

Vol A Part 5 1927.

Pp 57-63

SL /

A, d, m2, N1413

N27.10A.5

196358

49

Price, As. 8.]

[Vol. 10A, Part V, pp. 57 to 63.]


JOURNAL
OF THE
Indian Institute of Science.

CONTENTS.

isoERUCIC ACID.

BY

Tahilram Jethanand Mirchandani and John Lionel Simonsen.



O. FORSTER, F.R.S., CHAIRMAN OF EDITORIAL BOARD.

JOURNAL
OF THE
INDIAN INSTITUTE OF SCIENCE

VOLUME I.—(1914-18).

Parts I.-XXI.

VOLUME II.—(1918-19).

Parts I.-XV.

VOLUME III.—(1920).

Parts I.-IX.

VOLUME IV.—(1921).

Parts I.-XVI.

VOLUME V.—(1922).

Part I.—Studies in Acidolysis: I. The Equilibrium between Acetic Acid, Trichloroacetic Acid and their Ethyl Esters; II. The Equilibrium between Acetic Acid, Trichloroacetic Acid and their Methyl Esters. II.—An Attempted Synthesis of Norpinic Acid. III.—Experiments on the influence of *s*-Trinitrobenzene on Reactions in which Arylamines are involved. IV.—Induction Motors used as Synchronous Machines. V.—The Relation Between the Iodine Values and Refractive Indices of some Hardened Vegetable Oils, Part I. VI.—Performance of Three-phase Induction Motors on Unbalanced Voltages. VII.—The Production of Barium Salts from Barytes. VIII.—The Addition of Bromine to the α - and β -Chloro- and bromocinnamic Acids and their Esters. IX.—Oil Splitting by Castor-seed Lipase, Parts V—VII. X.—Some West Coast Vegetable Oils. XI.—Utilisation of Tamarind. XII.—West Australian 'Sandalwood Oil.' XIII.—The Extraction of Caffeine. XIV.—Mechanical Tests of Some Mysore Timbers.

VOLUME VI.—(1923).

Part I.—Mohua Oil. II.—Malabar Sardine Oil. III.—Studies in Esterification. IV.—The Action of Bases on $\alpha\alpha\beta$ - and $\alpha\beta\beta$ -Tribromo- β -phenylpropionic Acids and their Esters. V.—Hongay Oil. VI.—Cashew Kernel Oil. VII.—The Fruit of *Bassia Longifolia*. VIII.—Studies in Intensive Bacterial Oxidation, I. The Oxidation of Alcohol to Acetic Acid. IX.—Contributions to the Study of the Prickly Pear Problem. X.—A Chemical Examination of Samples of Water and of Shells from the Inle Valley. XI.—The Refining of Crude Saltpetre. XII. Enolising Action of Hydrogen Chloride on Aldehydes and Ketones. XIII.—The Conversion of Eugenol into Isoeugenol.

VOLUME VII.—(1924).

Part I.—Studies in Alcoholysis V. The Alcoholysis of Esters of $\alpha\beta$ -unsaturated Acids and of the corresponding saturated Esters. II.—Chemical Factors in Denitrification. III.—The Retting of Coconut Husk for the Production of Coir. IV.—Reactions of Chromates at High Temperatures. Part I. The Synthesis and Decomposition of Calcium, Sodium and Magnesium Chromates in Air. V.—Oil from *Mimusops Hexandra*: Rayan Oil. VI.—The Relation Between the Iodine Values and Refractive Indices of Hardened Oils. Part II. VII.—Contributions to the Scientific Study of the Lac Industry. Parts I—VIII. VIII.—I. The Systematic Nomenclature of Polycyclic Carbon Systems. II. Polycyclic and Cage Systems of Carbon Compounds. III. The Systematic Nomenclature of Heterocyclic Compounds including Polycyclic Structures. IX.—The Use of Mixed Catalysts in the Hydrogenation of Oils. X.—Condensation of Aromatic Amines with Chloroform or Carbon Tetrachloride in the Presence of Finely Divided Copper. XI.—I. Notes on Bixin, II. An Alkaloid from *Anona*

Price, As. 8.]

[Vol. 10A, Part V, pp. 57 to 63.]

JOURNAL

OF THE

Indian Institute of Science.

CONTENTS.

*iso*ERUCIC ACID.

BY

Tahilram Jethanand Mirchandani and John Lionel Simonsen.




DR. M. O. FORSTER, F.R.S., CHAIRMAN OF EDITORIAL BOARD.


*iso*ERUCIC ACID.¹

By Tahirram Jethanand Mirchandani and John Lionel Simonsen.

*iso*Erucic acid was first obtained by Alexandrov and Saytzev (*J. pr. Chem.*, 1894, **49**, 58) by treating iodobehenic acid, prepared from erucic acid, with potassium hydroxide. Saytzev (*ibid.*, 1894, **50**, 65) showed that it also resulted when brassidic acid was converted into iodobehenic acid and the halogen acid treated with alkali. *iso*Erucic acid, m.p. 54–56°, was characterised by the preparation of a dibromobehenic acid, m.p. 44–46°, and by oxidation in alkaline solution with potassium permanganate to a dihydroxybehenic acid, m.p. 86–88°. At the time of these investigations, erucic acid was considered to have the constitution (I), and the above-mentioned authors regarded it as probable that *iso*erucic acid was the structural isomeride (II).



It was subsequently established that erucic acid and brassidic acid were the *cis*- and *trans*-modifications respectively of an acid (III), since on oxidation with ozone, followed by chromic acid, they both yielded brassylic and pelargonic acids (compare Noller and Adams, *J. Amer. Chem. Soc.*, 1926, **48**, 1074; Jegorov, *J. pr. Chem.*, 1912, **86**, 539). The view of Alexandrov and Saytzev (*loc. cit.*) being accepted, it followed that *iso*erucic acid would in all probability have either formula (VI) or (VII), depending on whether iodobehenic acid from erucic and brassidic acids had formula (IV) or (V).



It was shown, however, by Ponzio (*Gazzetta*, 1904, **34**, ii, 50) that when *iso*erucic acid was oxidised by concentrated nitric acid, in a manner similar to that used in the oxidation of erucic acid (*J. pr. Chem.*, 1893, **48**, 323), it gave a mixture of acids from which brassylic and pelargonic acids were isolated and he showed further that the

¹ Reprinted from the *Journal of the Chemical Society*, 1927.

dibromo-acid (m.p. 44–46°), on treatment with alkali, yielded a behenolic acid identical with that obtained under the same conditions from the dibromo-acids derived from erucic and brassidic acids. As a result of these experiments he very legitimately concluded that *isoerucic* acid was not a structural isomere of erucic and brassidic acids. In subsequent years, a number of investigations were carried out with the object of determining by physico-chemical methods the relationship of these three acids (Mascarelli and Toschi, *Gazzetta*, 1915, **45**, i, 314; Mascarelli and Sanna, *ibid.*, 1915, **45**, ii, 208, 315; Mascarelli, *ibid.*, 1917, **47**, i, 160). The conclusion was reached that in all probability brassidic and *isoerucic* acids were dimorphic forms of the *trans*-acid (III), but all attempts to convert one acid into the other failed.

Notwithstanding Ponzio's experiments, it appeared to us possible that the original view of Alexandrov and Saytzev might be correct, since the oxidation of an unsaturated fatty acid with concentrated nitric acid, owing to the extreme violence with which it proceeds, cannot be regarded as an entirely satisfactory method of orienting a double linking, whilst the conversion of the dibromobehenic acid into the acetylenic acid is carried out under conditions which are recognised as favourable to isomeric change.

Repetition of the oxidation of *isoerucic* acid with nitric acid has confirmed the formation of brassylic acid as one of the products of the reaction, but the yield of this dibasic acid is extremely small, the main substances formed being acids of lower molecular weight. We decided, therefore, to oxidise *isoerucic* acid, in the form of its ester, with ozone and also with potassium permanganate under the standard conditions used by Armstrong and Hilditch (*J. Soc. Chem. Ind.*, 1925, **44**, 43 T) for the oxidation of methyl oleate.

The *isoerucic* acid used in these experiments was prepared by the method of Alexandrov and Saytzev (*loc. cit.*) and purified by repeated crystallisation from alcohol (95%). It melted sharply at 55–57° and appeared to be homogeneous. Treatment of the *ethyl* ester with ozone gave, in addition to the main oxidation products, which will be discussed later, a not inconsiderable quantity of *ethyl* arachidate. There is no doubt that the *isoerucic* acid used in the earlier experiments was contaminated with arachidic acid, although the acid had the correct m.p. and crystallised extremely well. In our later experiments, the *isoerucic* acid, after crystallisation from alcohol to constant melting point, was further purified by fractional precipitation of the zinc salt, and any arachidic acid was thus removed in the least soluble fractions.

When *ethyl isoerucate* was oxidised with ozone, under conditions which, with erucic acid, were found to yield a readily separable

mixture of brassylic and pelargonic acids (see p. 60), the oxidation product was a mixture which was somewhat difficult to resolve. As the quantity of this material was limited, we examined only the mixture of dibasic acids formed. We were able to isolate (a) decane-*α*-dicarboxylic acid, $\text{HO}_2\text{C} \cdot [\text{CH}_2]_{10} \cdot \text{CO}_2\text{H}$, m.p. 126–127°, identified by comparison with a synthetic specimen of the acid (Walker and Lumsden, *J.*, 1901, **79**, 1201), and (b) dodecane-*α*-dicarboxylic acid, $\text{HO}_2\text{C} \cdot [\text{CH}_2]_{12} \cdot \text{CO}_2\text{H}$, m.p. 121–122°, identified by analysis and by preparation of the methyl ester, m.p. 41°. 196358

This result of the oxidation of *ethyl isoerucate* with ozone was surprising, and appeared to show that *isoerucic* acid, in spite of its sharp m.p. and well-defined crystalline form, was not homogeneous but was a mixture of the two unsaturated acids (VI) and (VII).

To confirm this conclusion, a quantity of very carefully purified methyl *isoerucate* was oxidised in acetone solution with potassium permanganate. From the oxidation products we were able to separate the two dibasic acids mentioned above and also *n*-octoic acid. Unfortunately, *n*-decoic acid, which must also have been formed, could not be obtained in a pure state.

This experiment confirmed in all respects the results obtained by oxidation with ozone and seems to leave no doubt that *isoerucic* acid is not a homogeneous substance, but is a mixture of Δ^{11} -docosene-1-carboxylic acid and Δ^{10} -docosene-1-carboxylic acid. From the yields of the oxidation acids the two acids would appear to be present in approximately equal amounts.

Since *isoerucic* acid itself was an inseparable mixture, we examined a number of derivatives to see whether any of them showed signs of heterogeneity. The *methyl* ester is crystalline, m.p. 24–26°, resolidifying at 22°; the *anilide* and *p*-toluidide both crystallise extremely well and both before and after crystallisation have the same m.p.; the *zinc* salt appears to be homogeneous.

As has already been mentioned, *isoerucic* acid is stated to yield a dibromo-derivative, m.p. 42–44°, or 46–46.6° according to Macbeth and Stewart (*Proc. Chem. Soc.* 1912, **28**, 68). We have repeated the preparation of this substance several times, but in every case the bromination product had the appearance of a mixture and the m.p. was not sharp. We have not tried the oxidation of *isoerucic* acid to dihydroxybehenic acid, but it is probable that this is a mixture of two dihydroxy-acids.

The fact that *isoerucic* acid is an inseparable mixture of two acids yielding apparently homogeneous derivatives is somewhat unusual, but is not unique. Attention may be directed to the following

TL
Ard. m. 2, 1914
1973. 150.5

somewhat similar cases: mononitro-*o*-veratraldehyde, m.p. 88° (Perkin, Roberts, and Robinson, *J. Chem. Soc.*, 1914, **105**, 2389), has been shown to be a mixture of two isomeric nitroveratraldehydes of m.p. 110° and 113° respectively (Perkin, Robinson, and Stoye, *ibid.*, 1924, **125**, 2355); anisic and veratric acids form a mixture, m.p. 145°, inseparable by crystallisation (Cain and Simonsen, *ibid.*, 1912, **101**, 1061); the geometrical isomerides of camphorquinoneoxime (*isonitrosocamphor*) are inseparable by solvents and produce a mixture melting higher than the more fusible modification (Forster, *ibid.*, 1903, **83**, 535); benzoic and *p*-nitrobenzoic acids in the proportion 2 : 1 form a mixture indivisible by solvents and melting at 182–198° (Forster and Saville, *ibid.*, 1922, **121**, 823).

During our preliminary experiments we investigated the oxidation of erucic and brassidic acids under various conditions and found the nature of the dihydroxy-acid formed to be dependent on the conditions used. In view of the recent publication of Hilditch (*ibid.*, 1926, 1828) we do not intend to continue this side of the investigation. Experiments on the synthesis of the two unsaturated acids (VI and VII) are in progress and will, it is hoped, form the subject of a future communication.

EXPERIMENTAL.

*iso*Erucic acid prepared by Alexandrov and Saytzev's method (*loc. cit.*), after four crystallisations from alcohol, melted at 54–56°, but, as has already been mentioned, it was found as the result of oxidation experiments to contain a small percentage of arachidic acid. The acid was purified by fractional precipitation of the zinc salt; eight fractions were obtained, of which the first two were rejected, since they contained arachidic acid. After this purification, the acid had m.p. 55–57° and n_D^{20} 1.4469.¹

*iso*Erucic acid is not affected by treatment with nitrous acid under conditions which convert erucic acid into brassidic acid, nor does it undergo any change when an alcoholic solution containing a trace of iodine is exposed to bright sunlight for some days.

The *zinc* salt crystallised from benzene in thin leaflets, m.p. 109°. No depression of the m.p. occurs on admixture with an equal amount of the zinc salt of brassidic acid (m.p. 112–113°), but the crystalline forms of the two salts are different, the latter salt crystallising in needles.

¹ The refractive indices of erucic and brassidic acids at the same temperature were 1.4480 and 1.4472, respectively

The *anilide* separated from alcohol in fine needles, m.p. 77–78.5° (Found: N, 3.6. $C_{28}H_{47}ON$ requires N, 3.4 per cent.), and the *p*-toluidide in needles, m.p. 85–86° (Found: N, 3.5. $C_{29}H_{49}ON$ requires N, 3.3 per cent.).

The *methyl* ester, prepared by treating the acid with methyl alcohol and sulphuric acid, crystallised in needles, b.p. 240°/14 mm., m.p. about 24–26°, resolidifying at 22° (thermometer in liquid). It is somewhat sparingly soluble in methyl alcohol (Found: *M*, 351.1; iod. val., 71.9. $C_{23}H_{44}O_2$ requires *M*, 352; iod. val., 72). The *ethyl* ester is a viscid oil, b.p. 243°/8 mm.

Oxidation of Methyl isoErucate with Potassium Permanganate.—Methyl *iso*erucate (50 g.) was dissolved in acetone (500 c.c.), and potassium permanganate (168 g.) gradually added. The oxidation proceeded vigorously and sufficient heat was generated to keep the acetone boiling. The reaction mixture was heated on the waterbath for 15 hours, the acetone distilled off, and the manganese dioxide sludge dissolved by the addition of dilute sulphuric acid and sodium bisulphite. The oil which had separated was dissolved in ether, and the ethereal extract was washed with sodium carbonate solution until no further acid was removed, dried, and evaporated (A). The sodium carbonate extract was mixed with an excess of sodium hydroxide solution, boiled for one hour, and acidified. The liberated organic acids were dissolved in ether; on removal of the solvent a semi-solid, crystalline mass remained (47 g.). The mixture of acids was digested with light petroleum (b.p. 40–60°), and the boiling solution filtered (residue B), a small quantity of solid which separated on cooling being added to B. The light-petroleum extract was evaporated and the liquid acids (20 g.) were distilled; a large fraction was then obtained, b.p. 230–270°/685 mm. The residue in the distilling flask (4 g.) crystallised and was added to B. On refractionation, a fraction was obtained, b.p. 232–235°/685 mm., which was identified as *n*-octoic acid by titration (Found: *M*, 146. Calc.: *M*, 144) and by preparation of the *p*-toluidide, m.p. 67–69°. This m.p. was not depressed on admixture with an authentic specimen (Found: N, 6.1. Calc.: N, 6.0 per cent.). It was not found possible to separate a constant-boiling fraction from the residue.

The solid acid (B), m.p. 80–100°, was a mixture which could not be resolved by fractional crystallisation. Preliminary experiments having shown that one of the acids yielded a magnesium salt which was readily soluble in water, the mixture of acids was dissolved in dilute sodium hydroxide solution (phenolphthalein), and the boiling solution treated with an excess of magnesium sulphate. The sparingly soluble magnesium salt, B₁, was collected after 12 hours, and the

filtrate acidified; a voluminous, white precipitate then separated. This was taken up with ether (yield, 9 g.) and recrystallised from hot water and finally from ethyl acetate, flat needles being obtained (Found: C, 63.0; H, 9.6; M , 230.6. Calc.: C, 62.6; H, 9.5 per cent.; M , 230). It melted at 126–127°, alone or mixed with decane- $\alpha\kappa$ -dicarboxylic acid, prepared by Walker and Lumsden's method (*loc. cit.*).

The sparingly soluble magnesium salt B_1 , was decomposed with dilute sulphuric acid. The acid thus obtained (16 g.), m.p. 90–95°, softening at 80° (M , 255.6), was fractionally converted into the magnesium salt, four fractions being obtained. Titration showed that the last fraction, which was soluble in water, was mainly decane- $\alpha\kappa$ -dicarboxylic acid (M , 235); this was also present in not inappreciable amount in fraction 3 (M , 247.4). The first two fractions (M , 259; 256), which were evidently identical, were combined and recrystallised from ethyl acetate; an acid then separated in leaflets, m.p. 121–122° (Found: C, 65.1; H, 10.3; M , 258.9. Calc.: C, 65.1; H, 10.1 per cent.; M , 258). The barium salt also was analysed (Found: Ba, 34.8. Calc.: Ba, 34.9 per cent.). This acid has the properties of dodecane- $\alpha\mu$ -dicarboxylic acid (Brown and Walker, *Annalen*, 1891, **261**, 123); direct comparison was not possible, but the methyl ester prepared from the silver salt separated in needles, m.p. 41°, which is stated to be the m.p. of methyl dodecane- $\alpha\mu$ -dicarboxylate.

The neutral oil A (see above; 9 g.) consisted of unchanged methyl *isoerucate*.

The actual yields of the oxidation acids obtained and those calculated on the assumption that *isoerucic* acid is an equimolecular mixture of the two unsaturated acids, allowance being made for the weight of methyl *isoerucate* recovered, are: monobasic acids obtained, 16 g.; calc., 17.8 g.; dibasic acids obtained, 26 g.; calc., 28.2 g.

Oxidation of Ethyl isoErucate with Ozone.—In a preliminary experiment made with the object of determining the most suitable conditions for the oxidation of ethyl *isoerucate*, erucic acid (10 g.) was oxidised with ozone, followed by chromic acid, under the conditions described below. The mixture of acids was taken up with ether and on removal of the solvent an oil was obtained which partly crystallised. The solid, which melted at 109° after being washed with light petroleum and at 112–113° after crystallisation from benzene, was identified as brassylic acid by the method of mixed m.p. and by titration (Found: M , 243. Calc.: M , 244). The original filtrate, on dilution with light petroleum, deposited a further quantity of brassylic acid (m.p. 110°), and after the removal of this it was fractionated, b.p. 249–252°/685 mm., leaving a small residue which crystallised and

probably consisted of brassylic acid, but was not further examined. The distillate was identified as pelargonic acid by titration (Found: M , 159.5. Calc.: M , 158).

Ozone was passed through a well-cooled solution of ethyl *isoerucate* (10 g.) in acetic acid (70 c.c.) until a test portion no longer decolorised bromine. After the addition of formic acid (10 c.c.), the solution was heated on the water-bath for two hours, cooled, and treated with an aqueous solution of chromic acid sufficient in quantity to convert the aldehydes present into the corresponding carboxylic acids. The mixture of acids (from 30 g. of ethyl *isoerucate*) was dissolved in ether and, after removal of the solvent, converted into the ethyl esters. On distillation under diminished pressure (8 mm.), the following fractions were obtained: (i) up to 150° (14 g.), (ii) 150–180° (5 g.), (iii) 180–190° (8 g.), (iv) 190–210° (6 g.), (v) above 210° (3 g.). Fraction (i) consisted of the esters of the monobasic acids and was not further examined; fraction (v) crystallised on keeping and was identified as ethyl arachidate, m.p. 50° (m.p. of acid, 77°) (Found: M , 311. Calc.: M , 312).

Since on redistillation the intermediate fractions could not be obtained pure, they were combined and after hydrolysis the acids were separated by the magnesium salt method described above. The fractions of the soluble and the insoluble magnesium salts were converted into the methyl esters through the silver salts; the soluble magnesium salt then yielded a methyl ester, b.p. 175–177°/10 mm. This was methyl decane- $\alpha\kappa$ -dicarboxylate, since on hydrolysis it gave decane- $\alpha\kappa$ -dicarboxylic acid, m.p. 126–127° (Found: M , 230.6. Calc.: M , 230).

The methyl ester from the acid yielding the sparingly soluble magnesium salt distilled at 191–192°/10 mm. and solidified on keeping; it melted at 41° after crystallisation from methyl alcohol. The acid obtained on hydrolysis had m.p. 121° (Found: M , 257.6. Calc.: M , 258).

Oxidation of Erucic and Brassidic Acids.—Erucic acid, on oxidation in alkaline solution under the conditions used by Lapworth and Mottram (J., 1925, **127**, 1629) for oleic acid, gave the high-melting dihydroxybehenic acid (m.p. 127–128°; yield 36%). In cold acetone solution, it gave the low-melting isomeride, m.p. 99–100°. Under similar conditions, brassidic acid yielded only the low-melting dihydroxy-acid.

Department of Organic Chemistry,
Indian Institute of Science,
Bangalore.

[Accepted 14-4-27].

JL
A, d, m2, N 1423
N27.4A.4

PRINTED AND PUBLISHED
BY GEORGE KENNETH AT THE DIOCESAN PRESS
POST BOX 45, MADRAS-1927. C3169

Squamosa Leaves. XII.—Iodine as a Catalyst in Reactions Involving Elimination of Hydrogen Halides. XIII.—Studies Relating to the Symbiosis of Seeds and Bacteria. XIV.—The Bio-Genesis of Mahua Oil. XV.—Contributions to the Scientific Study of the Lac Industry. Parts IX—X.

VOLUME VIII (A)—(1925).

Part I.—The Photochemical Oxidation of Aromatic Hydrocarbons. I. II.—Kachi-Grass Oil. III.—Argemone Oil. IV.—Essential Oil of *Cyperus Rotundus*. V.—Dehydration of Rectified Spirit by means of Anhydrous Calcium Chloride. VI.—The Hydrolysis of the Amides of $\alpha\beta$ -Unsaturated Acids and of their Saturated Analogues. VII.—I. Studies Relating to the Acetone-Producing Organisms. II. Mahua Flowers as Raw Material for the Acetone-Fermentation Process. VIII.—Studies in Esterification. IX.—Constituents of the Marking-Nut: 'Semecarpus Anacardium' Linn. X.—Notes on Some Indian Essential Oils. XI.—Bromo-Derivatives of Para-Methoxycinnamic Acid. XII.—I. An Examination of Some Gum-Enzymes. II. Chemical Constitution of the Gum from 'Boswellia Serrata'. XIII.—Industrial Wastes as Manure. I. Ajowan and Mohua Cakes as Fertilisers. II. Utilisation of Refuse. XIV.—Constitution of Manasse's Hydroxycamphor. XV.—The Reaction Between Sodium Sulphite and Sulphur. XVI.—The Occurrence of Sylvestrene.

VOLUME VIII (B)—1925.

Part I.—Lecher Wire Measurements. II.—The Aluminium Anode Film Dielectric.

VOLUME IX (A)—1926.

Part I.—Contributions to the Scientific Study of the Lac Industry. Part XI. Early Recognition of Sex among Lac Insects, by S. Mahdihassan. Price, Rs. 4.
Part II.—Vegetable Oils Containing Glycerides of Erucic Acid, by J. J. Sudborough, H. E. Watson and P. Ramaswami Ayyar. Price, Rs. 2-8.
Part III.—The Production of Ether by Solid Catalysts, by S. K. Kulkarni Jatkar and H. E. Watson. Price Rs. 3.
Part IV.—The Constituents of Some Indian Essential Oils. Part XVIII. Derivatives of Abietic Acid, by Madyar Gopal Rau and John Lionel Simonsen. Price, As. 8.
Part V.—The Fat from 'Salvadora Oleoides': Khakan Fat, by C. K. Patel, S. Narayana Iyer, J. J. Sudborough and H. E. Watson. Price, As. 12.
Part VI.—The Constituents of Some Indian Essential Oils. Part XIX. The Essential Oil from the Rhizomes of *Kaempferia galanga* by Puthan Madhathil Bhaskara Panicker, B. Sanjiva Rao and John Lionel Simonsen. Part XX. The Essential Oil from the Rhizomes of *Curcuma Aromatica* Salisb., by B. Sanjiva Rao, Vishnu Purushottam Shintre and John Lionel Simonsen. Part XXI. The Essential Oil from the Wood of *Erythroxylon monogynum* Roxb., by B. Sanjiva Rao, Vishnu Purushottam Shintre and John Lionel Simonsen. Price, As. 12.
Part VII.—Reactions of Chromates at High Temperatures, Part II. The System $\text{CaO} - \text{Cr}_2\text{O}_3 - \text{O}_2$, by K. S. Nargund and H. E. Watson. Price, Re. 1.
Part VIII.—I. The Catalytic Hydrogenation of Carone, by Subramania Narayana Iyer and John Lionel Simonsen. II. Conessine, by Darab Dinsha Kanga, Panchandana Ramaswami Ayyar and John Lionel Simonsen. Price, As. 8.

Part IX.—Derivatives of Naphthaquinolines and Naphthaisoquinolines. By Charles Stanley Gibson, Kalvoi Venkatakrishna Hariharan, Kottiazath Narayana Menon and John Lionel Simonsen. Price As. 12.

Part X.—I. Isomeric Phenylserines. II. The Unstable Modification of *iso*Nitrosocamphor. III. *d*-Mannitol from *Gardenia Turgida*, by M. O. Forster and Keshaviah Aswath Narain Rao. Price, As. 12.

VOLUME IX (B)—1926.

Part I.—Double Synchronous Speed Alternators, by K. P. Roy. Price, Re. 1.

Part II.—The Characteristics of Beam Transmitting Aerials, by J. K. Catterson-Smith. Price, Re. 1.

Part III.—Aerial Testing, by J. K. Catterson-Smith. Price, Rs. 1-8.

Part IV.—Circulating Currents in Wave-Wound Armatures, by F. N. Mowdawalla and G. K. Pradhan. Price, Rs. 1-8.

Part V.—Madras (Fort) Radio Field Intensity Measurements at Bangalore, by K. Sreenivasan. Price, Rs. 3.

VOLUME X (A)—1927.

Part I.—I. Some Reactions of Carone, by Kottiazath Narayana Menon and John Lionel Simonsen. II. A Synthesis of Morindone, by Ramkanta Bhattacharya and John Lionel Simonsen. III. Derivatives of Acenaphthpyridine. Part I, by Sri Kumaran Unni Nair and John Lionel Simonsen. Price, As. 12.

Part II.—Oils and Fats from the Seeds of Indian Forest Plants. Part VIII. The Oil from the Seeds of *Thevetia Nerifolia* (Juss.). By Ramkanta Bhattacharya and P. Ramaswami Ayyar. Part IX. The Oil from the Seeds of *Cerbera Odollam* (Gaertn.). Part X. The Oil from the Seeds of *Holarrhena Antidysenterica*. Part XI. The Oil from the Seeds of *Anona Squamosa* (Linn.). By Ramachandra Vishnu Ghanekar and P. Ramaswami Ayyar. Price As. 12.

Part III.—Influence of the Sulphur Atom on Optical Rotatory Power. By P. P. Shukla. Price, As. 8.

Part IV.—The Constitution of the Acid formed by the Action of Sulphuric Acid on Camphorquinone. By Madhay Balaji Bhagvat and John Lionel Simonsen. Price, As. 8.

Part V.—*iso*Erucic Acid. By Tahilram Jethanand Mirchandani and John Lionel Simonsen. Price, As. 8.