

61

Price, As. 12.]

[Vol. 11A, Part III, pp. 23 to 29.]

JOURNAL

OF THE

Indian Institute of Science.

CONTENTS.

CONTRIBUTIONS TO THE STUDY OF SPIKE-DISEASE OF SANDAL
(SANTALUM ALBUM, LINN.).
PART I. DIASTATIC ACTIVITY OF THE LEAVES.

BY

M. Sreenivasaya and B. N. Sastri.

DR. M. O. FORSTER, F.R.S., CHAIRMAN OF EDITORIAL BOARD.

Printed

Price.

JOURNAL
OF THE
INDIAN INSTITUTE OF SCIENCE

- Parts I—XXI. VOLUME I.—(1914-18).
Parts I—XV. VOLUME II.—(1918-19).
Parts I—IX. VOLUME III.—(1920).
Parts I—XVI. VOLUME IV.—(1921).
Parts I—XIV. VOLUME V.—(1922).
VOLUME VI.—(1923).

Part I.—Mahua Oil. II.—Malabar Sardine Oil. III.—Studies in Esterification. IV.—The Action of Bases on $\alpha\alpha\beta$ - and $\alpha\beta\beta$ -Tribromo- β -phenylpropionic Acids and their Esters. V.—Hongay Oil. VI.—Cashew Kernel Oil. VII.—The Fruit of *Bassia Longifolia*. VIII.—Studies in Intensive Bacterial Oxidation, I. The Oxidation of Alcohol to Acetic Acid. IX.—Contributions to the Study of the Prickly Pear Problem. X.—A Chemical Examination of Samples of Water and of Shells from the Inle Valley. XI.—The Refining of Crude Saltpetre. XII.—Enolising Action of Hydrogen Chloride on Aldehydes and Ketones. XIII.—The Conversion of Eugenol into Isoeugenol.

VOLUME VII.—(1924).

Part I.—Studies in Alcoholysis V. The Alcoholysis of Esters of $\alpha\beta$ -unsaturated Acids and of the corresponding saturated Esters. II.—Chemical Factors in Denitrification. III.—The Retting of Coconut Husk for the Production of Coir. IV.—Reactions of Chromates at High Temperatures. Part I. The Synthesis and Decomposition of Calcium, Sodium and Magnesium Chromates in Air. V.—Oil from *Mimusops Hexandra*: Rayan Oil. VI.—The Relation Between the Iodine Values and Refractive Indices of Hardened Oils. Part II. VII.—Contributions to the Scientific Study of the Lac Industry. Parts I—VIII. VIII.—I. The Systematic Nomenclature of Polycyclic Carbon Systems. II. Polycyclic and Cage Systems of Carbon Compounds. III. The Systematic Nomenclature of Heterocyclic Compounds including Polycyclic Structures. IX.—The Use of Mixed Catalysts in the Hydrogenation of Oils. X.—Condensation of Aromatic Amines with Chloroform or Carbon Tetrachloride in the Presence of Finely Divided Copper. XI.—I. Notes on Bixin II. An Alkaloid from *Anona Squamosa* Leaves. XII.—Iodine as a Catalyst in Reactions Involving Elimination of Hydrogen Halides. XIII.—Studies Relating to the Symbiosis of Seeds and Bacteria. XIV.—The Bio-Genesis of Mahua Oil. XV.—Contributions to the Scientific Study of the Lac Industry. Parts IX—X.

VOLUME VIII (A)—(1925).

Part I.—The Photochemical Oxidation of Aromatic Hydrocarbons. I. II.—Kachi-Grass Oil. III.—Argemone Oil. IV.—Essential Oil of *Cyperus Rotundus*. V.—Dehydration of Rectified Spirit by means of Anhydrous Calcium Chloride. VI.—The Hydrolysis of the Amides of $\alpha\beta$ -Unsaturated Acids and of their Saturated Analogues. VII.—I. Studies Relating to the Acetone-Producing Organisms. II. Mahua Flowers as Raw Material for the A mentation Process.

Price, As. 12.]

[Vol. 11A, Part III, pp. 23 to 29.]

JOURNAL

OF THE

Indian Institute of Science.

CONTENTS.

CONTRIBUTIONS TO THE STUDY OF SPIKE-DISEASE OF SANDAL
(SANTALUM ALBUM, LINN.).
PART I. DIASTATIC ACTIVITY OF THE LEAVES.


BY

M. Sreenivasaya and B. N. Sastri.



DR. M. O. FORSTER, F.R.S., CHAIRMAN OF EDITORIAL BOARD.

12345



Am 2, N14

N28-11A-3

CONTRIBUTIONS TO THE STUDY OF SPIKE-DISEASE OF SANDAL (SANTALUM ALBUM, LINN.).

Part I. Diastatic Activity of the Leaves.

By M. Sreenivasaya and B. N. Sastri.

It is well known that in the case of many plant diseases the pathological condition is indicated by a marked disturbance in the enzymatic activity of the tissues. From this point of view the oxidising enzymes have been those most investigated and it has been found that the oxidase-activity of the diseased tissue is not infrequently intensified abnormally. Mosaic disease of tobacco, potato "Leaf-roll" disease and "Curly-top" of sugar-beet are examples in which this occurs. Other groups of enzymes have received comparatively little attention. The diastase-content of diseased potatoes has however been found greater than that of the normal tubers. Coleman investigated the diastatic activity of sandal-leaves, both healthy and spiked, in an endeavour to find an explanation for the strikingly abnormal accumulation of starch in the diseased leaves and other tissues of spiked sandal. He came to the conclusion, however, that the diastatic activity of spiked leaves was lower than that of the healthy ones, and explained starch-accumulation on the basis of defective translocation of the photosynthetic carbohydrates due to poor diastatic activity.

During the course of a detailed investigation of the various enzymes of sandal-tissues, both diseased and healthy, our experience has been entirely different and we have consistently found higher diastatic activity in the diseased leaves. Coleman's remark however that the spiked leaves are really comparable in composition with young healthy leaves holds true, not only with respect to their chemical composition but also with regard to enzymic activity. The spiked leaves are as rich in diastase as the growing points of a healthy sandal (see Table IV).

MATERIAL AND METHODS: COLLECTION OF SAMPLES.

Samples of leaves from spiked and healthy sandal trees were collected from two different areas, Uttarhalli and Ragihalli, the former lying at a distance of about 6 miles from Bangalore on the

Ka
sout
a nur
the U

the latter near Banneraghatta, about 13 miles from the Ragihalli area. Both the areas have been under observation for a number of years and the first symptoms of spike were observed in the Ragihalli area in the year 1914 (Coleman, *Spike Disease of Sandal*, 1917). No record is so far available to us with regard to the date of incidence of spike in the Ragihalli area. Samples were taken from a number of selected trees situated at different points on the area. The well-known external symptoms were taken as the criteria for determining the spiked condition of plants. In the present state of our knowledge we are not in a position to state definitely that the apparently disease-free plants from which we have collected our research material are absolutely healthy though we believe this to be the case. As they are growing in a spiked area it is possible they may already have become infected, though they show no symptom from which this could be deduced. Moreover, the healthy trees have, so far as all external appearance goes, remained healthy and, if infected at all, could only have been in the earliest stage of the disease.

Leaves and stems were collected between 8.30 and 9.30 a.m., and were taken from every one of the marked trees so that a representative sample was ensured. The material was placed in glass-stoppered bottles immediately after plucking, in order to minimise any loss of moisture due to transpiration, and carried to the laboratories in cold storage. In the case of the Uttaraghatta samples the time taken for transferring the bottle from the site of collection to the ice-chest was about 30 minutes, while roughly two hours elapsed with the samples from the Ragihalli area. The samples collected on any one day were at least comparable between themselves.

PREPARATION OF MATERIAL FOR ANALYSIS.

Expression of saps.—About 100 grams of leaves after separation from the stems were weighed into a nickel-wire-gauze holder which was afterwards immersed in liquid air and allowed to remain there until the liquid ceased to boil. The material was removed, and allowed to attain room-temperature in a closed bottle in order to prevent the condensation of atmospheric moisture. The material was then ground in a porcelain mortar, wrapped in a filter-cloth (previously wetted and pressed out at a pressure of one ton to the square inch) and pressed in a nickel-plated hydraulic press at one ton to the square inch.

To secure uniformity in composition and yield of sap from materials collected on different days the following arbitrary standards were fixed. The charge of leaves was 100 gms., pressure applied one ton; time allowed for expression 15 minutes. The sap expressed was weighed

also the leaf residue. The losses due to imperfect removal of the fluid from the parts of the press do not usually exceed one or two per cent. The sap was immediately centrifuged for 15 minutes at 3,000 revolutions per minute, toluene was added, and the sap preserved in the ice-chest. The conditions for the expression of saps from the stems were nearly the same, except that a pressure of 2 tons to the square inch was applied.

Preparation of leaf-powders.—The leaves were spread in thin layers and dried over freshly fused calcium chloride in a vacuum-desiccator, 24 hours being usually sufficient for complete drying. The material was then powdered in a porcelain mortar, passed through a 40-mesh sieve, and preserved in glass-stoppered bottles.

Methods of estimating diastatic activity.—The sap (2 c.c.) was pipetted into a 50 c.c. conical flask containing 25 c.c. of 2 per cent. soluble starch and a little toluene, the reaction being allowed to proceed at room-temperature. The necessary controls were run. After 20 hours the mixture was treated with 5 c.c. of dialysed iron and centrifuged. Sugars were estimated in an aliquot portion of the clear centrifugate by Bertrand's method and calculated as maltose.

The diastatic activity of leaf-powders was estimated similarly, 0.25 gm. of the dried powder being allowed to act upon 25 c.c. of 2 per cent. soluble starch solution. The necessary controls were run. The toluene added to the mixture serves not only as an antiseptic but also as an autolysing agent. After 20 hours the mixtures were treated with dialysed iron, centrifuged, and an aliquot portion of the clear fluid used for the estimation of reducing sugars by Bertrand's method. The coloration developed with iodine was noted in each case.

Results.—The diastatic activity of the saps, expressed in milligrams of maltose per cubic centimeter of sap, is shown in Table I. In every instance without exception the diastatic activity of saps from spiked leaves is definitely greater than that of healthy leaf-saps. In some cases the activity of spiked leaf-sap is found to be four times that of the healthy sample. Table II includes a few results shewing the diastatic activity of diseased and healthy stem-saps. A distinctly higher diastatic activity is again shown by the spiked specimens. Table III comprises the results obtained with the dry leaf-powders, diseased and healthy, expressed in grams of maltose produced per gram of leaf-powder. All the results point to a higher diastatic activity in the case of the spiked leaf. The colour reactions with iodine confirm this result.

TABLE I.
Diastatic Activity of Leaf-sap.

Date in 1927	Locality	Diastatic activity of 1 c. c. of sap			
		Healthy		Spike	
		Maltose in mgms.	Coloration with iodine	Maltose in mgms.	Coloration with iodine
April 27 ...	Uttarahalli	77.5	Red	161.0	None
May 12 ...	"	66.0	Blue	114.7	Red
" 19 ...	"	101.5	Deep Red	130.5	None
June 16 ...	"	66.0	"	144.0	"
" 23 ...	"	109.6	"	168.0	"
August 5 ...	"	75.0	"	152.0	"
" 11 ...	Ragihalli	28.0	Blue	126.0	"
October 27 ...	Uttarahalli	44.2	"	172.8	"
November 7 ...	Ragihalli	46.4	"	86.0	Blue

TABLE II.
Diastatic Activity of Stem-saps.


Date in 1927	Locality	Diastatic activity of 1 c. c. of sap			
		Healthy		Spike	
		Maltose in mgms.	Coloration with iodine	Maltose in mgms.	Coloration with iodine
July 20 ...	Uttarahalli	19.2	Blue	37.8	Blue
November 7 ...	Ragihalli	27.0	"	38.0	"

TABLE III.
Diastatic Activity of Leaf-powders.

Date in 1927	Locality	Activity of one gram of dry leaf-powder			
		Healthy		Spike	
		Maltose in gms.	Coloration with iodine	Maltose in gms.	Coloration with iodine
April 14 ...	Ragihalli	0.198	Reddish blue	0.980	Red
" 20 ...	Uttarahalli	0.204	"	0.356	"
May 5 ...	Ragihalli	0.610	Bluish Red	1.080	Slightly Red
" 12 ...	Uttarahalli	0.039	Reddish blue	0.906	Red
" 19 ...	"	0.192	Blue	0.294	Bluish Red
June 2 ...	"	* 0.664	Bluish Red	* 1.100	Colourless
" 16 ...	"	0.132	Reddish blue	0.287	Red

* Average of four determinations.

to expr



Gradient Study.—The variation in diastatic activity of the leaves according to the height of insertion in the stem was followed by a gradient-study. The results collected in Table IV reveal an almost entire absence of any gradient in the case of the spiked leaves, while a steady fall in the diastatic activity from the growing tip downwards is shewn in the healthy shoots. The results have also been represented graphically (Graph I).


TABLE IV.

Gradient of Diastatic Activity (June 2, 1927, Uttarahalli).

Sample	Activity of dry leaf-powders			
	Healthy		Spiked	
	Maltose in mgms.	Coloration with iodine	Maltose in mgms.	Coloration with iodine
Growing tips East...	120.0	Colourless	127.2	Colourless
I " ...	114.4	Red	128.0	"
II " ...	89.4	Red	126.4	"
III " ...	—	—	50.0	Red
Growing tips West...	102.0	Colourless	122.4	Colourless
I " ...	63.6	Blue	122.0	"
II " ...	—	—	127.2	"
III " ...	33.6	Blue	68.4	Red

TIME-COURSE STUDY.

A study of the time-course of diastatic action was made with the leaf extracts. Leaf-powder by itself could not be employed since the solid material interferes with the subsequent operations of drawing out definite quantities of the reaction mixture for analysis. The enzyme extract was prepared as follows: Desiccated leaf-powder (15 gms.) was digested at room-temperature with 150 c.c. of water containing a little toluene for 20 hours in a 250 c.c. glass-stoppered bottle shaken at frequent intervals. Hide-powder (5 gms.) was added for the abstraction of the tannins which might inhibit the activity of diastase, it having been previously ascertained that hide-powder did not adsorb any diastase. The whole digest was centrifuged, the supernatant liquid passed through a filter and the clear extract used for the experiments. The reaction mixture consisted of 100 c.c. of a 4 per cent. Lintner's soluble starch solution, 100 c.c. of the enzyme extract and 5 c.c. toluene. The hydrolysis was allowed to proceed at 30° in a thermostat. Definite quantities of the mixture were withdrawn from



the experimental and control flasks and examined polarimetrically, the reducing sugars being estimated by Bertrand's method.

An examination of the tables (V and VI) and the curves (graph II) brings out strikingly the higher diastatic activity of the diseased leaf extract. Whether these results indicate differences in the absolute amount of diastase or are explicable by the presence of inhibitors or activators in the respective samples is undoubtedly of great interest. The curves point however to a definitely lower enzyme-concentration.

TABLE V.


Time-course of Diastatic Action (Healthy Leaf Extract).

Time	Diastatic activity		Remarks
	Maltose in mgms.	Coloration with iodine	
30 minutes	2.4	Blue	Leaves collected at 8-30 A.M. on 4-11-27 from healthy sandal trees on the Institute Estate
60 "	4.7	Blue (tinged red)	
135 "	10.7	Bluish violet	
180 "	12.2	Violet	
300 "	18.0	Purple	
420 "	23.3	Brick red	
21.75 hours	58.3	Slight red	
47 "	78.5	Colourless	

TABLE VI.

Time-course of Diastatic Action (Healthy and Spike Leaf Extracts).

Time		Healthy		Spike	
		Maltose in mgms.	Coloration with iodine	Maltose in mgms.	Coloration with iodine
Hrs.	Mins.				
0	30	2.5	Blue	18.5	Blue
1	15	4.7	"	37.5	Reddish blue
2	15	8.9	Violet	58.0	Violet brown
3	15	11.5	Reddish violet	75.0	Dull red
4	15	19.2	"	77.0	"
5	45	24.5	Brown	89.0	Colourless
7	15	25.0	Red	96.0	"
8	45	27.0	"	97.0	"
10	15	32.4	"	103.0	"
12	45	50.4	"	114.0	"
22	30	60.0	Slight Red		"



in the healthy leaf extract. In addition to a higher enzyme-concentration in the spike-leaf extract however, the lower pH value (4.7), which approximates to the optimum reaction of plant amylases, the lower calcium content, and the higher amino-nitrogen value of the diseased leaf extract, possibly contribute towards the activation of the enzyme derived from the spiked leaf. Further work to examine the influence of these and other factors is in progress.

In conclusion it is our pleasant duty to express our gratitude to Professor Norris for the keen interest he has evinced in the work and for his many helpful suggestions.

*Department of Bio-Chemistry,
Indian Institute of Science,
Bangalore.*

[Accepted, 14-12-27.]



*Am 2, 114
28.10.3*

PRINTED AND PUBLISHED
BY GEORGE KENNETH AT THE DIOCESAN PRESS,
POST BOX 433, MADRAS-1928. C4561

VIII.—Studies in Esterification. IX.—Constituents of the Marking-Nut: 'Semecarpus Anacardium' Linn. X.—Notes on Some Indian Essential Oils. XI.—Bromo-Derivatives of Para-Methoxycinnamic Acid. XII.—I. An Examination of Some Gum-Enzymes. II. Chemical Constitution of the Gum from 'Boswellia Serrata'. XIII.—Industrial Wastes as Manure. I. Ajowan and Mohua Cakes as Fertilisers. II. Utilization of Refuse. XIV.—Constitution of Manasse's Hydroxycamphor. XV.—The Reaction Between Sodium Sulphite and Sulphur. XVI.—The Occurrence of Sylvestrene.

VOLUME VIII (B)—1925.

Part I.—Lecher Wire Measurements. II.—The Aluminium Anode Film Dielectric.

VOLUME IX (A)—1926.

- Part I.—Contributions to the Scientific Study of the Lac Industry. Part XI. Early Recognition of Sex among Lac Insects, by S. Mahdihassan. Price, Rs. 4.
- Part II.—Vegetable Oils Containing Glycerides of Erucic Acid, by J. J. Sudborough, H. E. Watson and P. Ramaswami Ayyar. Price, Rs. 2-8.
- Part III.—The Production of Ether by Solid Catalysts, by S. K. Kulkarni Jatkar and H. E. Watson. Price, Rs. 3.
- Part IV.—The Constituents of Some Indian Essential Oils. Part XVIII. Derivatives of Abietic Acid, by Madyar Gopal Rau and John Lionel Simonsen. Price, As. 8.
- Part V.—The Fat from 'Salvadora Oleoides': Khakan Fat, by C. K. Patel, S. Narayana Iyer, J. J. Sudborough and H. E. Watson. Price, As. 12.
- Part VI.—The Constituents of Some Indian Essential Oils. Part XIX. The Essential Oil from the Rhizomes of *Kaempferia galanga*, by Puthan Madhathil Bhaskara Panicker, B. Sanjiva Rao and John Lionel Simonsen. Part XX. The Essential Oil from the Rhizomes of *Curcuma Aromatica* Salisb., by B. Sanjiva Rao, Vishnu Purushottam Shintre and John Lionel Simonsen. Part XXI. The Essential Oil from the Wood of *Erythroxylon monogynum* Roxb., by B. Sanjiva Rao, Vishnu Purushottam Shintre and John Lionel Simonsen. Price, As. 12.
- Part VII.—Reactions of Chromates at High Temperatures, Part II. The System $\text{CaO} - \text{Cr}_2\text{O}_3 - \text{O}_2$, by K. S. Nargund and H. E. Watson. Price, Re. 1.
- Part VIII.—I. The Catalytic Hydrogenation of Carone, by Subramania Narayana Iyer and John Lionel Simonsen. II. Conessine, by Darab Dinsha Kanga Panchanada Ramaswami Ayyar and John Lionel Simonsen. Price, As. 8.
- Part IX.—Derivatives of Naphthaquinolines and Naphthaisoquinolines. By Charles Stanley Gibson, Kalvoi Venkatakrishna Hariharan, Kottiazath Narayana Menon and John Lionel Simonsen. Price As. 12.
- Part X.—I. Isomeric Phenylserines. II. The Unstable Modification of isoNitroso-camphor. III. *d*-Mannitol from *Gardenia Turgida*, by M. O. Forster and Keshaviah Aswath Narain Rao. Price, As. 12.

VOLUME IX (B)—1926.

- Part I.—Double Synchronous Speed Alternators, by K. P. Roy. Price, Re. 1.
- Part II.—The Characteristics of Beam Transmitting Aerials, by J. K. Catterson-Smith. Price, Re. 1.
- Part III.—Aerial Testing, by J. K. Catterson-Smith. Price, Rs. 1-8.
- Part IV.—Circulating Currents in Wave-Wound Armatures, by F. N. Mowdawalla and G. Pradhan. Price, Rs. 1-8.

Part V.—Madras (Fort) Radio Field Intensity Measurements at Bangalore, by K. Sreenivasan. Price, Rs. 3.

VOLUME X (A)—1927.

- Part I.—I. Some Reactions of Carone, by Kottiazath Narayana Menon and John Lionel Simonsen. II. A Synthesis of Morindone, by Ramkanta Bhattacharya and John Lionel Simonsen. III. Derivatives of Acenaphthpyridine. Part I, by Srikumaran Urni Nair and John Lionel Simonsen. Price, As. 12.
- Part II.—Oils and Fats from the Seeds of Indian Forest Plants. Part VIII. The Oil from the Seeds of *Thevetia Neriifolia* (Juss.) By Ramkanta Bhattacharya and P. Ramaswami Ayyar. Part IX. The Oil from the Seeds of *Cerbera Odollam* (Gaertn.). Part X. The Oil from the Seeds of *Holarrhena Antidysenterica*. Part XI. The Oil from the Seeds of *Anona Squamosa* (Linn.). By Ramachandra Vishnu Ghanekar and P. Ramaswami Ayyar. Price As. 12.
- Part III.—Influence of the Sulphur Atom on Optical Rotatory Power. By P. P. Shukla. Price, As. 8.
- Part IV.—The Constitution of the Acid formed by the Action of Sulphuric Acid on Camphorquinone. By Madhav Balaji Bhagvat and John Lionel Simonsen. Price, As. 8.
- Part V.—*iso*Erucic Acid. By Tahilram Jethanand Mirchandani and John Lionel Simonsen. Price, As. 8.
- Part VI.—Note on the Sorting, Classification and Briquetting of Chrome and Manganese Ores. By K. R. Krishnaswami. Price, As. 8.
- Part VII.—Esterification in Mixed Solvents, by B. V. Bhide and H. E. Watson. Price, As. 8.
- Part VIII.—Nitrogen-Fixation by 'Azotobacter Chroococcum'. By S. Ranganathan and Roland V. Norris. Price, Rs. 1-4.
- Part IX.—Studies in intensive Bacterial Oxidation. The Oxidation of Ammonia to Nitric Acid, Parts I—IV. By Gilbert J. Fowler, Y. N. Kotwal, Roland V. Norris, S. Ranganathan and M. B. Roy. Price, Re. 1.
- Part X.—The Decomposition of Alkaline Earth Sulphates. By L. A. Bhatt and H. E. Watson. Price, As. 12.

VOLUME X (B)—1927.

- Part I.—Tests on Suspension Insulators After Ten Years' Service. By A. S. Venkateswaran and U. Ganguly. Price, Re. 1.
- Part II.—The Measurement of Voltage Gradient on a String of Suspension Insulators. By G. Yoganandam and R. K. Sen. Price, Re. 1.
- Part III.—Intensity Variations of Madras (Fort) Radio Station. By K. Sreenivasan. Price, As. 8.
- Part IV. Suspension Insulator Testing. By G. Yoganandam. Price, As. 8.

VOLUME XI (A)—1928.

- Part I.—I. Studies on Invertase, Part I. Preparation and Purification of the Enzyme. By B. N. Sastri and Roland V. Norris. II. Note on a Simple Method for concentrating Enzyme Solutions. By B. N. Sastri. Price, Re. 1.
- Part II.—The Bleaching of Lac. By M. Venugopalan. Price, As. 8.
- Part III.—Contributions to the Study of Spike-Disease of Sandal (*Santalum Album*, Linn.). Part I. Diastatic Activity of the Leaves. By M. Sreenivasaya and B. N. Sastri. Price, As. 12.