

Price, Re. 1.]

[Vol. 11A, Part XIX, pp. 241 to 247.]

12

JOURNAL

OF THE

Indian Institute of Science.

CONTENTS.

I. CONTRIBUTIONS TO THE STUDY OF SPIKE-DISEASE OF
SANDAL (SANTALUM ALBUM, LINN.).

PART IV. CHEMICAL COMPOSITION OF HEALTHY AND
SPIKED SANDAL STEMS.

BY

D. A. Rama Rao, and M. Sreenivasaya.

II. CONTRIBUTIONS TO THE STUDY OF SPIKE DISEASE OF
SANDAL (SANTALUM ALBUM, LINN.).

PART V. TRANSMISSION OF SPIKE BY BUDDING.

BY

M. Sreenivasaya and G. Gopalaswami Naidu.

DR. M. O. FORSTER, F.R.S., CHAIRMAN OF EDITORIAL BOARD.

JOURNAL
OF THE
INDIAN INSTITUTE OF SCIENCE

- Parts I—XXI. VOLUME I.—(1914-18).
 Parts I—XV. VOLUME II.—(1918-19).
 Parts I—IX. VOLUME III.—(1920).
 Parts I—XVI. VOLUME IV.—(1921).
 Parts I—XIV. VOLUME V.—(1922).
 Parts I—XIII. VOLUME VI.—(1923).
 Parts I—XV. VOLUME VII.—(1924).
 Parts I—XVI. VOLUME VIII (A)—(1925).
 VOLUME VIII (B)—1925.
 Part I.—Lecher Wire Measurements. II.—The Aluminium Anode Film Dielectric.
 VOLUME IX (A)—1926.
 Parts I—X. VOLUME IX (B)—1926.
 Part I.—Double Synchronous Speed Alternators, by K. P. Roy. Price, Re. 1.
 Part II.—The Characteristics of Beam Transmitting Aerials, by J. K. Catterson-Smith.
 Price, Re. 1.
 Part III.—Aerial Testing, by J. K. Catterson-Smith. Price, Rs. 1-8.
 Part IV.—Circulating Currents in Wave-Wound Armatures, by F. N. Mowdawalla and
 G. K. Pradhan. Price, Rs. 1-8.
 Part V.—Madras (Fort) Radio Field Intensity Measurements at Bangalore, by K. Sree-
 nivasan. Price, Rs. 3.
 VOLUME X (A)—1927.
 Part I.—I. Some Reactions of Carone, by Kottazath Narayana Menon and John Lionel
 Simonsen. II. A Synthesis of Morindone, by Ramkanta Bhattacharya
 and John Lionel Simonsen. III. Derivatives of Acenaphthpyridine.
 Part I, by Srikumaran Unni Nair and John Lionel Simonsen. Price,
 As. 12.
 Part II.—Oils and Fats from the Seeds of Indian Forest Plants. Part VIII. The Oil
 from the Seeds of *Thevetia Nerifolia* (Juss.) By Ramkanta Bhatta-
 charya and P. Ramaswami Ayyar. Part IX. The Oil from the Seeds of
Cordia Odollam (Gaertn.). Part X. The Oil from the Seeds of
Holarrhena Antidysenterica. Part XI. The Oil from the Seeds of *Anona*
Squamosa (Linn.). By Ramachandra Vishnu Ghanekar and P. Ramaswami
 Ayyar. Price, As. 12.
 Part III.—Influence of the Sulphur Atom on Optical Rotatory Power. By P. P. Shukla.
 Price, As. 8.

Price, Re. 1.]

[Vol. 11A, Part XIX, pp. 241 to 247.

JOURNAL

OF THE

Indian Institute of Science.

CONTENTS.

- I. CONTRIBUTIONS TO THE STUDY OF SPIKE-DISEASE OF
 SANDAL (SANTALUM ALBUM, LINN.).
 PART IV. CHEMICAL COMPOSITION OF HEALTHY AND
 SPIKED SANDAL STEMS.

BY


D. A. Rama Rao and M. Sreenivasaya.

- II. CONTRIBUTIONS TO THE STUDY OF SPIKE DISEASE OF
 SANDAL (SANTALUM ALBUM, LINN.).
 PART V. TRANSMISSION OF SPIKE BY BUDDING.

BY

M. Sreenivasaya and G. Gopalaswami Naidu.

DR. M. O. FORSTER, F.R.S., CHAIRMAN OF EDITORIAL BOARD.



Part IV.—Chemical composition of healthy and spiked Sandal stems.

By D. A. Rama Rao and M. Sreenivasaya.

Sandal stems 0.20 to 0.50 cm. in diameter derived from diseased and healthy sandal trees of the same area were dried at 70° and analysed for the main mineral constituents, total nitrogen and starch. The dried substance was ground to pass through a 30-mesh sieve and preserved in stoppered bottles for subsequent analysis. Methods employed for the estimation of total nitrogen, phosphorus, sulphur, calcium and potassium were the official methods recommended by the Association of Agricultural Chemists (official and tentative methods of the A.O.A.C., Washington). Starch was estimated after thorough extraction of the weighed sample in a Soxhlet with 90 per cent. alcohol for six hours, by the taka-diastase method (Davis and Daish, *J. Agric. Sci.*, 1914, 6, 152).

From the results tabulated below it will be seen that the diseased stems have usually a low ash content, a uniformly high nitrogen value calculated on the moisture-free material, a low calcium and potash content and a higher percentage of phosphorus, all the three being calculated on the weight of ash. The starch content of the diseased stems is also higher than that of the healthy stems as already found by Coleman and his colleagues (Spike-disease of Sandal, 1917, Bulletin No. 3, Mycological Series, Department of Agriculture, Mysore).

TABLE I.

Total Ash and Nitrogen Content of Sandal Twigs.

Date in 1927	Total ash per cent.		Total nitrogen in mgms. per 100 gms.	
	HEALTHY	SPIKED	HEALTHY	SPIKED
UTTARAHALLI AREA				
July 15 ...	2·950	2·611	554·0	887·0
August 18 ...	2·618	3·069	752·0	1157·0
September 1 ...	5·214	4·645	1279·0	1548·0
„ 15 ...	1·960	1·954	645 0	1021·0
Average for the season ...	3·186	3·070	807·5	1153·3


Total Ash and Nitrogen Content of Twigs.—(continued).

Date in 1927	Total ash per cent.		Total nitrogen in mgms. per 100 gms.	
	HEALTHY	SPIKED	HEALTHY	SPIKED
RAGIHALLI AREA				
May 5 ...	6·870	4·550	1092·0	1960·0
September 8 ...	2·177	2·050	566·0	642·0
„ 13 ...	2·598	2·348	665·0	666·0
October 14 ...	2·113	2·304	577·0	635·0
November 23 ...	2·770	2·320	651·0	724·5
December 19 ...	2·000	2·220	646·2	831·7
Average for the season ...	3·088	2·632	699·5	909·5

TABLE II.

Ash Constituents of Sandal Twigs.

Date in 1927	Ca per cent.		K ₂ O per cent.		P ₂ O ₅ per cent.	
	HEALTHY	SPIKED	HEALTHY	SPIKED	HEALTHY	SPIKED
UTTARAHALLI AREA						
July 15 ...	31·93	17·47	13·67	12·59	2·00	5·37
August 18 ...	24·20	19·48	27·87	41·77	5·27	2·70
September 1 ...	13·75	8·21	17·63	17·20	4·43	5·22
„ 15 ...	21·43	22·82	46·22	25·44	8·06	8·15
Average for the season ...	22·83	16·99	26·35	24·25	4·94	5·36
RAGIHALLI AREA						
May 5 ...	14·78	12·18	21·84	26·52	0·24	2·55
September 8 ...	17·90	23·66	21·07	15·00	1·53	1·94
„ 13 ...	19·43	30·83	30·27	22·50	2·41	2·61
October 14 ...	1·28	1·00	27·54	22·68	3·23	4·13
November 23 ...	12·00	11·40	52·35	33·62	2·20	2·65
December 19 ...	18·20	17·30	54·50	36·50	1·31	1·67
Average for the season ...	13·92	13·11	34·60	26·14	1·82	2·59



The figures indicate that the more important constituents nitrogen, calcium, phosphorus and potash in the diseased plant are equally subject to seasonal influences with the healthy plant itself; but the differences are consistently maintained throughout the season under experiment. The chart of the phosphate values brings out this point quite clearly.

A study of the figures in the two tables also shows that the average value for the season for a particular constituent, differs for different areas. Taking total nitrogen into consideration, Uttarhalli samples show a higher nitrogen content than those of Ragihalli. The ecological factors, more especially the flora associated with sandal, have a distinct influence in controlling the composition of the parasitic sandal. Each area has therefore its own definite average value with regard to every one of its constituents. It would be misleading therefore to compare the values of one area with those of another.

SUMMARY.

Diseased sandal stems have usually a lower ash, calcium and potash content than the healthy ones, but a definitely higher nitrogen, phosphorus and starch content in any one area.

The constituents are equally subject to seasonal variations in the diseased as well as the healthy stems, there being a consistent variation between them.

The average value for the season for a particular constituent varies for different areas.

*Department of Bio-Chemistry,
Indian Institute of Science,
Bangalore.*

[Accepted, 11-10-28.]

CONTRIBUTIONS TO THE STUDY OF SPIKE DISEASE OF SANDAL (*SANTALUM ALBUM*, LINN.).

Part V.—Transmission of Spike by Budding.

By M. Sreenivasaya and G. Gopalaswami Naidu.

The development of a suitable and reproducible technique for artificial transmission of the disease usually forms an important step in the investigation of infectious diseases of plants. Virus diseases have been communicated to healthy stocks in various ways depending upon the virulence of the virus. In the case of several mosaic diseases transmission can be effected through mere pin pricks with an infected needle, while in other cases a hypodermic injection of the infectious sap, or a transplantation of the diseased tissue on to the susceptible region of the healthy stock is necessary to induce the disease. In yet other cases, an actual grafting or budding of the diseased tissue on to the healthy stock may be necessary. Coleman has transmitted spike disease of sandal to healthy stocks through grafting. According to him an intimate organic connection of the tissues is necessary for the disease transference to occur. In his Bulletin (Spike Disease of Sandal, Bulletin No. 3, 1917, Mycological Series, Department of Agriculture, Mysore), he does not, however, mention the percentage of success he was able to achieve by his method of grafting nor does he refer to the behaviour of any controls that may have been run. His experiments appear therefore to be mainly qualitative in character and the effect of hosts on the susceptibility of sandals to graft-infection is not considered. The graft-infections were carried out on stocks growing under natural, i.e., uncontrolled conditions in the soil.

With regard to budding, however, he says, 'Butler, in 1904, following the work of Smith on "Peach Yellows" attempted to produce the disease by budding. For this purpose he took buds from diseased trees and grafted them on to healthy ones. Although he tried some one hundred and fifty budding experiments, none of the buds grew, so that the results of the experiments gave no information of value. In 1914, similar experiments were commenced by us in Bangalore, but it was soon found that budding was not a suitable method of grafting for sandal as only one of the buds formed in connection with the healthy stock and even this one died soon after.' So far the disease has not been communicated by an injection of the maceration extract of the diseased tissues.

Results of experiments on the artificial transmission of spike by budding are recorded in this paper.

EXPERIMENTAL.

The experimental stocks were raised from seeds collected from one of the plantations near Chickballapur. *Acacia farnesiana*, nine months old was the host provided to the sandal after about eight weeks' growth. The plants were grown in a cast iron pot with an internal diameter of 85 cms. and a depth of 40 cms. (see Plate I). The soil consisted of equal parts of red-earth, sand and well-rotted horse manure.


Buds were derived from a scion brought on 13th February, 1928 from the diseased portion of a partially spiked plant in the Uttarahalli area. This plant showed the external symptoms of spike about the middle of February. The scion was about 1 cm. in diameter and 30 cms. long. The scion was first used for a bottle-graft on a different stock and in the course of four days commenced to send forth the dormant buds on both sides of the point of grafting. On 20th February, one of the buds below the grafting point was taken and budded on to one of the lower branches of the experimental stock. The usual mixture of clay and cowdung was not used to dress the wound, but only sterilised wax cloth. The terminal foliage of the branch was cut down while the rest of the plant was left undisturbed.

Four days after the operation, the bud burst and measured 1.5 mm. in height. Subsequently it began to make slow growth. The following are the dates on which various operations were carried out:—

Seed sown	...	...	7-3-1927
Germination occurred	...	...	6-4-1927
Host given	...	...	30-4-1927
Transplanted	...	...	1-9-1927
Budded	...	...	20-2-1928
First photograph taken	...	...	20-4-1928
Symptoms of spike observed on stock	...	...	1-7-1928

As a control, a healthy bud was grafted upon a healthy stock, the same technique being employed. The bud was derived from a healthy plant brought from the Uttarahalli area which was planted in mixed earth in a pot. Six days after, the dormant buds began to burst from several portions of the plant. One of these was used for the budding experiment.

The stock was a comparatively small one growing in association with an *Acacia farnesiana* plant in a flower pot. Budding was carried



out on 26th April, 1928 and the first photograph taken on 19th May, 1928. The period under experiment had the following average values for temperature and humidity in the greenhouse where all the buddings were conducted.

1928		Temperature, F.	Humidity, per cent.
February	...	79.6	54.8
March	...	81.8	44.6
April	...	85.0	53.0
May	...	...	...
June	...	78.6	66.0

Percentage of Success.—In the case of spiked buds about 10 per cent. succeeded while in the case of healthy specimens the figure was only 8 per cent. The number of spiked buds grafted was 79 and healthy buds 75. The dry heat during the early part of the period under experiment was chiefly responsible for the high mortality of the buds concerned. There is no doubt that with the advance of the season, the percentage of success will increase.

Type of Growth.—The growth of a spiked bud is entirely different from the corresponding growth of a healthy bud. The rate of growth in the case of spike is much slower than with a healthy bud. Plate V clearly shows the difference between a spiked bud which has grown for one hundred and twenty-eight days and a healthy natural bud allowed to grow for only 10 days.

The healthy leaves are broad and light green, internodes longer, and the healthy shoot does not show any tendency to branch off laterally. The growth of the spiked bud on the other hand, is quite characteristic. The leaves are small and deep green; the internodes are very short and the branching is proliferous, imparting a typical bushy appearance to the entire growth (see Plate V). The spiked shoot may also be compared with a healthy shoot budded on another stock (see plate VII). This represents a growth of only 68 days.

TRANSMISSION OF THE INFECTIVE PRINCIPLE TO THE STOCK.

The natural bud on the branch of the stock used for budding, which had been allowed to grow since 5th June, 1928, ceased growing after a fortnight. On 1st July the internodal buds began to burst, though the apical bud did not grow any further. The pale green

colour of the leaves on the shoot intensified to a dark green during the course of the week, and on 7th July the characteristic growth of spiked leaves was first observed, i.e., one hundred and thirty-one days after the bud was inserted. During the weeks that have followed the spike has taken full possession of the shoot (see Plate VIII) and thus the original suspicion has been fully confirmed. It is possible that infection might have been observed at an earlier date if the normal shoot had been allowed to grow earlier than 5th June.

SUMMARY.

Buds derived from spiked stems can be successfully budded on to healthy stocks growing in pots under controlled conditions.

This method provides us with a technique by which the transmission of spike can be effected in young sandals 6–9 months old. Controls can also be run with equal success.

The stock under experiment caught the infection 131 days after the spiked bud was inserted in the stock.

The susceptibility of sandals growing under different controlled conditions to bud infection is being investigated.

Our grateful thanks are due to Professor Norris for his helpful criticisms and the keen interest he has evinced during the course of the work.

*Department of Bio-Chemistry,
Indian Institute of Science,
Bangalore.*

[Accepted, 11-10-28.]

PRINTED AND PUBLISHED
BY GEORGE KENNETH AT THE DIOCESAN PRESS,
POST BOX 433, MADRAS-1928. C 6339.



The experimental stock growing with *Acacia farnesiana* as host. Observe the buds, healthy and spiked, enclosed in the marked rectangle and marked A and B.

PLATE II.



Spiked bud, two months after grafting.

PLATE III.



The spiked bud at a later stage, three weeks later than in Plate II.

PLATE IV.




The spiked bud, three months after grafting.



A healthy natural bud A after ten days' growth and the grafted spiked bud B after 128 days growth.

PLATE VI.



Control experiment with a healthy bud. Photographed three weeks after grafting.



Control experiment. The healthy bud $2\frac{1}{2}$ months after grafting.

PLATE VIII.



Compare Plate V. The normal bud A has now become infected.

- Part IV.—The Constitution of the Acid formed by the action of Sulphuric Acid on Camphorquinone. By Madhav Balaji Bhagvat and John Lionel Simonsen. Price, As. 8.
- Part V.—*iso*Erucic Acid. By Tahilram Jethanand Mirchandani and John Lionel Simonsen. Price, As. 8.
- Part VI.—Note on the Sorting, Classification and Briquetting of Chrome and Manganese Ores. By K. R. Krishnaswami. Price, As. 8.
- Part VII. Esterification in Mixed Solvents. By B. V. Bhide and H. E. Watson. Price, As. 8.
- Part VIII.—Nitrogen-Fixation by 'Azotobacter Chroococcum'. By S. Ranganathan and Roland V. Norris. Price, Rs. 1-4.
- Part IX.—Studies in intensive Bacterial Oxidation. The Oxidation of Ammonia to Nitric Acid, Parts I—IV. By Gilbert J. Fowler, Y. N. Kotwal, Roland V. Norris, S. Ranganathan and M. B. Roy. Price, Re. 1.
- Part X.—The Decomposition of Alkaline Earth Sulphates. By L. A. Bhatt and H. E. Watson. Price, As. 12.

VOLUME X (B)—1927.

- Part I.—Tests on Suspension Insulators after Ten Years' Service. By A. S. Venkateswaran and U. Ganguly. Price, Re. 1.
- Part II.—The Measurement of Voltage Gradient on a String of Suspension Insulators. By G. Yoganandam and R. K. Sen. Price, Re. 1.
- Part III.—Intensity Variations of Madras (Fort) Radio Station. By K. Sreenivasan. Price, As. 8.
- Part IV.—Suspension Insulator Testing. By G. Yoganandam. Price, As. 8.

VOLUME XI (A)—1928.

- Part I.—I. Studies on Invertase, Part I. Preparation and Purification of the Enzyme. By B. N. Sastri and Roland V. Norris. II. Note on a Simple Method for concentrating Enzyme Solutions. By B. N. Sastri. Price, Re. 1.
- Part II.—The Bleaching of Lac. By M. Venugopalan. Price, As. 8.
- Part III.—Contributions to the Study of Spike-Disease of Sandal (*Santalum Album*, Linn.). Part I. Diastatic Activity of the Leaves. By M. Sreenivasaya and B. N. Sastri. Price, As. 12.
- Part IV.—A Micro-Method for the Determination of Enzyme Activity. By B. N. Sastri and M. Sreenivasaya. Price, As. 12.
- Part V.—The Dielectric Constants of Ammonia, Phosphine and Arsine. By H. E. Watson. Price, Re. 1.
- Part VI.—The Fermentation of Toddy and an Account of the Micro-Organisms producing it. By M. Damodaran. Price, As. 12.
- Part VII.—I. Estimation of Iodine in Soils. By Roland V. Norris and D. A. Rama Rao. II. Carbohydrate Changes during the Ripening of Plantains. By S. Ranganathan (Junior). Price, As. 12.
- Part VIII.—I. The Oxidation of Sulphur in Suspensions of Activated Sludge and its Influence on the Solubilisation of Mineral Phosphates. By C. V. Ramaswami Ayyar, T. S. S. Perumal and Roland V. Norris. II. Studies in the Proteins of Indian Foodstuffs. Part I. The Proteins of Ragi (*Eleusine coracana*): Eleusin, the Alcohol-Soluble Protein. By Nuggehalli Narayana and Roland V. Norris. Price, Re. 1.

- Part IX.—I. Contributions to the Study of Spike Disease of Sandal (*Santalum album*, Linn.). Part II. Analysis of Leaves from Healthy and Spiked Trees. By A. V. Varadaraja Iyengar. II. Contributions to the Study of Spike Disease of Sandal (*Santalum album*, Linn.). Part III. Physico-Chemical Study of the Leaf-Sap. By A. V. Varadaraja Iyengar. Price, Re. 1.
- Part X.—I. Studies on Soil Protozoa. Part I. Protozoan Fauna of Some Mysore Soils. By H. S. Madhava Rao. II. Studies on Soil Protozoa. Part II. The Function of Mitochondria in Some Soil Protozoa. By H. S. Madhava Rao. Price, Re. 1.
- Part XI.—I. Studies in Enzyme Action. Part I. Amylase from Cholam (*Sorghum vulgare*). By V. N. Patwardhan and Roland V. Norris. II. Studies in Enzyme Action. Part II. The Nature of Amylase. By D. Narayana-murti and Roland V. Norris. Price, Rs. 1-8.
- Part XII.—A Bio-Chemical Study of some Soil Fungi with special reference to Ammonia Production. By A. K. Thakur and Roland V. Norris. Price, Re. 1.
- Part XIII.—The Atomic Weight of Antimony from Different Sources. By K. R. Krishnaswami. Price, Re. 1.
- Part XIV.—Oil from the Seeds of *Adenanthera Pavonina*. A Source of Lignoceric Acid. By S. M. Mudbidri, P. Ramaswami Ayyar and H. E. Watson. Price, As. 8.
- Part XV.—I. Constituents of some Indian Essential Oils. Part XXII. The Essential Oil from the Flower Heads of *Cymbopogon Coloratus*, Stapf. By P. P. Pillay, B. Sanjiva Rao and J. L. Simonsen. Part XXIII. The Essential Oil from the Fruits of *Piper Cubeba*, Linn. By B. Sanjiva Rao, V. P. Shintre and J. L. Simonsen. Part XXIV. The Essential Oil from the Rhizomes of *Curcuma Zedoaria*, Roscoe. By B. Sanjiva Rao, V. P. Shintre and J. L. Simonsen. II. The Constituents of Indian Turpentine from *Pinus Longifolia*, Roxb. Part IV. By P. Parameswaran Pillay and J. L. Simonsen. Price, Rs. 1-4.
- Part XVI.—I. α -isoPropylglutaconic Acid. By K. V. Hariharan, K. N. Menon and J. L. Simonsen. II. Derivatives of Methyl 2:2-Dimethylcyclo-Pentan-3-one-1-Carboxylate. By C. S. Gibson, K. V. Hariharan and J. L. Simonsen. Price, As. 12.
- Part XVII.—I. Thiophthalic Acids. Part I. By Gopâl Chandra Chakravarti. II. Organic Cyclic Polysulphides: Condensation of Ethylene Mercaptan with Di- and Trichloroacetic Acids. By Gopâl Chandra Chakravarti and Jogendra Mohan Saha. Price, As. 8.
- Part XVIII.—Bis-Semidine Inversion in Aromatic Dihydrazo-Compounds. By Praphulla Chandra Guha and Hirendra Kumar Banerjee. Price, As. 8.
- Part XIX.—I. Contributions to the Study of Spike-Disease of Sandal (*Santalum Album*, Linn.). Part IV. Chemical Composition of Healthy and Spiked Sandal Stems. By D. A. Rama Rao and M. Sreenivasaya. II. Contributions to the Study of Spike Disease of Sandal (*Santalum Album*, Linn.). Part V. Transmission of Spike by Budding. By M. Sreenivasaya and G. Gopalaswami Naidu. Price, Re. 1.