

Verified
M
Journal of the Indian Institute
of Science.

V. 16 A Part 12. 1933.

Pp 129-138

32

A. d. m. 2, N. 14. 15
N 33. 16A. 12

196396

956

956

Price, As. 12.]


[Vol. 16A, Part XII, pp. 129 to 138.]

JOURNAL

OF THE

Indian Institute of Science

CONTENTS



SYNTHESIS OF PROTEINS IN PLANTS.

PART I. CONVERSION OF NITRATES INTO PROTEIN IN
HELIANTHUS ANNUS, LINN.

BY

K. S. Varadachar.

JOURNAL
OF THE
INDIAN INSTITUTE OF SCIENCE

Volume 1 included 21 issues during 1914-18. Thereafter one volume was published annually until 1925, when Volumes 8 (A) and (B), followed by 9 (A) and (B) (1926) and 10 (A) and (B) (1927) appeared.

VOLUME 11 (A)—1928.

Parts I—XIX.

VOLUME 12 (A)—1929.

Parts I—XX.

VOLUME 13 (A)—1930.

Parts I—XV.

VOLUME 14 (A)—1931.

Part I.—Studies on the chemical composition and physical properties of plant tissue fluids.
Part I. Effects of age and environment on the tissue fluids of French beans (*Phaseolus Vulgaris*). By C. Narasimha Acharya and B. N. Sastri. Price, As. 12.

Part II.—The Photochemical Oxidation of Organic Hydrocarbons. Part II. Toluene, and the Oxidation of Benzaldehyde. By C. J. Kothari and H. E. Watson. Price Re. 1-4.

Part III.—I. Cantharidin from *Mylabris Pustulata* Fb., India. By B. H. Iyer and P. C. Guha.
II. The Fatty acids from oil of Cantharis (*Mylabris Pustulata* Fb., India) By B. H. Iyer and P. Ramaswami Ayyar. Price, Re. 1.

Part IV.—I. Amylase from Rice. II. Dialysis of Some Cereal Amylases. By Vinayak Narayan Patwardhan and Dattatreya Vishnu Karmarkar. Price, As. 12.

Part V.—Contributions to the Study of Spike-Disease of Sandal (*Santalum Album*. Linn.),
Part XII. Ecology of Sandal. By M. Sreenivasaya and S. Rangaswami. Price, As. 8.

Part VI.—Studies in Electro-Culture. Part I. Introduction. By K. Venkata Giri, T. J. Mirchandani and V. Subrahmanyam. Studies in Electro-Culture. Part II. Influence of Electrical Treatment on the Germination of Barley and the Diastatic Activity of Malt. By K. Venkata Giri and V. Subrahmanyam. Price, Rs. 2.

Part VII.—The Origin and Nature of the Peaty Soils of Travancore. By T. R. Narayana Pillai and V. Subrahmanyam. Price, Re. 1.

Part VIII.—The Relation between Seeds and Micro-Organisms. By T. R. Sathe and V. Subrahmanyam. Price, Rs. 1-4.

Part IX.—The Reactivity of Dimethyldihydroresorcin. Part I. Condensation with Aromatic Aldehydes. By Gopál Chandra Chakravarti, Haripada Chattopadhyaya and Prafulla Chandra Ghosh. Price, Re. 1.

Part X.—The Reactivity of Dimethyldihydroresorcin. Part II. Behaviour towards Ortho-Nitro- and Ortho-Aminobenzaldehydes. By B. H. Iyer and G. C. Chakravarti. Price Re. 1.

Price, As. 12.]

[Vol. 16A, Part XII, pp. 129 to 138.]

JOURNAL

OF THE

Indian Institute of Science


CONTENTS



SYNTHESIS OF PROTEINS IN PLANTS.
PART CONVERSION OF NITRATES INTO PROTEIN IN
***HELIANTHUS ANNUS*, LINN.**

BY

K. S. Varadachar.



SYNTHESIS OF PROTEINS IN PLANTS. PART I. CONVERSION OF NITRATES INTO PROTEIN IN *HELIANTHUS ANNUS*, LINN.

By K. S. VARADACHAR.

Although the study of synthesis of proteins in plants has drawn the attention of a large number of workers, yet very little definite information is available about this important branch of plant physiology. This is chiefly due to want of adequate technique for controlling the intake of nitrogen by plants as also to lack of accurate methods for estimating small quantities of nitrogenous substances formed at various stages in the plant.

A number of useful observations have been recorded by individual workers but the uncertainty regarding the first product of photosynthesis has invariably led to a good deal of confusion in interpreting these observations. At various times hydrocyanic acid, ammonia or potassium nitrite has been mentioned as reacting with formaldehyde and thence the mechanism of protein-synthesis elaborated. The fact that protein-synthesis is not dependent on light but can proceed even in the dark has been a great drawback in these explanations. Several theories have also been postulated on the basis of observations made under *in vitro* conditions but most of them have only academic interest.

The theories regarding the mechanism of protein-synthesis can be broadly divided into three groups—(1) those that assume the preliminary formation of the several amino acids and their subsequent condensation to proteins, (Trier, G. *Einfachen-flanzenbasen und ihre Beziehungen zum Aufbau der Eiweisstoff und Lecithin*, Berlin, 1912; Baudisch u. Meyer, *Central Bakt*; Abt. II. 1912, **45**, 1771; Baudisch u. Klinger, *Ibid.*, 1916, **49**, 1167; Lob, *Zeit. f. Elektrochem.*, 1906, **12**, 282), (2) those that assume direct interaction between the sugars formed during photosynthesis and potassium nitrite or ammonia arising out of the decomposition of absorbed nitrate, resulting in the synthesis of protein in plant (Baly *et al.*, *J.C.S.*, 1922, **121**, 1078; Huppert, *Konstitution und Konifiguration der Eiweisstoff*, Leipzig, 1925; Loew, *Biochem. Z.*, 1912, **41**, 224; *Chem. Ztg.*, 1912, No. 7 and *Ber.*, 1913, **46**, 684), and (3) those that assume a preliminary formation of amides, especially asparagine and their subsequent transformation into proteins (Muller, *Landwirtschaft Versuchs-St.*, 1886, **33**, 311; Nakamura, *Bull.*

TL
A. d. m. 2, N14 I 2
N33 16A 12

Agric. Coll. Tokyo, 1897, 465; Saposchnikow, *Bot. Zent.*, 1895, **63**, 246; Prinanischnikoff, *Russ. J. Expt. landwirt*, 1912, **13**, 5; *Rev. gen. Bot.* 1913, **25**, 5).

The object of the present investigation is to throw some fresh light on this important problem by directly feeding the plant with various nitrogenous constituents and following the attendant biochemical changes. The present paper relates to the changes undergone by nitrates in the sunflower plant (*Helianthus annuus*, Linn.).

EXPERIMENTAL.


The experiments were conducted with sand-cultured plants in glazed earthenware pots. The sand was repeatedly washed with water and, after drying, filled in at the rate of 60 lbs. per pot. It was then mixed with lime at 50 g. per pot and left to weather for about 4-5 weeks, after which potassium acid phosphate (4.8 g.) and potassium sulphate (2.7 g.) were added.

The sunflower plants (Sutton's Giant Yellow variety) were raised on manured beds until the stems attained a thickness of about 8 mm. and then transplanted into the pots. The potted plants were watered at the rate of 2 litres for every alternate day.

After about 15 days when the plants had established themselves, they were injected with distilled water and their ability to take in aqueous nutrients tested from day to day. The technique of injection developed after numerous trials and adopted for this investigation was as follows :—

A glass tube about 1 cm. in diameter and 10 cms. long was fused to another one of diameter 3 mm. The smaller tube was bent twice at right angles. To the end of the smaller tube was attached one inch of rubber tubing and to this was fixed an 'Esco' record syringe needle of stainless steel. To inject a plant, the needle was inserted slantwise into the stem as near the soil as possible so that it was well fixed in the plant. It was then removed and the bore cleaned with a brass wire and reinserted. The point at which the injection was made was protected with a little beeswax. Distilled water was placed in the side tube which acted as a reservoir and observed for a couple of days to ensure that the uptake was continuous and that there was no choking in the needle. When required, the distilled water was pipetted out and the experimental solution introduced in known quantities. The

tube was held in position by means of a zinc clip attached to an iron rod and a test tube inverted over the reservoir to minimise evaporation.



It was observed that almost all the plants took in water quite readily and continued to grow healthy though the leaves soon began to turn yellow owing to lack of nitrogen.

After ensuring that the passage of water into the plant was steady and continuous, the distilled water was pipetted out and known quantities of standard nitrate solution (2.5 per cent.) placed in each of the reservoir tubes. Several plants were left with water only to serve as controls.

The above treatment with nitrate solution was continued for about a week. Specimens of whole plants were collected at intervals, biological activity stopped by heating for 4 hours at 100° and drying completed at 60°. The dried specimens were analysed for total weight, total nitrogen including nitrates, humin N, amide N, basic N, and non-basic N. The quantity of nitrate absorbed by each plant at the time of observation was also determined by the difference between the quantities introduced at the beginning and that left unabsorbed when sampling.

A preliminary trial was conducted in July 1930 and subsequently repeated in greater detail in February 1931 and 1932 respectively. It was observed that the plants recovered their normal colour within 48 hours and remained healthy throughout the experiment. The results show that the injection method of feeding plants is a highly efficient form of introducing large quantities of nutrients into plants without any detrimental effect.

The total nitrogen was found according to the Arnold-Gunning modification of the Kjeldahl method. Considerable difficulty was experienced in selecting suitable methods for estimating small quantities of the different forms of nitrogen in plant material, owing to the amount of material available for analysis being small. Thus the estimation of nitrates presented great difficulty and several standard methods were tried and found unreliable. The most consistent results were obtained by repeated extraction with hot water followed by reduction with Devarda's alloy.

In regard to the distribution of nitrogen in the plant, the method of differentiating it in terms of solubility in various solvents like water, alcohol and sodium chloride would not lead to any conclusive figures. Van Slyke's method of analysis is not applicable to a material like dried plant. In view of the above the micro method of determining Hausmann numbers developed by Thiemann (*Biochem. J.*, 1926, 20, 1190) was employed. The material was dropped into boiling 20 per cent. hydrochloric acid and hydrolysed for 36 hours. The hydrolysate was then separated into humin, amide, basic and non-basic fractions. The nitrate content of each sample was also determined.

A large mass of useful data were collected during the three seasons in which the experiments were conducted but with a view to facilitating discussion only the more representative figures have been given in the following tables:—

TABLE I.
Effect of nitrate injection on the dry weights of plants.

No. of days under injection	TREATED PLANTS		CONTROL PLANTS	Percentage increase in dry weight over controls
	Nitrate absorbed in mg.	Dry weight in gms.	Dry weight in gms.	
1	8.5	11.78	10.8	+ 10.0
2	14.9	7.51	6.42	+ 16.9
3	6.4	6.00	7.73	- 23.0
4	7.9	11.21	11.85	- 5.4
5	11.1	11.27	9.09	+ 23.8
6	16.9	9.66	6.60	+ 46.3
7	13.2	13.28	8.40	+ 58.0
8	4.1	11.02	9.69	+ 13.4
11	3.6	12.45	6.69	+ 86.4
12	7.3	10.11	7.05	+ 43.4
13	5.7	8.27	9.91	- 17.0
14	6.5	10.61	7.51	+ 41.3
15	9.8	12.65	7.44	+ 70.0
17	?	10.82	7.71	+ 40.2
18	8.1	12.18	9.79	+ 24.5
19	10.4	11.63	8.44	+ 37.9
20	?	17.63	11.24	+ 57.0
21	11.5	13.02	9.59	+ 35.5
22	12.6	13.71	11.20	+ 22.3
23	9.4	9.58	9.14	+ 5.8
24	0.2	12.25	7.87	+ 55.5
25	9.9	13.82	10.47	+ 32.0
26	11.7	16.39	12.74	+ 28.7
27	5.9	9.59	9.65	- 0.7
28	9.7	11.74	10.27	+ 14.3

Average increase in dry weight - 25.6 per cent.

TABLE II.

Season—1931.

Number of days under injection	Total nitrogen in plant	Nitrate nitrogen in plant
	mgms.	mgms.
2	67.1	10.7
4	92.5	4.2
5	112.6	10.6
7	105.4	13.2
11	205.2	26.4
12	126.9	1.9
15	122.6	7.2
18	99.5	2.3
20	233.2	19.2
21	101.5	5.5

TABLE III (a).

Season—1930.

Date of commencement of injection, 25-8-1930.

Number of days under injection	Total nitrogen per cent.	Humin nitrogen per cent.	Percentage of humin N. in total N.
4	0.79	0.33	41.0
9	0.94	0.40	43.2
15	0.83	0.34	41.0

TABLE III (b).

Season—1931.

Number of days under injection	Percentages on dry weight			
	Humin N.	Amide N.	Diamino N.	Non-basic N.
2	0.11	0.10	0.15	0.40
4	0.11	0.06	0.10	0.53
5	0.12	0.06	0.10	0.67
6	0.14	?	0.13	0.61
7	0.12	0.05	0.13	0.39
8	0.11	0.01	0.13	0.50
11	0.12	0.15	0.11	1.77
12	0.12	0.08	?	0.91
13	0.13	0.05	0.13	0.45
15	0.14	0.05	0.09	0.67
17	0.13	0.04	0.14	0.42
19	0.12	?	0.11	0.57
20	0.16	0.05	0.09	0.51
22	0.11	0.09	?	0.65
23	0.13	0.08	0.16	0.45
25	0.12	0.06	?	0.47
26	0.13	0.07	0.14	0.88
27	0.14	0.08	0.12	1.77

The effect of nitrate injection on the dry weights of plants was determined as follows: pairs of plants which were very nearly alike with regard to height, girth and general development at the time of the commencement of the experiment were taken out at stated intervals and their respective dry weights compared. Since the specimens were independent plants and had absorbed varying amounts of nitrate, the quantitative effect of the treatment with time could not be evaluated. The plants were therefore treated as representatives of two sets of population, one subjected to injection and the other left untreated and the effects on corresponding specimens compared. The results (Table I) show that the treatment has been beneficial and that the majority of

plants have shown distinct increase in dry weight as the result of nitrate injection. The above observation being perhaps the first record of plant growth being favoured by direct feeding with a mineral nutrient, the experiments were repeated carefully during successive seasons and the results confirmed.

The absorption of nitrate (as shown in column 2, table I) being a mechanical process dependent on such variable factors as temperature, humidity, transpiration etc. does not show any regularity. The results show, all the same, that the plants have absorbed useful amounts of nitrogen and have grown normally without, in any way, being affected by either the injection or the high concentration of the nutrient fed into it.

From column 3 of table II, it might be seen that there is increase in the nitrate content upto the 11th day followed by during successive days. There is however no regularity about this accumulation nor does the quantity bear any proportion to the total nitrogen in the plant thereby suggesting that there is continuous transformation of nitrate into other forms of nitrogen.

The large difference with regard to the humin nitrogen obtained during the 1930 and 1931 series is quite striking. In the former, the humin [table III (a), column 3] accounted for about 40 per cent. of the total nitrogen while in the latter [table III (b), column 2] it was less than a third of that quantity. The difference between the two sets of figures is quite inexplicable particularly when considering that the hydrolyses were conducted under identical conditions. Such high figures as those obtained in the first season were not repeated at any later stage but even those of the latter ones should be regarded as excessive so that the accuracy of the distribution of nitrogen has to a large extent been affected by the precipitation of quite considerable quantities in the form of the comparatively unknown entity humin.

The variation observed with regard to the figures for amides [column 3, table III (b)] suggest the formation of those compounds as intermediary products, in the building up of proteins. This supports one of the existing theories (*loc. cit.*) regarding the mechanism of protein synthesis. Whether this is due to the formation of asparagine or any other amide, it is not possible to say at this stage of the investigation. It is hoped that later research will throw more light on this aspect of the problem.

The figures for diamino nitrogen [table III (b), column 4] representing the combined values for arginine, histidine, lysine and cystine

fractions do not lead to any conclusion. Fractionation of the constituents by micro methods is now in progress and will form the subject of a later communication.

The figures for non-basic nitrogen (column 5) show a correlation with those for amide. The rise and fall correspond inversely to similar alteration in the amide value showing a relation between those two classes of substances. Thus, from table III (b), it is clear that the transformation of nitrate in the plant is accompanied by periodic accumulation and disappearance of amides and corresponding reverse changes in the non-basic fractions.

With a view to shortening the interval between the observations the whole series of experiments were repeated in 1932 sampling the plants day and night at intervals of two hours. The first sample was taken 2 hours after the injection of nitrate and the last 50 hours later. The figures for the analysis of the various samples collected at six-hour intervals are given in table IV.

TABLE IV.

Time of sampling hrs.	Total dry weight of plant in gms.	Total N in whole sample in mg.	Nitrate N in whole sample in mg.	Humin N	Amid N	Diamino N	Non-basic N
0	14.62	118.2	0.6	0.16	0.13	0.16	0.41
0	14.08	124.9	2.8	0.16	0.12	0.14	0.32
6	19.05	144.0	0.8	0.15	0.13	0.13	0.35
12	16.05	202.7	1.3	0.19	0.12	0.16	0.79
18	18.94	...	3.7	0.12	0.09	0.12	0.52
24	17.66	...	2.1	0.17	0.16	...	0.66
30	17.24	159.1	0.7	0.16	0.11	0.17	0.48
36	16.20	372.6	4.5	0.30	0.19	0.37	1.40
42	37.65	349.7	1.5	0.14	0.09	0.13	0.57

It may be seen from the above that the nitrate content in the plant remains comparatively small throughout the period of observation. The total nitrogen does however show a tendency to increase, thereby indicating that conversion of the nitrogen into other forms takes place at a very rapid rate so that nitrate content is not high at any stage during the experiment.

The amide nitrogen under observation shows the same type of rise and fall as that observed before thereby indicating that the nitrate passes through the amide stage. But there is no correlation between the amide content of the plant and the figures for non-basic nitrogen as was noted before.

Although the foregoing observations have not revealed any striking feature in the mechanism of synthesis of proteins in plants they have yet shown that the technique of direct feeding offers immense possibilities for future study. Further work is therefore in progress with a view to refining the technique and standardising the conditions for estimation of the different forms of nitrogen.

SUMMARY.

A direct method of feeding of plants with nutrients is described. This technique has been applied to the study of protein synthesis by feeding nitrogen-starved sunflower plants with potassium nitrate solution and studying the attendant changes.

There is evidence to suggest that there is continuous conversion of nitrate into other forms of nitrogen. The added nitrogen would appear to pass through the amide stage before being converted into protein.


In conclusion the author wishes to express his thanks to Dr. V. Subrahmanyam, D.Sc., F.I.C., for suggestions and criticism.

[Accepted, 24-7-1933.]

Department of Biochemistry,

Indian Institute of Science,

Bangalore.



3L

A, d, m 2, N 14 13

N 32.16A.12

VOLUME 15 (A)—1932.

- Part I.—Salt Stains on South Indian Hides and Skins. By V. N. Patwardhan and M. Subramania Sastry. Price, As. 12.
- Part II.—Chemical Examination of the Root-Bark of *Plumbago Rosea*, Linn. By M. C. Tummin Katti and V. N. Patwardhan. Price, As. 8.
- Part III.—Dilatometric Studies in Enzyme Action. Part II. Contraction Constants of Enzyme-Substrate Reactions. By M. Sreenivasaya and H. B. Sreerangachar. Price, As. 12.
- Part IV.—Aryl Di- and Poly-Stibinic Acids, Distibinous Oxides and Distibino-Compounds. By Sohrab M. Mistry and Praphulla Chandra Guha. Price, Re. 1.
- Part V.—Bhadravati Wood-Tar and its Utilization. By Y. K. Raghunatha Rao, B. Sanjiva Rao and H. E. Watson. Price, Re. 1.
- Part VI.—The Equilibrium between Dimethyl Ether, Methyl Alcohol and Water. By N. G. Gajendragad, S. K. Kulkarni Jatkar and H. E. Watson. Price, Re. 1.
- Part VII.—Studies in Indian Essential Oils. I. Essential Oil from Leaves of *Litsæ Zeylanica* Linn. By B. Sanjiva Rao. II. Essential Oil from Flower-Heads and Stalks of *Andropogon Kuntzeanus*, Hack, *Var. Foveolata*, Hack. By B. Sanjiva Rao. III. Essential Oil from Leaves of *Thymus Serpyllum*, Linn. By Jagjit Singh and B. Sanjiva Rao. IV. Essential Oil from Leaves of *Cinnamomum Zeylanicum*, Breyn. By (the late) V. P. Shintre and B. Sanjiva Rao. Price, Re. 1.
- Part VIII.—Organic Manure from Sewage, Town Refuse and Waste Vegetation. By J. Jagannatha Rao and V. Subrahmanyam. Price, Rs. 1-8.
- Part IX.—I. Studies in Enzyme Action. Part V. Ageing of Amylases in Aqueous Solution. By K. Venkata Giri and V. Subrahmanyam. II. Studies in Enzyme Action. Part VI. Heat Inactivation of Pancreatic Amylase. By K. Venkata Giri. III. Studies in the Proteins of Indian Foodstuffs. Part IV. The Proteins of Fenugreek (*Trigonella Foenum Graecum*). By Y. V. Srinivasa Rau and M. Sreenivasaya. Price Rs. 1-8.
- Part X.—Studies in Bridge Formation. Part I. Attempts to Synthesise Bicyclic Terpene Derivatives by a New Method. By P. P. Patel and P. C. Guha. Part II. Formation and Stability of a Bridged *cycloHexenone*, 1-Dimethylethane—^{1:2:3} III ^{2:3:4} *cyclobutane-2-one*. By P. S. Mayuranathan and P. C. Guha. Price, Re. 1.

VOLUME 15 (B)—1932.

- Part I.—A New Permeameter. By K. V. Karantha. Price, Rs. 1-8.
- Part II.—Measurement of Power Factor and Loss in Dielectrics. By T. J. Mirchandani, G. Yoganandam, S. K. Roy and N. V. Narayanaswami. Price, Rs. 1-4.
- Part III.—Transients in Negative Constant Series Circuits. By L. C. Verman. Price, Rs. 1-4.
- Part IV.—Telephony by Carrier and One Side Band. By S. P. Chakravarti. Price, Re. 1.

VOLUME 16 (A)—1933.

- Part I.—Rhaponticin, and Anthraquinone Derivatives from *Rheum Emodi*, Wall. (Indian, or Himalayan Rhubarb). By M. Ghouse Mohiuddin and M. C. Tummin Katti. Price, As. 12.
- Part II.—Constitution of the So-Called Dithiourazole of Martin Freund. Part V. Isomerism of Hydrazodithiodicarbonamides, Iminothiolthiobiazoles and Di-R-Iminothiobiazoles. By Shaha L. Janniah and P. C. Guha. Part VI. Isomerism of Hydrazomonothiodicarbonamides, Iminothiobiazolones and Monothiourazoles. By S. L. Janniah and P. C. Guha. Price, Re. 1-4.

- Part III.—Indian Medicinal Plants. Part I. Withania Somnifera. Part II. Swertia Chirreta. By D. N. Majumdar and P. C. Guha. Price, As. 12.
- Part IV.—I. The Colouring Constituents of Alkanet Root (*Anchusa Tinctoria*, Lam.). Part I. The Constitution of Alkannin. By Mangesh V. Betrabet and Gopāl Chandra Chakravarti. II. Note on the Constituents of the Wax from Alkanet Root. By M. V. Betrabet and G. C. Chakravarti. Price, Re. 1.
- Part V.—The Velocity of Alcoholysis of Esters of Dibasic Acids. By Z. H. Patel and H. E. Watson. Price, As. 12.
- Part VI.—I. Dilatometric Studies in Enzyme Action. Part III. Contraction Constants of Enzyme-Substrate Reactions. By H. B. Sreerangachar and M. Sreenivasaya. II. Physiological Products of the Lac Insect. Part I. A Preliminary Investigation. By N. K. Ranga Rao and M. Sreenivasaya. Price, Re. 1.
- Part VII.—I. Studies in the Proteins of Indian Foodstuffs, Part V. The Alcohol Soluble Protein of Fenugreek (*Trigonella Foenum Graecum*). By Y. V. Sreenivasa Rao, B. N. Sastri and N. Narayana. II. The Mucilage of Fenugreek (*Trigonella Foenum Graecum*). By C. R. Harihara Iyer and B. N. Sastri. Price, As. 8.
- Part VIII.—I. Contributions to the study of the Spike Disease of Sandal (*Santalum Album*, Linn.), Part XIII. Investigation of the Hexone Bases. By Y. V. Sreenivasa Rao. II. Contributions to the Study of the Spike-Disease of Sandal (*Santalum Album*, Linn.), Part XIV. Study of Mosaics Associated with Spiked Areas. By Y. V. Sreenivasa Rao. Price, As. 8.
- Part IX.—Physiological Products of the Lac Insect. Part II. Investigation of the Water Soluble Nitrogenous Constituents. By N. K. Ranga Rao. Price, As. 8.
- Part X.—Extension of Michael's Reaction. Part III. By Tejendra Nath Ghosh and Praphulla Chandra Guha, Price, As. 12.
- Part XI.—Studies in Bridge Formation. Part III. The Influence of the Methyl and Carbethoxy-groups in the Internal Dehydration of *gem*-Disubstituted Dihydroresorcinol Esters. Part IV. Tautomerism in isophorone. By P. S. Mayuranathan. Price, Re. 1.
- Part XII.—Synthesis of Proteins in Plants. Part I. Conversion of Nitrates into Protein in *Helianthus Annus*, Linn. By K. S. Varadachar. Price, As. 12.

VOLUME 16 (B)—1933.

- Part I.—Taylor's Frequency Tripler. By P. Rajagopal Pillai and F. N. Mowdawalla. Price, Re. 1.