

Va-piel
12

Journal of the Indian Institute
of Science

V.16 A Part 14, 1933

Pp 153-165

JL

A.d.m.2, NIVIS

N33.16A.14

196348

280


280

Price, Re. 1.]

[Vol. 16A, Part XIV, pp. 153 to 165.]

JOURNAL
OF THE
Indian Institute of Science

CONTENTS



THE SOLUBILITY OF SILVER CHLORIDE

BY

P. C. Dave and K. R. Krishnaswami

JOURNAL
OF THE
INDIAN INSTITUTE OF SCIENCE

Volume 1 included 21 issues during 1914-18. Thereafter one volume was published annually until 1925, when Volumes 8 (A) and (B), followed by 9 (A) and (B) (1926) and 10 (A) and (B) (1927) appeared.

VOLUME 11 (A)—1928.

Parts I—XIX.

VOLUME 12 (A)—1929.

Parts I—XX.

VOLUME 13 (A)—1930.

Parts I—XV.

VOLUME 14 (A)—1931.

Part I.—Studies on the chemical composition and physical properties of plant tissue fluids.
Part I. Effects of age and environment on the tissue fluids of French beans (*Phaseolus Vulgaris*). By C. Narasimha Acharya and B. N. Sastri. Price, As. 12.

Part II.—The Photochemical Oxidation of Organic Hydrocarbons. Part II. Toluene, and the Oxidation of Benzaldehyde. By C. J. Kothari and H. E. Watson. Price Re. 1-4.

Part III.—I. Cantharidin from *Mylabris Pustulata* Fb., India. By B. H. Iyer and P. C. Guha.
II. The Fatty acids from oil of Cantharis (*Mylabris Pustulata* Fb., India) By B. H. Iyer and P. Ramaswami Ayyar. Price, Re. 1.

Part IV.—I. Amylase from Rice. II. Dialysis of Some Cereal Amylases. By Vinayak Narayan Patwardhan and Dattatreya Vishnu Karmarkar. Price, As. 12.

Part V.—Contributions to the Study of Spike-Disease of Sandal (*Santalum Album*. Linn.),
Part XII. Ecology of Sandal. By M. Sreenivasaya and S. Rangaswami. Price, As. 8.

Part VI.—Studies in Electro-Culture. Part I. Introduction. By K. Venkata Giri, T. J. Mirchandani and V. Subrahmanyam. Studies in Electro-Culture. Part II. Influence of Electrical Treatment on the Germination of Barley and the Diastatic Activity of Malt. By K. Venkata Giri and V. Subrahmanyam. Price, Rs. 2.

Part VII.—The Origin and Nature of the Peaty Soils of Travancore. By T. R. Narayana Pillai and V. Subrahmanyam. Price, Re. 1.

Part VIII.—The Relation between Seeds and Micro-Organisms. By T. R. Sathe and V. Subrahmanyam. Price, Rs. 1-4.

Part IX.—The Reactivity of Dimethyldihydroresorcin. Part I. Condensation with Aromatic Aldehydes. By Gopāl Chandra Chakravarti, Haripada Chattopadhyaya and Prafulla Chandra Ghosh. Price, Re. 1.

Part X.—The Reactivity of Dimethyldihydroresorcin. Part II. Behaviour towards Ortho-Nitro- and Ortho-Aminobenzaldehydes. By B. H. Iyer and G. C. Chakravarti. Price, Re. 1.

Price, Re. 1.]


[Vol. 16A, Part XIV, pp. 153 to 165.]

JOURNAL

OF THE

Indian Institute of Science



CONTENTS



THE SOLUBILITY OF SILVER CHLORIDE

BY

P. C. Dave and K. R. Krishnaswami



THE SOLUBILITY OF SILVER CHLORIDE

By P. C. DAVE and K. R. KRISHNASWAMI.

The solubility of silver chloride in water, which at 25° is of the order of 2 milligrams per litre, is sufficiently great to cause appreciable errors in accurate analytical work if not allowed for, while in atomic weight determinations involving the precipitation of silver chloride, a knowledge of the solubility is essential. The relation between the solubility determined by the conductivity method and by other means is also of theoretical importance and it is therefore not surprising to find that numerous papers on the subject have been published. Of these the majority refer to the solubility in water, and very little information is available regarding the solubility in nitric acid or salt solutions in which silver chloride is usually precipitated.

The available data for water are shown in Table I, the solubility being expressed in moles. per litre $\times 10^3$. For the purpose of comparison, the values at 25° are given in column 6; these when not recorded directly, have been obtained either by interpolation from the results at other temperatures or, when these are not available, by assuming that the change in solubility in the region under consideration is the same as the one found in our experiments.

The variations in these figures are distinctly larger than might be expected and suggest that factors are present which have not been taken into consideration when presenting the results. One possibility is that some of the methods are inherently wrong or are subject to constant errors. A comparison of results obtained by the same method shows however that this is not a sufficient reason and it is much more probable that the explanation is to be found in the varying solubility of silver chloride according to its method of preparation and subsequent treatment. As long ago as 1871, Stas (*Compt. rend.*, 1871, **73**, 1000) observed that precipitated silver chloride could exist in four different forms which he designated as gelatinous, flocculent, powdery or granular. The solubility of these varied from 19mg./l. for the gelatinous variety to 0.02 mg./l. for the granular. Although the results obtained by Stas cannot be considered very accurate and his observation regarding the low solubility of the granular variety has not been confirmed, his remarks suggest that unless similar specimens are employed for solubility determinations, the values obtained will not be comparable.

Richards and Wells (*J. Amer. Chem. Soc.*, 1905, **27**, 485) observed that if silver chloride was filtered immediately after precipi-

36

A. d. 20.2.1923

N 23.16A 15

TABLE I.

Solubility of silver chloride in water

No.	Authors	Method employed	Temperature	Solubility	Solubility at 25°	Reference
1	Holleman	Conductivity	13.8 26.5	0.97 1.82	1.74	<i>Z. physikal Chem.</i> , 1893, 12 , 132.
2	Kohlrausch and Rose.	do.	2.0 10.0 18.0 26.0 34.0 42.0	0.53 0.68 1.06 1.56 2.11 2.82	1.50	<i>ibid.</i> , 1893, 12 , 242.
3	Böttger	do.	19.95	1.07	1.34	<i>ibid.</i> , 1903, 46 , 603.
4	Kohlrausch	do.	18.0	1.12	1.56	<i>ibid.</i> , 1905, 50 , 356.
5	Do.	do.	1.55 4.68 9.97 17.51 18.0 25.86 34.12	0.39 0.46 0.62 0.91 0.93 1.35 1.91	1.30	<i>ibid.</i> , 1908, 64 , 168.
6	Melcher	do.	18.0	1.05	1.42	<i>J. Amer. Chem. Soc.</i> , 1910, 32 , 50.
7a	Van Rossem	do.	18.0 25.0	1.04 1.43	1.43	<i>Chem. Weekblad.</i> , 1912, 9 , 396.
7b	Do.	E. M. F.	18.0 25.0	1.02 1.43	1.43	<i>idem.</i>
8	Goodwin	do.	25.0	1.25	1.25	<i>Z. physikal Chem.</i> , 1894, 13 , 645.
9	Thiel	do.	25.0	1.41	1.41	<i>Z. anorg. Chem.</i> , 1900, 24 , 57.
10	Abegg and Cox...	do.	25.0	1.25	1.25	<i>Z. physikal Chem.</i> , 1903, 46 , 1.
11	Pinkus and Berkolaiko.	E. M. F. and Nephelometric.	25.0	1.42	1.42	<i>J. Chim. physique</i> , 1930, 27 , 364.
12	Scott and Johnson.	Nephelometric	0.5	0.50	...	<i>J. Amer. Chem. Soc.</i> , 1930, 52 , 3586.
13	Rolla	Optical	18.0	1.20	1.57	<i>Atti. R. Acad. Lincei.</i> 1913, 22 , 2.
14	Hill	Volumetric	25.0	1.60	1.60	<i>J. Amer. Chem. Soc.</i> 1908, 30 , 74.
15	Glowczynski	do.	25.0	1.20	1.20	<i>Kolloidchem. Beih.</i> , 1914, 6 , 147.
16	Whitby	Colorimetric	21.0	1.07	1.29	<i>Z. anorg. Chem.</i> , 1910, 67 , 108.

No.	Authors	Method employed	Temperature	Solubility	Solubility at 25°	Reference
17	Popoff and Neuman.	Tyndal beam	25.0	1.278	1.278	<i>J. Physical Chem.</i> , 1930, 1853.
18	Eversole and McLachlan.	do.	25.0	1.52	1.52	<i>J. Amer. Chem. Soc.</i> , 1932, 54 , 867.
19	Neuman	do.	25.0	1.273	1.273	<i>ibid.</i> , 1932, 54 , 2195.

tation the filtrate contained 1.5 mg./l. of silver chloride, whereas if filtration was deferred for several days, the quantity in solution fell to 1.1 mg./l. The difference is almost certainly due to aggregation of the particles and this aspect of the question has been discussed by Glowczynski (*loc. cit.*) who pointed out that a decrease in the degree of dispersion would lower the solubility.

The first seven determinations given in Table I were all made by the conductivity method, but the first two of them were not carried out under favourable experimental conditions. Determinations Nos. 3 and 5 were conducted with silver chloride precipitated from cold dilute solutions and washed with cold water. The conductivity was determined at closely regulated temperatures and the solution was protected from actinic light during the experiments. Böttger's value was the mean of thirty-two separate and concordant determinations and is in good agreement with the value 1.04 interpolated from Kohlrausch's figures. Melcher and Van Rossem's determinations (Nos. 6 and 7) appear to have been made with as much care as those of Kohlrausch or Böttger, but the silver chloride employed was obtained by precipitation from boiling solutions, washing with hot water and drying. The results of Melcher are in agreement with those of Van Rossem but not with those of Böttger or Kohlrausch.

Determinations Nos. 7b, 8, 9 and 10 were carried out by the electromotive force method, but except in the case of No. 7b no details are given of the way in which the silver chloride was prepared. Abegg and Cox made a large number of determinations and obtained the value 1.25×10^{-5} M./L. (moles. per litre) identical with Goodwin's figure; Van Rossem, however, after a prolonged series of measurements arrived at the figure 1.43×10^{-5} M./L. identical with the value he had obtained by the conductivity method and in good agreement with the result obtained by Thiel. It appears highly probable that the discrepancy between the two sets of values is due to differences in the methods of preparing the silver chloride. The wide difference between the results of Glowczynski and of Hill who both made use of the thiocyanate method is probably to be ascribed to the same cause.

Scott and Johnson and also Pinkus and Berkolaiko employed the nephelometric method. The former authors described their silver chloride as 'flocculent' but as they worked only at 0.5° it is not possible to compare the results. Pinkus and Berkolaiko prepared silver chloride by the addition of a slight excess of 0.1N hydrochloric acid to silver nitrate solution and boiling the mixture. The precipitate was washed with cold water and the solubility determined within five hours of precipitation. Their results by the nephelometric and E.M.F. methods were in good agreement.

Popoff and Neuman and subsequently Neuman made several determinations by observing the concentration at which the Tyndal beam due to turbidity just made its appearance. A very high order of accuracy, viz. $1 \times 10^{-8} M./L.$ was claimed for this method, and the two sets of results are in close agreement, but the mean value 1.273×10^{-5} is at great variance with the figure 1.52×10^{-5} obtained by Eversole and McLachlan under similar experimental conditions. It therefore appears probable that the method involves some source of error which has not yet been determined.

The available data for the solubility of silver chloride in nitric acid and sodium nitrate of various concentrations are given in Table II. The concentration of nitric acid and sodium nitrate is expressed in moles. per litre and the solubility of silver chloride in $M./L \times 10^{-5}$

TABLE II.

Solubility of silver chloride in nitric acid and sodium nitrate solutions.

Glowczynski ...	...	25°	HNO ₃	0.0005	0.001	0.01	0.3
			AgCl	1.15	1.19	1.24	1.54
Eversole and McLachlan.	25°		HNO ₃	0.0238	0.119	0.238	0.52
			AgCl	1.54	1.62	1.70	2.35
Scott and Johnson	...	0.5°	HNO ₃	0.11	0.24	0.49	0.99
			AgCl	0.628	0.698	0.711	0.732
Mulder	...	15-20°	NaNO ₃	0.047	0.093	0.355	...
			AgCl	6.7	9.3	17.6	...

The results of Mulder (*Die Silberprobirmethoden, Leipzig, 1859*) are very old and the exact temperature is not specified. They serve to indicate, however, that the solubility of silver chloride is largely increased by the addition of sodium nitrate, and the values are of the same order as those we have obtained.

The solubility of silver chloride in aqueous solutions of nitrates has also been studied by Popoff and Neuman (*loc. cit.*) but their work, was confined to solutions of very low concentrations of the order

0.00001-0.01 M. as the object in view was the study of the applicability of modern theories of strong electrolytes to such solutions.

To supplement these results we have determined the solubility of silver chloride in water and in solutions of nitric acid and sodium, potassium and ammonium nitrates over a range of temperature from 0 to 50°.

EXPERIMENTAL.

Preparation and purification of materials.—The pure water used in all the experiments was obtained by redistillation of the laboratory distilled water from a dilute, slightly alkaline solution of potassium permanganate. The distillation was carried out in an all-glass apparatus, with a ground-in condenser of fused transparent silica. The water was prepared immediately before use and stored in bottles of resistance glass. It was found to be halogen free when tested in a nephelometer and had a specific conductivity of about 1×10^{-6} mhos.

The silver employed for the preparation of silver chloride was of a high degree of purity (99.996 per cent.) and when dissolved in pure nitric acid the solution was always clear and halogen-free.

The alkali nitrates were purified by repeated evaporation to dryness with concentrated nitric acid to remove traces of chloride which were found to be present in some of them. Three or four digestions with nitric acid always yielded halogen-free material which was then recrystallised and employed for preparing solutions of the required concentrations.

The silver chloride required for the solubility determinations was prepared as follows:—An accurately weighed quantity of silver (1.0788 g.) was placed in a conical flask which was provided with a funnel to collect the spray. Dilute nitric acid (20 c.c. of 1:3) was introduced into the flask which was then gently warmed. After the silver had dissolved, the liquid in the flask was diluted by the addition of 100 c.c. of water through the funnel, and boiled for about 10 minutes. The liquid was then cooled, transferred to a litre measuring flask and made up to the correct volume. A solution of sodium chloride of exactly equivalent concentration to the silver solution, i.e. containing 0.5846 g. of NaCl per litre was next prepared. Equal volumes (200 c.c.) of the solutions of silver and sodium chloride were mixed in a stoppered conical flask and shaken vigorously for about 10 minutes,

Shaking was continued for 5 minutes at intervals of $\frac{1}{2}$ hour for a period of about 10 hours and the flask then set aside. At the end of about 24 hours it was found that the silver chloride had settled completely leaving the supernatant liquid quite clear and it was possible to pour off all but 20 c.c. 500 c.c. of water was then poured into the flask, shaken with the precipitate periodically, allowed to settle and poured off. The silver chloride remaining after four such washings was considered to be sufficiently pure. When the solubility was to be measured in nitric acid or a salt solution, a portion of the solution was used for the last two washings instead of water. The precipitation as well as the washing of the silver chloride was carried out at room temperature in a room illuminated by ruby light and the precipitated silver chloride always remained moist.

In order to determine the solubility, the moist silver chloride (0.1-0.2 g.) and the solvent (100 c.c.) were kept in separate vessels in a thermostat at the required temperature for about $\frac{1}{2}$ hour, after which the silver chloride was added to the solvent and agitated frequently with a glass stirrer during 4 hours. After standing for periods which varied from 24 to 48 hours a portion of the clear supernatant liquid was decanted into another vessel kept in the thermostat and used for determining the solubility. It was found that the solubility did not vary with the period of standing. For temperatures from 25 to 30° the thermostat employed was capable of being regulated to within 0.02°, but for higher temperatures the variation reached 0.5°. For temperatures below that of the room, a large Dewar vessel containing cold water was employed and the temperature regulated by slowly adding iced water. In this case also the variation was 0.5°.

Since the solubility of silver chloride in the solvents employed increases appreciably with temperature it was necessary to dilute all solutions prepared at temperatures higher than that of the room to prevent separation of silver chloride when they were cooled for examination. This was done as follows:—The saturated solution of silver chloride was added to a known volume of water contained in a calibrated 125 c.c. measuring flask with 0.1 c.c. graduations on the neck until the level was near the top of the graduations, and the flask was stoppered, shaken and allowed to attain the temperature of the laboratory. The level of the liquid was then read, giving the quantity of saturated solution added, volume changes on mixing being negligible even with the most concentrated solutions employed. The quantity of water in the flask was sufficient to prevent precipitation on cooling and was calculated from the results of preliminary solubility determinations.

Estimation of the dissolved silver chloride. The nephelometer employed for the determination of silver chloride was of the Kober type with a Klett top-reader. Plain glass cups 10 cm. long provided with opaque glass bottoms contained the solutions during measurements of opalescence.

The standard solutions of silver chloride required for comparisons of opalescence were prepared thus. 2.00 c.c. of the standard silver nitrate solution was accurately measured by means of a micro-pipette, mixed with the same volume of the standard solution of sodium chloride and shaken vigorously. The precipitated silver chloride was just redissolved by gradual addition of a slight excess of very dilute ammonia and the ammonical solution so obtained was diluted to exactly 250 c.c. Each c.c. of this solution contained 0.00318 mg. of silver chloride. In a few cases where higher concentrations of silver chloride had to be estimated, standard solutions with higher concentrations of silver chloride were prepared.

The generation of opalescence for examination in the nephelometer was performed as follows:—An accurately measured quantity (2.00-5.00 c.c.) of the solution to be examined was introduced into the nephelometer cup, diluted with water (2-5 c.c.) acidified with 0.5N. nitric acid (1-2 c.c.) and the silver chloride precipitated by slow addition of 1N. sodium chloride (1-2 c.c.) from a pipette, the liquid in the cup being stirred by a thin glass rod bent into a ring at one end. This procedure is recommended by Scott and Moilliet. (*J. Amer. Chem. Soc.*, 1932, 205). An identical procedure was adopted for generating opalescence in the standard solution of silver chloride. The concentrations of the two solutions were made approximately equal so that the heights of the liquids in the two cups were of the same order when measured. All the precautions necessary for accurate work (Richards and Wells, *loc. cit.*) were strictly observed.

The results obtained in the various experiments carried out in different concentrations of the solvent and at various temperatures are assembled in Tables III and IV. The mixed solvents employed in the experiments detailed in Table IV were prepared by mixing the necessary amounts of the two solvents of known molarities. The table gives the molarity of each of the nitrates and the solubility *S* of the silver chloride.

Each of the values given in the tables is the mean of two or three independent determinations. In each of these the value was deduced from at least four nephelometer readings which were found to be concordant to about 1 part in 400. The variation between different determinations was about 0.5 per cent.

3L

A. d. m. a. n. i. 4. 15

N 32. 16 A. 14

TABLE III.

Solubility of silver chloride, $M./L. \times 10^{-5}$.

Solvent	Molarity	0°	8°	18°	25°	30°	40°	50°
¹ Water	...	0.49	...	1.01	1.39	1.69	2.63	3.82
HNO ₃	0.11	0.60	...	...	1.51	...	...	...
	0.48	0.71	0.89	1.38	1.64	1.84	3.98	7.69
	0.81	0.77	...	...	1.68	...	...	...
	1.00	0.79	...	...	1.74	...	...	...
² NaNO ₃	0.0001	...	...	...	...	1.69	...	...
	0.001	...	...	...	...	1.91	...	...
	0.01	...	...	...	...	2.80	...	...
	0.05	...	...	...	7.35	...	...	...
	0.10	1.35	3.00	4.94	10.1	10.1	10.2	10.8
	0.50	...	...	...	13.5	...	...	...
	0.75	...	...	...	15.5	...	...	...
	1.00	...	...	...	17.1	...	...	...
	3.00	...	...	...	17.5	...	...	...
	Sat. soln.	...	...	...	18.4	...	...	...
KNO ₃	0.10	0.79	0.84	1.42	2.71	2.93	4.30	6.05
	0.50	...	...	...	...	5.15	...	...
	0.75	...	...	...	...	5.92	...	...
	1.00	...	...	...	...	6.93	...	...
NH ₄ NO ₃	0.10	0.68	0.82	1.44	3.27	3.28	4.84	6.75
	0.50	...	...	...	5.23	...	...	...
	0.75	...	...	...	6.33	...	...	...
	1.00	...	...	...	7.40	...	...	...

TABLE IV.

Solubility of silver chloride in mixed solvents at 30°.

KNO ₃ + HNO ₃		S.	NH ₄ NO ₃ + HNO ₃		S.	NaNO ₃ + HNO ₃		S.
0.10	0.25	2.94	0.10	0.25	3.29	0.10	0.25	10.1
0.10	0.50	2.98	0.10	0.50	3.32	0.10	0.50	10.2
0.50	0.25	5.19	0.50	0.25	5.25	0.50	0.25	13.5
0.50	0.50	5.21	0.50	0.50	5.26	0.50	0.50	13.8

The solubility of silver chloride in water in the range studied increases with rise in temperature but it was found that the results could not be expressed by a parabolic equation of the usual type. The expression

$$\log S = 0.01816 t - 0.318$$

calculated by the method of least squares, S being the solubility, fits the experimental results within the limit of error as shown in Table V.

¹ The solubility of silver chloride in water at 10° was found to be 0.72.

² The solubility of silver chloride in 0.1 M. solutions of sodium nitrate at 22 and 23.5° was found to be 5.86 and 6.56 respectively.

Fig. 1.




TABLE V.


Temp.	...	...	0	10	18	25	30	40	50
S calc.	...	...	0.48	0.73	1.02	1.37	1.69	2.56	3.89
S found.	...	...	0.49	0.72	1.01	1.39	1.68	2.63	3.82

The addition of nitric acid or nitrates in small quantities up to a concentration of 0.1 M. results in a marked increase in solubility as may be seen from fig. 1 which shows the results at 25°. On increasing the concentration, the solubility continues to rise but not nearly as rapidly as with more dilute solutions. The effect with nitric acid is the least and with sodium nitrate the greatest. Potassium and ammonium nitrates behave in a similar manner to each other.

The temperature-solubility curves are somewhat remarkable owing to their irregular nature as may be seen from fig. 2. It will be observed that in two cases an abrupt rise is found very near to 25°. As this is the lowest temperature at which an automatically controlled

Fig. 2.

The solubility of silver chloride at various temperatures in water and in 0.1 M. solutions of the nitrates of sodium, ammonium and potassium



thermostat was used it was suspected that the discontinuity might be due in part to change in the method of temperature control, but several additional determinations made just below and just above 25° in the hand-controlled bath confirmed the previous results.

Determinations of solubility by the conductivity method.—Several determinations of the conductivity of silver chloride solutions have been carried out from time to time as will be seen from Table I, but the results obtained do not exhibit satisfactory agreement for the reasons already explained. We have therefore carried out measurements of the conductivity of saturated solutions of silver chloride prepared under the definite conditions described earlier with a view to compare the values obtained by this method with those determined nephelometrically.

The apparatus employed for the conductivity measurements was of the usual type, comprising a Leeds and Northrup bridge. The electrolytic cell was made of pyrex and the electrodes were platinum discs of about 18 mm. in diameter and 3 mm. apart. The determination of the cell constant and the calculation of the molar concentration of silver chloride in the solvent employed were all performed in the usual manner, assuming that silver chloride was completely dissociated in solution, and adopting Johnston's values (*J. Amer. Chem. Soc.*, 1909, **31**, 1015) for the equivalent conductances of ions.

The mean value at 25° obtained from three separate experiments involving about 20 determinations was 1.373×10^{-5} M./L. which is in fair agreement with the value 1.388×10^{-5} M./L. obtained nephelometrically.

COMPLEX FORMATION

The large increase in solubility produced by the addition of salts points to complex formation, the extent of which can be calculated by means of the equation given by Pinkus and Berkolaiko (*loc. cit.*) who have calculated the degree of complexity of solutions of silver chloride in potassium chloride.

According to these authors if α_{Ag} , V_{Ag} and $[Ag^+]$ are the activity, the coefficient of activity and the concentration respectively of Ag^+ ion, S the solubility and x the degree of complexity.

$$S = [Ag^+] + Sx \text{ or } x = 1 - \frac{\alpha_{Ag}}{V_{Ag} \times S} \text{ since } [Ag] = \frac{\alpha_{Ag}}{V_{Ag}}$$

Since S is known and the values of α_{Ag} and V_{Ag} may be obtained

from tables of activity coefficients (Lewis and Randall, *Thermodynamics*, p. 382) x may be calculated for the various solutions examined. The results so obtained are given in Table VI.

TABLE VI.

Complexity of silver chloride solutions.

Solvent	Conc.	Temp.	$S \times 10^5$	γ_{Ag}	$\alpha_{Ag} \times 10^6$	x
NaNO ₃	0.1	0	1.35	0.77	2.27	0.78
	"	18	4.94	"	2.65	0.93
	0.05	25	7.35	0.80	3.27	0.94
	0.1	"	10.07	0.77	2.48	0.97
	0.001	30	1.91	0.97	14.25	0.23
	0.01	"	2.81	0.90	10.47	0.59
	0.1	"	10.09	0.77	3.41	0.96
	0.1	40	10.23	"	8.79	0.89
	"	50	10.78	"	17.60	0.79
NH ₄ NO ₃	0.1	0	0.69	0.77	4.43	0.17
	"	18	1.44	"	9.11	0.18
	"	25	3.27	"	7.65	0.70
	"	30	3.29	"	10.44	0.59
	"	40	4.84	"	18.6	0.50
	"	50	6.75	"	28.12	0.46
KNO ₃	0.1	0	0.79	0.77	3.88	0.36
	"	18	1.42	"	9.22	0.16
	"	25	2.71	"	9.21	0.56
	"	30	2.93	"	11.71	0.48
	"	40	4.30	"	20.91	0.37
	"	50	6.18	"	30.68	0.36

These figures indicate that (1) the degree of complexity is a maximum at 25°, (2) it is much greater for solutions of sodium nitrate than for ammonium and potassium nitrates and (3) in the case of sodium nitrate it increases with the concentration of the solution at constant temperature.

SUMMARY

1. The solubility of silver chloride in water and in aqueous solutions of nitric acid and the nitrates of sodium, potassium and ammonium has been determined by a nephelometric method over a temperature range 0-50° under conditions approximating to those prevailing during analytical work.


2. The solubility in water at 25° has been deduced from conductivity measurements giving the value $1.373 \times 10^{-5} M/L$ as compared with $1.388 \times 10^{-5} M/L$ by the nephelometric method.

3. The solubility of silver chloride in solutions of sodium nitrate has been found to be much higher than in the other solutions studied and this is due to the existence as complex ions of a large proportion of the silver ions.

We desire to express our gratitude to Professor H. E. Watson for his constant interest and valuable suggestions during the progress of this work.

Department of General Chemistry,
Indian Institute of Science,
Bangalore.

Accepted 20-7-33.]



JL

A, d, m, n, v, i, u, s

N 33 16 A. 14

PRINTED AND PUBLISHED
BY V. M. PHILIP AT THE DIOCESAN PRESS,
POST BOX 455, MADRAS—1934. C9037.

VOLUME 15 (A)—1932.

- South India Hides and Skins. By V. N. Patwardhan and M. Subra-
a Sastry. Price, As. 12.
- Lat Examination of the Root-Bark of *Plumbago Rosea*, Lion. By M. C.
Tummin Katti and V. N. Patwardhan. Price, As. 8.
- Dilatometric Studies in Enzyme Action. Part II. Contraction Constants of Enzyme-
Substrate Reactions. By M. Sreenivasaya and H. B. Sreerangachar. Price,
As. 12.
- Part IV.—Aryl Di- and Poly-Stibinic Acids, Distibinous Oxides and Distibino-Compounds. By
Sohrab M. Mistry and Praphulla Chandra Guha. Price, Re. 1.
- Part V.—Bhadravati Wood-Tar and its Utilization. By Y. K. Raghunatha Rao, B. Sanjiva Rao
and H. E. Watson. Price, Re. 1.
- Part VI.—The Equilibrium between Dimethyl Ether, Methyl Alcohol and Water. By N. G.
Gajendragad, S. K. Kulkarni Jalkar and H. E. Watson. Price, Re. 1.
- Part VII.—Studies in Indian Essential Oils. I. Essential Oil from Leaves of *Litsie Zeylanica*
Linn. By B. Sanjiva Rao. II. Essential Oil from Flower-Heads and Stalks
of *Andropogon Kuntzeanus*, Hack, *Var. Foveolata*, Hack. By B. Sanjiva Rao.
III. Essential Oil from Leaves of *Thymus Serpyllum*, Linn. By Jagjit Singh
and B. Sanjiva Rao. IV. Essential Oil from Leaves of *Cinnamomum Zeylani-*
cum, Breyn. By (the late) V. P. Shintre and B. Sanjiva Rao. Price, Re. 1.
- Part VIII.—Organic Manure from Sewage, Town Refuse and Waste Vegetation. By J. Jagan-
natha Rao and V. Subrahmanyam. Price, Rs. 1-8.
- Part IX.—I. Studies in Enzyme Action. Part V. Ageing of Amylases in Aqueous Solution.
By K. Venkata Giri and V. Subrahmanyam. II. Studies in Enzyme Action.
Part VI. Heat Inactivation of Pancreatic Amylase. By K. Venkata Giri. III.
Studies in the Proteins of Indian Foodstuffs. Part IV. The Proteins of
Fenugreek (*Trigonella Foenum Graecum*). By Y. V. Srinivasa Rau and M.
Sreenivasaya. Price Rs. 1-8.
- Part X.—Studies in Bridge Formation. Part I. Attempts to Synthesise Bicyclic Terpene Deri-
vatives by a New Method. By P. P. Patel and P. C. Guha. Part II. Formation
and Stability of a Bridged cycloHexenone, 1-Dimethylethane—III
cyclobutane-2-one. By P. S. Mayuranathan and P. C. Guha. Price, Re. 1.

VOLUME 15 (B)—1932.

- Part I.—A New Permeameter. By K. V. Karantha. Price, Rs. 1-8.
- Part II.—Measurement of Power Factor and Loss in Dielectrics. By T. J. Mirchandani,
G. Yoganandam, S. K. Roy and N. V. Narayanawami. Price, Rs. 1-4.
- Part III.—Transients in Negative Constant Series Circuits. By L. C. Verman. Price, Rs. 1-4.
- Part IV.—Telephony by Carrier and One Side Band. By S. P. Chakravarti. Price, Re. 1.

VOLUME 16 (A)—1933.

- Part I.—Rhaponticin, and Anthraquinone Derivatives from *Rheum Emodi*, Wall. (Indian, or
Himalayan Rhubarb). By M. Ghouse Mohiuddin and M. C. Tummin Katti.
Price, As. 12.
- Part II.—Constitution of the So-Called Dithiourazole of Martin Freund. Part V. Isomerism
of Hydrazodithiodicarbonamides, Iminothiolthiobiazoles and Di-R-Iminothio-
biazoles. By Shaha L. Janniah and P. C. Guha. Part VI. Isomerism of
Hydrazomonothiodicarbonamides, Iminothiothiazolones and Monothiothiazoles. By
S. L. Janniah and P. C. Guha. Price, Re. 1-4.

- Part III.—Indian Medicinal Plants. Part I. *Withania Somnifera*.
By D. N. Majumdar and P. C. Guha. Price, As. 12.
- Part IV.—I. The Colouring Constituents of Alkanet Root (*Anchusa Tinctoria*).
The Constitution of Alkannin. By Mangesh V. Betrabet and
Chakravarti. II. Note on the Constituents of the Wax from A
By M. V. Betrabet and G. C. Chakravarti. Price, Re. 1.
- Part V.—The Velocity of Alcoholysis of Esters of Dibasic Acids. By Z. H. Patel
H. E. Watson. Price, As. 12.
- Part VI.—I. Dilatometric Studies in Enzyme Action. Part III. Contraction Constants of
Enzyme-Substrate Reactions. By H. B. Sreerangachar and M. Sreenivasaya.
II. Physiological Products of the Lac Insect. Part I. A Preliminary Investiga-
tion. By N. K. Ranga Rao and M. Sreenivasaya. Price, Re. 1.
- Part VII.—I. Studies in the Proteins of Indian Foodstuffs, Part V. The Alcohol Soluble
Protein of Fenugreek (*Trigonella Foenum Graecum*). By Y. V. Sreenivasa Rao,
B. N. Sastri and N. Narayana. II. The Mucilage of Fenugreek (*Trigonella*
Foenum Graecum). By C. R. Harihara Iyer and B. N. Sastri. Price, As. 8.
- Part VIII.—I. Contributions to the study of the Spike Disease of Sandal (*Santalum Album*,
Linn.), Part XIII. Investigation of the Hexone Bases. By Y. V. Sreenivasa
Rao. II. Contributions to the Study of the Spike-Disease of Sandal (*Santalum*
Album, Linn.), Part XIV. Study of Mosaics Associated with Spiked Areas.
By Y. V. Sreenivasa Rao. Price, As. 8.
- Part IX.—Physiological Products of the Lac Insect. Part II. Investigation of the Water
Soluble Nitrogenous Constituents. By N. K. Ranga Rao. Price, As. 8.
- Part X.—Extension of Michael's Reaction. Part III. By Tejendra Nath Ghosh and
Praphulla Chandra Guha, Price, As. 12.
- Part XI.—Studies in Bridge Formation. Part III. The Influence of the Methyl and Carbethoxy-
groups in the Internal Dehydration of *gem*-Disubstituted Dihydroresorcinol
Esters. Part IV. Tautomerism in isophorone. By P. S. Mayuranathan.
Price, Re. 1.
- Part XII.—Synthesis of Proteins in Plants. Part I. Conversion of Nitrates into Protein in
Helianthus Annus, Linn. By K. S. Varadachar. Price, As. 12.
- Part XIII.—Contributions to the Study of Spike-Disease of Sandal (*Santalum Album*, Linn.)
Part XV. The role of plant acids in health and disease. By A. V. Varadaraja
Iyengar. Price, Re. 1.
- Part XIV.—The Solubility of Silver Chloride. By P. C. Dave and K. R. Krishnaswami.
Price, Re. 1.

VOLUME 16 (B)—1933.

- Part I.—Taylor's Frequency Tripler. By P. Rajagopal Pillai and F. N. Mowdawalla. Price,
Re. 1.