

1590

Journal of.
South Indian medicine.
vol-I NO-6

15.6.1935



SL

-m21, N35 S2

N35.1. 6.

147296

INDEX NUMBER

REGD. No. M. 3201

**JOURNAL OF
SOUTH INDIAN MEDICINE**

JUNE 15, 1935

ISSUED BY THE SOUTH INDIAN MEDICAL UNION, MADRAS

VOLUME 1

NUMBER 6



KESARI KUTEERAM
EGMORE :: :: MADRAS.

JOURNAL OF SOUTH INDIAN MEDICINE



MÖLLER'S A. B. C. D. VITAMIN FOOD

EACH FLUID OUNCE REPRESENTS:

VITAMIN

- A 11 320 U. S. P. Units
- B 5 66 Rat Units (Complex)
- C 28 3 International Guinea pig Units
- D 3 396 U. S. P. Units

Dose: CHILDREN— $\frac{1}{2}$ teaspoonful 3 times a day.

ADULTS—1 to 2 teaspoonfuls 3 times a day.

Sole Agents for India, Burma & Ceylon

The South Indian Mercantile Co.,

P. O. NO. 505. 12, MASSARU STREET. P. T. MADRAS.



JOURNAL OF SOUTH INDIAN MEDICINE

VOLUME ONE

CONTENTS

NUMBER 1	JANUARY, 1935
Foreword, Major-General Sir F. P. Connor, Kt., D.S.O., K.H.S., I.M.S., ...	1
Surgery of the Breast, Lt. Col. V. Mahadevan, I. M. S. ...	3
Some Common Drugs of Cardiac Practice, T. Krishna Menon, M. B. ...	17
Some Recent Advances in Medicine, R. Viswanathan, M. B., M. R. C. P. ...	23
<i>Retrospect</i> ...	35
<i>The Late Dr. T. Krishna Menon</i> ...	37
Dr. T. Krishna Menon ...	39
NUMBER 2	FEBRUARY, 1935
Bacillus Coli Infections, T. Satakopan, M. D. ...	41
Streptococcal Disease, G. Raman Pillai, M. B. Ch. B. (Edin) ...	58
Radium Therapy, P. Rama Rau, D. M. R., (Vienna) ...	68
GLEANINGS FROM MEDICAL LITERATURE	
(1) Status Lymphaticus ...	81
(2) Carcinoma of the Male Breast ...	82
(3) A Case of Evipan Paralysis ...	83
<i>Rural Medical Relief</i> ...	85
<i>The Building Craze</i> ...	86
News and Notes ...	88
NUMBER 3	MARCH, 1935
A Note on a Case of Tropical Typhus occurring in Madras, Capt. R. D. Alexander, I. M. S. ...	89

147296

A Note on Tropical Typhus Fever, Major G. R. McRobert, I. M. S.	...	...	93
Gastric Neuroses, T. Satakopan, M. D.	...	...	97
Headache: Its Differentiation and Significance, T. S. Duraiswami.	...	...	104
GLEANINGS FROM MEDICAL LITERATURE—			
(4) Spinal Anaesthesia : Nervous System Sequelae	...	...	118
(5) A Lowered Mortality in Acute Appendicitis	...	...	119
(6) Chronic Appendicitis and the Right Lower Quadrant Syndrome	...	...	120
(7) Chronic Amoebiasis	...	...	121
(8) Alcohol and Automobile Accidents	...	...	123
(9) Studies on Reticulocytes	...	...	123
(10) Treatment of Fracture of Spinal Column	...	...	124
Research and Record	...	...	125
Problems on Filariasis	...	...	127
NUMBER 4			
APRIL 1935			
The Diagnosis of Kala-Azar, K. C. Paul, M. D., and V. Govindarajulu, M. B. B. S.,	...	...	129
The Leukaemias, S. Thambiah, M. C., M. R. C. P., D. T. M. & H., F. D. S.,	...	...	145
The Problem of Birth Control, T. S. Balasubramania Iyer, M. B. B. S.	...	...	152
GLEANINGS FROM MEDICAL LITERATURE—			
(11) Fibrocystic Disease of Bone	...	...	160
(12) Pathology and Treatment of Lymphadenoma	...	...	161
(13) The Management of Urologic Complications in Injuries to the Spine	...	...	161
(14) Epidural and Perineural Injections in the Treatment of Peripheral Pain	...	...	162
(15) Oral Urography	...	...	163
(16) Recurrent Dislocation of the Shoulder	...	...	163
REVIEWS	...	...	165
Budget Speeches	...	...	167
MEDICAL PERIODICALS	...	...	169
NUMBER 5			
MAY, 1935			
Ligature of the Femoral Vein in Diabetic and Senile Gangrene, P. Govinda Rau, M.B.B.S.	...	...	179

The Need for a National Anti-Cancer Campaign in India, P. Rama Rau, D. M. R. (Vienna)	...	...	183
The All-India Ophthalmological Society, Fourth Conference, Welcome Address—Dr. E. V. Srinivasan, M.B.,C.M.	...	...	195
Opening Speech—Sir Mahomed Usman, B.A., K.C.I.E.	...	...	199
Presidential Address—Lt. Col. R. E. Wright, I.M.S.	...	...	202
Rural Medical Practitioners' Fifth Provincial Conference, Presidential Address—D. S. Ramachandra Rao, M.A., M.D.,	...	...	209
GLEANINGS FROM MEDICAL LITERATURE—			
(17) Strychnine and the Barbiturates: Mutual Antidotes	...	...	213
(18) Artificial Pneumothorax in the Treatment of Pneumonia	...	...	213
(19) Treatment of Varicosities	...	...	215
(20) Diagnosis and Treatment of Intestinal Obstruction	...	...	215
Election to the Indian Medical Council	...	...	217
Control of Cancer	...	...	218
The Rural Medical Scheme	...	...	219
MEDICAL PERIODICALS	...	...	221
NUMBER 6			
JUNE, 1935			
A Search for New Types of Tropical Splenomegaly, T. Bhaskara Menon, M.D., M.R.C.P.	...	...	227
Note on a Case of Popliteal Aneurysm Treated by Hunter's Method, B. M. Sundaravadanan, B.A., M.B.B.S., F.R.C.S. (Eng.)	...	...	233
Development of the Heart, Normal and Abnormal, P. K. Koshy, B.A., M.B., F.R.C.P. (E.)	...	...	236
GLEANINGS FROM MEDICAL LITERATURE—			
(21) Epilepsy and its Treatment	...	...	248
(22) Nervous Breakdown	...	...	249
(23) Acute Infections of the Upper Respiratory Tract, Middle-Ear and Mastoid	...	...	251
(24) Radiation in the Treatment of Cancer	...	...	253
(25) Endocrine Effects of Certain Ovarian Tumours	...	...	254
(26) The Treatment of Angina Pectoris	...	...	255
Clinical Memoranda	...	...	257
The Treatment of Small-pox, V. Govindan Nair, M. R. C. P. (E.)	...	...	257
REVIEWS	...	...	258
The Problem of Tuberculosis	...	...	259
The Vizagapatam Medical College	...	...	260
Colonel E. W. C. Bradfield, O. B. E., C. I. E., I. M. S.	...	...	261
Retrenchment in Hospital Expenditure.	...	...	261
MEDICAL PERIODICALS	...	...	263

JOURNAL OF SOUTH INDIAN MEDICINE
(INCORPORATING THE BULLETIN OF THE SOUTH INDIAN MEDICAL UNION)

VOLUME I

JUNE, 1935

NUMBER 6

**A SEARCH FOR NEW TYPES OF
TROPICAL SPLENOMEGALY ***

T. BHASKARA MENON, M.D. (Madras), M.R.C.P. (Lond.)
Pathologist, Government Royapuram Hospital, Madras.

Nearly seven years ago, when I was working in the pathology department of the Madras Medical College, Dr. Krishna Menon showed me a series of case histories of patients with chronic enlargement of the spleen, where the enlargement could not be classified on clinical grounds. Complete clinical investigation had failed to reveal the cause of the enlargement. The question was whether these cases belonged to a new type. Was there an unknown type of tropical splenomegaly? In the absence of any definite post-mortem evidence I could not answer the question.

In 1932, De published his analysis of splenic enlargement in Bengal and postulated the existence of a new type, with definite clinical associations and a definite histological picture. In the same year, Hughes and Shrivastava in Lahore brought forward some evidence for the existence of a type of post-malarial haemolytic anaemia in a group of cases. In 1933, Sir Frank Connor's paper at the British Medical Association dealt with a survey of splenic enlargement in Bengal, Bombay, and Rangoon, covering nearly 1,200 cases. He raised the question of surgical intervention in cases of post-malarial and possibly post-kala-azar types where the infections had died down, but left enlarged spleens.

I attempted an analysis of the cases from old autopsy records of the Madras Medical College. These were kindly lent to me by Dr. Goyle to

* Abstract of a paper read before the South Indian Branch of the British Medical Association, 1934.

whom my thanks are due. I have included cases from the more recent autopsy records from the Royapuram Medical School. The records from the Medical College I have analysed date from 1889 to 1929, a period of 40 years. The figures relate to 5,034 autopsies out of which 1,101 showed splenic enlargement as judged by our standards. This gives a percentage of 21.9 as compared with 50 per cent in De's figures for Bengal and 40 per cent in Gharpure's figures for Bombay.

This analysis and a discussion on that form the subject of a paper in the *Indian Journal of Medical Research* (Bhaskara Menon, 1934). I might be pardoned if I quote extensively from this paper and if I lay more stress on the pathological rather than on the clinical aspects of the question.

In the first group of my classification I have included the types of acute enlargement—the septic spleen of the pathologist. I shall not comment on this group of 399 cases except to point out the high figure for amoebic abscess of the liver and amoebic dysentery—a factor which is of interest to workers in the tropics.

In the second group are the types of chronic venous congestion including one case of portal thrombosis and another of thrombosis of the splenic vein.

In the third group are included all cases of cirrhosis of the liver showing splenic enlargement. In portal cirrhosis the splenic enlargement is slight in a large proportion of cases, but others show marked enlargement. These are cases that merge almost imperceptibly into Banti's disease. There are two instances of this condition. That splenic enlargement in cirrhosis is not a matter of mere chronic venous congestion is now certain. The large spleens met with in some cases have been the subject of much discussion. The old idea of a splenomegalic cirrhosis presupposes toxins formed in the spleen travelling to the liver and causing damage. McNee regards this type as due to toxins which are dammed back into the spleen, owing to obstruction to the portal circulation. It seems to me that toxins might have varied reactions on the parenchyma and reticulo-endothelium. Those that affect the parenchymatous cells may cause cirrhosis, and those which act more on the reticulum may induce splenic enlargement. The cirrhosis is slight and occurs only late in the disease.

Malarial cirrhosis is a controversial theme. On the one side Osler from Johns Hopkins, Rogers from Calcutta, Tirumurthi and Radhakrishna

Rao from Vizagapatam all deny its existence. Rolleston and McNee doubt whether it is a coincidence. I would not dare to venture out against this formidable array, but Sinton and Hughes have demonstrated the effect of malaria in causing damage to the liver, and Hughes from Lahore cites cases where the gradual change could be followed from malaria with splenic enlargement to cirrhosis with ascites. I shall only remark there is nothing incongruous in the idea of a malarial cirrhosis. Maximow has taught us that the reticulum cell is capable of being transformed into hyperplastic endothelium and even into collagen fibre if the stimulus is chronic. With kala-azar cirrhosis we are also on uncertain ground. Rogers has described an intercellular cirrhosis and in addition a portal type. Shanks and De have found fibrotic changes in the liver by special staining in 30 per cent of their cases, though they conclude that this is not a regular event in kala-azar.

The next group of splenic enlargement, in disease of the haemopoietic system including mostly leukaemias, is of no special significance. In the five cases of ancylostomiasis I have recorded, the enlargement may be due to toxic absorption from the intestine.

Chronic hyperplasia from the infective granulomata form a well recognised group. Malaria and kala-azar are of special importance here. Malarial enlargement is responsible for 11.3 per cent of the total and kala-azar has even a higher figure of 14.3 per cent. It is noteworthy that De reports only 6.5 per cent of malarial spleens in his series. Tirumurthi and Radhakrishna Rao from Vizagapatam point out the low incidence of malarial spleen in the autopsy records as compared with clinical findings. Hughes and Shrivastava report a high incidence in Lahore.

When gametocytes are absent in the blood, the clinical diagnosis of chronic malaria rests mostly on the history. Some would lay a good deal of stress on the finding of pigment-containing mononuclear cells. Our experience has been the reverse. Such pigmented monocytes are infrequently met with even in cases with parasistes in the blood. Spleen puncture is no doubt a risky procedure, but I believe it is much more certain in its results. To the pathologist, the appearance of the spleen is sufficiently indicative. Histological sections of the spleen and liver are of great confirmatory value, since haemozoin can be demonstrated. In doubtful cases, the ammonium sulphide reaction may be tried, as haemozoin is dissolved by ammonium sulphide.

Uncommon types form the next group in this analysis. Chronic hyperplastic peritonitis of obscure origin (the endemic ascites of Megaw, the superior peritonitis of Sprawson) is a type that is of interest. Enlargement with primary carcinoma of the liver is a feature to which I believe little attention has been drawn so far. Enlargement due to follicle hyperplasia is a curious condition first described in 1926 in America. These are instances of reticulo-endotheliosis, the so-called "reticulosis" of modern writers. I have been frequently asked by clinicians if there is any evidence of the Gaucher type here. So far I have met with none. There is one case, however, in my series of localised lipoid deposits in diabetes.

I have grouped together 45 cases, where the cause of the enlargement is obscure. The existence of this group, I believe, is mostly due to imperfect investigation rather than to the presence of any well-defined unknown types. I have attempted to sub-divide this group on a morphological basis. Of these, the first 14 were probably unrecognised kala-azar. These cases were recorded before the discovery of the parasite. Another 5 form a series in which the appearance suggests kala-azar, but no parasites were found. To my mind, these are cases where the parasites have undergone post-mortem disintegration till they are no longer recognisable. One other case seems to be an unrecognised leukaemia. Three others showed a late cirrhosis of the liver. One other was probably of the same kind before the occurrence of the cirrhosis. Two more cases suggest the co-existence of malaria and some acute infection. Another was due to perisplenic suppuration. In the remaining 18 cases which could not be classified, I could find no common factor since the cases have been imperfectly investigated. Presumably, the enlargement is due to a combination of factors. I have cited an instance in my first paper of a case where there was a large spleen with no history of malaria, yet malaria pigment was found in the spleen puncture. A strongly positive Wassermann and Kahn reaction suggested the co-existence of syphilis.

When we consider the possibilities of new types in South India, we have to take into account two types, one described by De in Bengal and the other by Hughes and Shrivastava in Lahore. If we take up De's type of Bengal splenomegaly, according to De, these cases come from West Bengal and the lower parts of Eastern Bengal. They all give a history of irregular fever which is often diagnosed as malaria, kala-azar, or enteric fever, but which is not amenable to quinine or antimony. There is a gradual and

progressive splenic enlargement which is followed by secondary anaemia. The liver is enlarged in the early stages, but cirrhosis gradually supervenes, followed by ascites. Histologically, coarse fibrosis of the trabeculae, reticular fibrosis and red cell phagocytosis are the prominent features.

A study of this picture reminds me of another which was painted and repainted by Banti and commented on and criticised by a whole continent. I refer to the syndrome of Banti, so-called Banti's disease, where the condition starts at first as a splenic anaemia with occasional irregular fever and ends up in cirrhosis of the liver. That the histological picture is not constant is borne out by Banti's subsequent descriptions and by a whole host of critics, Durr, Nagaeli, and Krumbhaar, to cite a few. The reticular fibrosis that De emphasises seems to me nothing more than the *fibro-adenie* of Banti. Red cell phagocytosis and deposit of blood pigment, though denied by Banti, has been noted by a number of observers. When we remember that the spleen is the charnel house of the erythrocyte, red cell phagocytosis in the spleen is neither distinctive nor significant.

There are those who might say that Banti's disease has no meaning and it is only a syndrome. Well, then, call it what we like, an uncommon type of portal cirrhosis, as McNee would have it. Yet we cannot deny that there is such a clinical picture.

If we pass on to the type described by Hughes and Shrivastava, are we dealing with a new type? They believe that repeated attacks of malaria with chronic enlargement of the spleen is followed by a hyperactivity of the reticulo-endothelium so that the phagocytic endothelial cells would pick up normal erythrocytes and induce a haemolytic anaemia. They cite cases where the bilirubin content of the serum is high. Fragility changes in the red cells are exceptional in their series. The malarial infection is believed to have died down since the parasites were absent in the blood and there was no fever. It is possible, however, that in such cases the malarial infection might have been latent and still smouldering in the spleen.

Ross's studies on malaria have taught us that 25,000 parasites per c. mm. are necessary to affect the temperature curve and induce clinical malaria. In latent malaria, parasites would be found in such scanty numbers that demonstration in a drop of blood on a slide would be difficult, if not impossible. In such cases, are we justified in concluding that the malarial infection has died down and that the haemolytic anaemia is due

to the hyperactivity of the reticulo-endothelium on the normal erythrocyte? It is only by a coordinated investigation between the clinician, the bio-chemist and the pathologist that this question can be settled.

To revert to the original theme, are there new types of tropical splenomagaly in South India? Are there cases masquerading as chronic malaria, kala-azar and splenic anaemia? The history of kala-azar itself suggests the possibility of such new types. But from the analysis that I have attempted, based mostly on dead pages, I am not convinced that any well-defined unknown type exists here in South India.

REFERENCES

- KRISHNA MENON, T. (1927), Personal Communication.
 DE, M. N. (1932), *Ind. Journ. Med. Res.*, XIX, 1029.
 HUGHES, T. A. and SHRIVASTAVA, D. L. (1931),
Ind. Journ. Med. Res., XIX, 565.
 CONNOR, Sir F. P. (1933), *Ind. Med. Gaz.*, LXVIII, 35.
 GHARPURE, (cit. Connor).
 BHASKARA MENON, T. (1934), *Ind. Journ. Med. Res.*, XXI, 695.
 McNEE, J. W. (1932), *Brit. Med. Journ.*, i, 1007.
 OSLER, W. (Quoted by Rolleston and McNee 1929).
 ROGERS, L. (1932), *Lancet*, ii, 124.
 TIRUMURTHI, T. S. and RADHAKRISHNA RAO, M. V. (1933),
Journ. Ind. Med. Assoc., II, 269.
 ROLLESTON, H. and McNEE, J. W. (1929), *Disease of the Liver, Gall-Bladder, and Bile-Ducts*, London, 332.
 SINTON, J. A. and HUGHES, T. A. (1933),
Ind. Journ. Med. Res., XII, 409.
 MAXIMOW, A. A. (1933), In Cowdry's *Special Cyology*, II, 618.
 ROGERS, L. (1930), In Rogers and Megaw: *Tropical Medicine*, 507, 508.
 SHANKS, B. and DE, M. N. (1934), *Ind. Journ. Med. Res.*, XIX, 457.

NOTE ON A CASE OF POPLITEAL ANEURYSM TREATED BY HUNTER'S METHOD.

B. M. SUNDARAVADANAN, B.A., M.B.B.S., L.R.C.P. (Lond.), F.R.C.S. (Eng.)

Honorary Surgeon, Government Royapettah Hospital, Madras.

Popliteal Aneurysm is by far the commonest type of aneurysm occurring in the extremities, and also the most amenable to surgical treatment.

CASE.—Mr. S.....Hindu, aged 28, was seen by me about the middle of February 1935, his only complaint being that he had a little pain in his right knee of about three months' duration. There was no history of having had any sore on his penis, but he admitted having exposed himself to the liability of venereal infection.

On examination, the patient was a well-built man. The right lower extremity as a whole was better developed than the left one, and this, he said, was the case from his boyhood, and he had always felt the left one weaker than the right. On the back of the right knee-joint, a characteristic *expansile swelling* of the size of a large hen's egg could be palpated. There was a definite *thrill* palpable over the swelling which occupied practically the whole of the popliteal fossa. The pulsation was arrested by pressure on the femoral artery above, and a definite *systolic bruit* was heard over the tumour. There was no congestion or oedema of the foot and leg. There was no limitation of movement of the knee-joint, and, apart from the slight pain in the joint, the patient was not troubled about the swelling even in the least. The heart and the other systems were normal. The blood pressure was not raised and the Wassermann reaction was strongly positive.

The patient was put on potassium iodide, 15 gr. per day, at first, and this was increased to 60 gr. per day in a week's time. The pain was immediately relieved.

Surgical Treatment.—Hunter's method of treatment consists in ligaturing the artery at some distance from the aneurysmal sac, a branch intervening between the aneurysm and the site of ligature. The object

of ligaturing the artery at a distance is not to cut off the entire blood flow into the aneurysmal sac, but to slow down the blood-stream to such an extent that clotting is promoted inside the sac. For cases of aneurysms which do not allow of ligature of the main artery, just proximal to the sac, Hunter's method is best suited. The great objection to Hunter's method of proximal ligature at a distance from the sac is that two sets of collateral circulation have to be developed, one around the sac and the other around the site of ligature. If the collateral circulation is not adequate, gangrene is very likely to ensue. In about 25 per cent of cases of aneurysms of the popliteal artery treated by Hunter's method, gangrene has occurred.

In the above case Hunter's method was decided on, as the patient was quite a young man with good physique and the likelihood of efficient collateral circulation establishing itself was very good. In doubtful cases, intermittent compression of the artery above helps to establish a collateral circulation.

Under spinal anaesthesia, using novocaine, the femoral artery was ligatured in Hunter's canal, and the limb, after the patient was returned to bed, was covered with cotton wool and put in a box-splint.

The patient suffered from severe pain, and the limb, which was pale and cold at first, soon became warm, and, in about twenty-four hours, the toes regained their normal colour and warmth. The pain was controlled by morphine and the sutures were out on the seventh day. It took nearly a fortnight after the operation for the patient to walk about. There was no pulsation immediately after the operation, and the pulsation never returned any time afterwards. There was a firm hard swelling in place of the aneurysmal sac—evidently the sac had become completely filled with blood-clot and this slowly decreased in size. Three months after the operation, the hard swelling disappeared completely.

The patient was put on a complete course of antisyphilitic treatment.

Other Methods of Treatment.—Proximal ligature of the artery close to the sac is possible only when the aneurysm is situated low down in the popliteal fossa.

Complete excision of the sac is not possible in late cases as there are considerable adhesions between the sac and the surrounding structures,

but this method of treatment should be attempted wherever it is possible. After the sac is excised, it may be possible to bring the two ends of the artery together; otherwise, a piece of a vein is grafted between the two ends of the artery and the continuity is established.

Matas's operations of *Endo-Aneurysmorrhaphies* have not come into general use, although in his own hands and in those of a few American surgeons, they have given good results. Most of Matas's operations have been performed on popliteal aneurysms. The cases we see here come very late, and the sac has grown into such a big size and has caused all the neighbouring nerves to be firmly adherent to it, that reconstructive and restorative methods of aneurysmorrhaphy are out of question. The obliterative method may be tried in selected cases, but the risks of gangrene setting in are great.


Hunter's operation is contraindicated where serious cardiac disease exists, where pressure over the vessel does not control the circulation through the sac, where the peripheral vessels are extensively calcified, or where gangrene of the limb is threatening or is present.

DEVELOPMENT OF THE HEART, NORMAL AND ABNORMAL.

P. K. KOSHY, B.A., M.B.B.S., F.R.C.P. (E.)
Professor of Anatomy, Madras Medical College.


It will be advantageous to bear in mind from the outset a general truth that malformations of the body including those of the heart are almost invariably due to an arrest of natural processes brought about by obscure forces.

The heart is of mesodermal origin, being a modified blood vessel formed by the fusion of two bilateral primitive arteries called dorsal aortae. It appears in the third week of intra-uterine life, in the cervical region, and is then below the mandible and ventral to the pharynx. An embryonic partition called *septum transversum* bounds it caudally. A cross-section of it at this stage has the appearance shown in Diagram 1.



It will be seen that the organ is composed of two parts, an inner one representing the endothelial lining of the adult heart, and an outer, the rest of the organ, viz., the musculature and the visceral layer of pericardium. The inner comes from the vasoformative mesenchymal cells


which are present throughout the embryonic body and the outer from the splanchnic layer of mesoderm. The bilateral origin of the heart, its position within the pericardium, and its connection with the pharynx are also seen. By its cranial end it becomes continuous with the dorsal aortae which connect it with the placenta through the umbilical arteries and by the opposite extremity it receives the umbilical and vitelline veins (see Diagram 2).



The heart begins to beat late in the 3rd week of intra-uterine life. At this stage it is pre-eminently a pump for the placenta, sending blood to it by way of the umbilical arteries and receiving it back through the umbilical veins. Up to the time of birth this continues to be the most essential function of the organ.

The next stage in the organisation of the heart is highly characteristic. The organ now consists of a twisted loop divisible into a number of parts each of which has some share in the structure of the adult organ. The parts as enumerated from the caudal to the cranial extremity are: 1. sinus venosus; 2. atrium or auricle; 3. atrial or atrio-ventricular canal; 4. ventricle; 5. bulbus cordis; 6. truncus arteriosus (see Diagram 3). It

DIAGRAM 3



Heart in Fifth Week (only the endothelial lining of the Cardiac tube is shown).


will be seen that the sinus venosus and the truncus arteriosus are bound to the pericardium by folds of mesoderm, called respectively *venous* and *arterial mesocardium*, and that the points of their attachment are separated at this stage by a considerable interval occupied by what is called the *dorsal mesocardium*. As development proceeds, this interval will ultimately be reduced to a potential space, the *transverse sinus* of the adult heart—the venous mesocardium in relation to the superior vena cava and the right atrium having met the arterial mesocardium enclosing the pulmonary artery and the aorta. (These structures will have taken the place of the sinus venosus and the truncus arteriosus by that time). Three pairs of venous channels now empty into the sinus venosus: two vitelline veins, two umbilical veins, two ducts of Cuvier. The last pair, i.e., the ducts of Cuvier, carries venous blood from the region of the trunk. These vessels have to traverse the septum transversum before they enter

the sinus venosus, and it may be noted that this part of the embryonic partition will become a portion of the pericardium, viz., the one which is attached to the central tendinous division of the diaphragm. From the truncus arteriosus at the cranial end of the embryonic heart the aortic arches come out to connect it with the dorsal aortae.

We shall now proceed to trace the growth changes of each of the six subdivisions of the primitive cardiac tube.

1. *The sinus venosus*.—It remains as a separate structure in lower vertebrates but in mammals is absorbed into the atria (auricles). Its entrance into the atrial portion of the heart tube is guarded by two valves which prevent regurgitation of blood into the sinus. The valves meet above and below, forming the *superior* and the *inferior fornix* respectively (see Diagram 4). The inferior fornix can be traced as far as the

DIAGRAM 4



atrio-ventricular node (the starting point of the A-V bundle of His—see below). This fact has an important bearing on the transmission of the waves of contraction of the heart. The rudimentary valves at the mouth of the inferior vena cava and coronary sinus represent remnants of the right sinus valve, the crista terminalis of the right atrium being its persisting muscular rudiment. The left sinus valve takes part in the formation of the limbus fossa ovalis on the inter-atrial septum.


The portions of the adult atria which are derived from the sinus venosus are the smooth posterior areas where the great veins open, the

anterior regions, including the auricular appendages where the musculi pectinati occur, being developed from the atrial portion of the cardiac tube. In each atrium the junction between the morphologically different elements entering into its composition is indicated by a ridge, the *crista terminalis*. The musculature of the sinus is a very specialised tissue containing a large number of nerve cells. In the course of development towards the adult stage of the human heart, the muscle disappears entirely except at the mouth of the superior vena cava and the atrial wall in front of it. This remnant of sinus musculature constitutes the *sino-auricular node*, the "pace-maker" of the heart.

2. *The atrium (auricle).*—To start with, it consists of one undivided chamber, continuous caudally with the sinus venosus, and cranially with the ventricular division of the embryonic cardiac tube through the atrial canal (see Diagram 3). It grows dorsally, while, as we shall see presently, the ventricle expands in the opposite direction, i.e., ventrally. A partition grows down from the dorsal wall, and, advancing towards the atrio-ventricular region, ultimately fuses with the endocardial cushions which are forming there (see Diagram 4). This partition is called the *primary septum*. The temporary aperture between the advancing primary septum and the atrio-ventricular endocardial cushions is called the *primary foramen ovale* marked X in Diagram 4. This aperture disappears as a normal event. If it persists, as it does rarely, there will be free communication between the right and left atria. Before the primary septum meets the atrio-ventricular cushions, the dorsal portion of it breaks down producing the (*secondary*) *foramen ovale*. A second partition, having a crescentic free border, now appears on the right side of the first (see Diagrams 4 and 5). The anterior

and posterior free ends of the crescent fuse with the atrio-ventricular cushions, but the intermediate part is simply applied to the right face of the primary septum without fusing with it, so that blood can pass from the right to the left side of the atrium through the (*secondary*) foramen ovale. This state of things continues normally till birth, when in the majority of individuals the secondary septum blends with the first obliterating the (*secondary*) foramen ovale permanently. The primary septum forms that part of the interatrial wall of the adult which lies at the bottom of the fossa ovalis, and the secondary septum gives rise to the limbus fossa ovalis.



DIAGRAM 5



Showing the formation of the various septa of the heart.
Right face of the septa drawn.

grow. This fact is indicated throughout adult life by the interventricular sulcus. The septum, in the first instance, is an incomplete one and, consequently, the two compartments of the ventricle communicate with each other above the free upper border of it. This communication is called the *interventricular foramen*, and is ordinarily a temporary opening only (see Diagram 5). It is closed by the fusion of the ventricular septum with the bulbar septum (see below) and the atrio-ventricular endocardial cushions. The structure which thus fills the gap forms the membranous portion of the interventricular septum in contradistinction to the muscular part, the remaining and major subdivision of it. The interventricular foramen may not be closed owing to the non-fusion of the various embryonic rudiments mentioned just now. Such a defect constitutes one of the most common malformations of the heart and occurs usually with congenital stenosis of the pulmonary artery. The remaining unabsorbed portion of the spongy interior of the ventricular rudiment is transformed into the chordae tendinae, muscoli papillares, trabeculae carnae, and, in the case of the right ventricle, the moderator band in addition.

DIAGRAM 6

(a)
Heart in Third Week(b)
Heart in Third Month


1. Sinus venosus.
2. Atrium.
- 3' Left ventricle. 3' right ventricle.
4. Bulbus cordis. 4' Part of right ventricle derived from bulbus cordis. 4' Part of left ventricle derived from bulbus cordis.
5. Truncus arteriosus.
- 5' Pulmonary artery. 5' Aorta.
- 6 is placed between the contiguous walls of bulbus cordis and ventricle. Note that these walls have been absorbed in stage reached in (b).
7. Anterior cusp of mitral valve which intervenes between the mitral orifice and the aortic opening.
8. Interventricular septum.

5. *Bulbus cordis*.—This is a separate unit in the embryonic cardiac tube distinctly marked off from the ventricle proximally and the truncus arteriosus distally (see Diagram 3). But in the course of its further development, it is absorbed into the ventricles, chiefly the right. We have seen that a similar fate overtakes the sinus venosus, which, losing its distinctness as such, is merged into the atria. A series of changes, too complicated to be described here in detail, take place in the bulb whereby (a) the contiguous walls of the bulb and the ventricle are absorbed resulting in the union of the two cavities (see Diagram 6), and (b) a septum called *bulbar septum* is developed inside this subdivision of the primitive heart, which joins up, proximally, with the interventricular septum and the posterior of the two endocardial cushions in the region of the atrio-ventricular canal (see Diagram 5) and, distally, with the spiral septum (see below) coming into existence inside the truncus arteriosus. This septum in the course of the bulb is instrumental in dividing it into two parts, a bigger one added to the right ventricle and a smaller to the left. The respective areas of the adult ventricles derived from the bulbus cordis are the infundibulum, in the case of the right, and the aortic vestibule, in that of the left ventricle. Again, the territorial boundary between the bulbar and ventricular rudiments of the primitive tube is indicated by the position of the moderator band. Defective transformation of the bulb into the infundibulum is responsible for congenital pulmonary stenosis, which, together with a persistent interventricular foramen—an associated condition—furnishes by far the greatest number of congenital anomalies of the heart.

6. *Truncus arteriosus*.—By the development and fusion of endocardial thickenings in this subdivision of the cardiac tube, a spiral septum comes into existence along the whole length of it which fuses proximally with the bulbar septum. The truncus is thus divided into two channels, the pulmonary artery in front and aorta behind (see Diagram 6). The normal disposition of the septum is such that the aorta gets connected with the left ventricle and the pulmonary artery with the right. If the spiral is directed in the opposite way the aorta will be in communication with the right ventricle and the pulmonary artery with the left. Rarely, the proximal end of the septum is placed at right angles to the ventricular septum, i. e., midway between the first two positions. In that case, each ventricle on contraction will be pumping blood into both the arteries. The morphological junction of the bulbus cordis and truncus arteriosus in the adult heart is the region of the aortic and pulmonary valves.

With the formation of the spiral septum in the course of the truncus arteriosus and its joining up with the bulbar septum, the heart is divided completely into right and left parts, the right part subserving the pulmonary circulation and the left one the systemic. When once the partition is effectively formed the communication between the two halves is only an indirect one, viz., through the capillaries of the lungs. It will be useful to recapitulate the various elements of this remarkable partition or septum. They are (a) the interatrial septum composed chiefly of the primary and secondary septa fusing with (b) the atrio-ventricular endocardial cushions which in their turn blend with (c) the interventricular septum and (d) the bulbar septum. Lastly (e) the spiral septum of the truncus arteriosus comes in and, joining up with the bulbar septum, completes the partition.

The Formation of the Semilunar Valves, aortic and pulmonary. Four thickenings, anterior, posterior, right and left, arise in the course of the endocardium lining the cardiac tube at the junction of the bulbus cordis and the truncus arteriosus, as shown in Diagram 7 (a). The spiral septum



mentioned in connection with the truncus cuts across the right and left thickenings and divides each of them into an anterior and a posterior part [see Diagram 7 (b)]. The anterior divisions of the right and left thickenings together with the undivided anterior thickening get converted into the semilunar valves of the pulmonary arteries. Similarly, the two posterior divisions together with the unsplit posterior thickening get fashioned into the valves of the aortic opening.

The Atrio-Ventricular Bundle (of His)—In lower vertebrates, the atrial or atrio-ventricular canal is surrounded by a collar of muscle tissue which is continuous proximally with the musculature of the atrium, and distally with that of the ventricle. This tissue is responsible for the transmission of the waves of contraction from the atria to the ventricles. In the mammalian heart, including the human, this muscular collar entirely disappears and is replaced by a fibrous ring, except in one place where a small fasciculus of the original muscle remains. This remnant now forms the only muscular connection between the atrial and ventricular portions of the heart. In the adult stage, this strand (A - V bundle of His) takes origin in a special group of cells called the *atrio-ventricular node* situated in the septal wall of the right atrium near the opening of the coronary sinus. Therefrom the bundle proceeds distally along the upper border of the interventricular septum and divides into right and left branches for the corresponding ventricles. Reference has already been made to the fact that the right sinus valve arising in the region of the sino-auricular node (pace-maker of the heart) ends in the posterior of the two endocardial cushions of the atrio-ventricular canal. This actually is in the close vicinity of the node, the starting point of the A-V bundle.

Reference may now be made to an embryonic structure which, though not strictly a part of the primitive cardiac tube, comes into importance in association with developmental defects of the organ. That structure is the *ductus arteriosus*. The vessel in the embryo short-circuits most of the blood in the pulmonary artery to the aorta, there being no necessity at that stage for all the pulmonary blood to circulate through the lungs. That being no longer the case after birth, i. e., as it is imperative for all the blood in the pulmonary artery to circulate through the lungs, there is no further need for this embryonic channel. It should disappear. Normally, it is reduced to a fibrous remnant connecting the aortic arch with the left branch of the pulmonary artery. In company with other abnormalities this channel may remain open.

Having thus reviewed the embryology of the heart we shall now consider some of the common abnormalities.

Persistent primary foramen ovale.—This is due to failure of union between the primary septum and the atrio-ventricular endocardial cushions. The defect may be so gross as to involve the whole of the partition between the atria, with the result that there will be only one chamber in the atrial division of the heart.

Patent (secondary) foramen ovale.—It may occur alone or in company with other defects and is due to the non-fusion of the primary and secondary septa. In a considerable proportion of adult bodies this defect is noticed—not usually during life, but accidentally at post-mortem examinations. In other cases it is a compensatory arrangement in connection with defects like congenital pulmonary stenosis.

Atrio-ventricular passage.—Instead of two canals, there may be only one undivided passage due to failure of fusion between the two endocardial thickenings of that region of the primitive cardiac tube.

Congenital pulmonary stenosis.—This is one of the most, if not the most, common of the congenital defects of the heart and is the result of faulty development of the bulbus cordis. There is invariably associated with it a cavity of considerable size in the region of the infundibulum of the right ventricle. As a rule, the interventricular foramen is patent and blood, therefore, can pass from the right ventricle to the left. The ductus arteriosus is also open in such cases and part of the aortic blood goes that way to the lungs for their nutrition. The state of the ventricles will differ materially according to whether the stenosis takes place before the interventricular foramen is closed or after.

If before the right ventricle gets hypertrophied and the left atrophied, the septum is shifted to the left and the aorta is wholly or partially connected with the right ventricle, the foramen ovale and the ductus arteriosus are open. The lungs get blood from the aorta through the ductus arteriosus and not to any extent from the pulmonary artery. The whole of the systemic and almost the entire pulmonary circulation are carried on by the right ventricle.

If, on the other hand, the stenosis occurs after the interventricular foramen is closed, the left ventricle enlarges and hypertrophies; and the foramen ovale and the ductus arteriosus remain patent.

Atresia or complete obliteration of the pulmonary artery may occur, but is a much rarer condition than the preceding. If it occurs after the interventricular foramen is closed, the right ventricle atrophies and ultimately the right atrio-ventricular opening gets obliterated.

The interventricular septum may be absent wholly or in part. Partial deficiency occurs in the upper portion where the membranous rudiment has not developed, and an opening, the interventricular foramen, is left in its place. But sufficient has been said about it already.

Complete absence of the septum is rarer. When that happens, the ventricle is composed of a single chamber as is the case in the frog.

Stenosis and atresia of the aorta are rarer forms of congenital defects than those of the pulmonary artery. Anyone of three points in the course of the artery may be affected: at its origin from the left ventricle, just proximal to its junction with the ductus arteriosus, or just distal to the last. In the *first* variety the left ventricle remains undeveloped and the systemic circulation is mostly carried on by the pulmonary artery through the ductus arteriosus. The *second* type occurs along with other malformations such as pulmonary stenosis, and most of the systemic circulation is done by the right ventricle through the ductus arteriosus which remains open. The *third* kind is more common. The aorta is stenosed below the junction with the ductus arteriosus. A collateral circulation develops in these cases and compensates the defect in some measure.

An *abnormal communication* may be present *between the aorta and the pulmonary artery*, the result of defective development of the spiral septum in the truncus arteriosus.

The abnormal positions of the aorta and the pulmonary artery in relation to the ventricles, in certain cases, have been mentioned already. No satisfactory explanation of such phenomena is forthcoming. *Congenital defects of the atrio-ventricular valves* may be present, the number of cusps being either increased or decreased. The tricuspid valve is more commonly affected than the mitral.

Transposition of the heart to the right side, constituting the condition called 'dextro-cardia,' may occur. It is almost invariably a part of a general reversal in position of the organs and is detected either on X-ray examination or at autopsies.

Lastly, a word might be said about a rare condition called 'ectopia cordis.' Here, the organ lies outside the thoracic cavity. It may be placed in the neck, in front in the chest, or in the abdomen. When it is in the pectoral region, the sternum is found to be cleft in the middle line or partially absent, the rudimentary organ protruding through the deficiency. In extreme cases, only the dorsal wall of the pericardium is present, which is flush with the general surface of the body. Such cases remind one of 'ectopia vesicae' where the whole of the urinary bladder except the trigone is missing and the trigone itself with the orifices of the ureters presents at the surface.

GLEANINGS FROM MEDICAL LITERATURE

21

EPILEPSY AND ITS TREATMENT

JAMES COLLIER

The Practitioner, July 1934, p. 37.

Epilepsy is common in birds and mammals. Heredity is a factor in the transmission of the disease but it is not so great as is usually supposed. The malady may begin at any period of life, but occurs with greater frequency in early childhood, with the onset of puberty, and in old age. It is generally classified as *organic or symptomatic epilepsy* and *idiopathic epilepsy*. Epilepsy-like phenomena which follow local diseases of the brain, kidney, thyroid and liver, those occurring in infantile malnutrition, infection and intoxication are called symptomatic or organic epilepsy.

Recent investigations point to the conclusion that all epilepsies are the result of one and the same underlying fault in metabolism, and that local lesions of the brain, uraemia, and intoxications act as secondary factors in raising the epilepsy-producing potentiality of this metabolic fault. It seems that the community might be divided into three classes: (1) Those in whom no trace of metabolic fault is present—they cannot be provoked into epilepsy by local lesions of the brain or other organs or intoxications; (2) Those in whom the metabolic fault is present in minor degree—in them a second factor such as disease of the brain will cause epilepsy; (3) Those with the fault in high degree—they are the idiopathic epileptics. The clinical facts that make the above hypothesis acceptable are: (1) the periodicity of the attack and their association with sleep and moments of bodily and mental inactivity when oxidative processes are deemed to be low; (2) occurrence of fits in certain cases for a few days followed by freedom from attacks for months and years, the explanation being that the fault is present in sufficient degree only at the period when the fits occurred; (3) epileptic women are nearly always free from epilepsy during pregnancy, it being supposed that the foetus corrects the metabolic fault of the mother.

This evidence of a metabolic dyscrasia is not very clear or convincing. In epilepsy, when the attacks are frequent, the alkali reserve is increased and there is great variation in the value of the blood pH from day-to-day. O. Foerster states that in epileptics hyperventilation of the lungs by so simple a method as making the patient breathe deeply as fast and as long as he can will bring about an attack at once by the excessive removal of carbon dioxide. It has been noted that a condition of *status epilepticus* develops in patients not previously epileptic by the use of alkalis in the treatment of gastric ulcer. Epilepsy is incompatible with acidosis, and this may account for the rarity of epilepsy in diabetics. The good effect of a ketogenic diet seems to suggest that alkalosis plays some part in the production of epilepsy.

A sudden drop in blood pressure precedes or accompanies an attack of epilepsy. This is seen clinically as a momentary facial pallor. It was observed by Foerster Kennedy as a sudden blanching of the brain when a general epileptic convulsion occurred during an oper-

ation in which the surface of the brain was widely exposed. Epilepsy seems to be a protective phenomenon to avert a dangerously falling blood pressure and dangerous alkalosis as the muscular convulsion with the arrest of respiration concentrates the carbon dioxide ratio and lessens the alkalosis. Epileptics belong to the vagotonic group of Eppinger and often manifest such vagotonic phenomena as urticaria, asthma, and migraine.

The hypothesis that the epileptic attack is an expression of excitation of the brain has been abandoned, and, with Hughlings Jackson, it is regarded as due to loss of function, local or general, in the brain. The positive events such as hallucinations and convulsions are regarded as release phenomena in lower regions of the nervous system following loss of function of the higher regions.

Treatment.—Epilepsy is fostered by inactivity of mind and body and warded off by activity. Hence, the excitation theory of the later Victorian period, which advocated rest and quiet, has given place to one of activity of body and mind. When such conditions as rickets, renal diseases, are present, they must be corrected. A ketogenic diet, rich in protein and fat and deficient in carbohydrate, is definitely beneficial. The medicinal treatment of epilepsy consists in the exhibition of Luminal, Bromides, and Paraldehyde, and in rare cases of Zinc and Belladonna. To get the maximum benefit, the drug must be exhibited before the anticipated attack, and if the remedies are not effective in ordinary doses, they will not be effective even in larger doses. Luminal is the drug of choice and the maximum dose for the 24 hours must not exceed 3 grains. Bromides may be used when Luminal fails or as an adjunct to Luminal; the maximum useful dose is not more than 30 grains per day for an adult. Where these fail, Zinc oxide, in 3 to 5 grains doses, in capsules, and Tincture Belladonna, in 10 minim doses, may be tried. Paraldehyde in four drachm doses dissolved in twice the amount of olive oil per rectum is effective in cases of *status epilepticus*.

V. G.

22

NERVOUS BREAKDOWN

C. P. SYMONDS

Guy's Hospital Gazette, Feb. 2, 1935.

The term covers an illness, or episode of disease, which is mental rather than physical, characterized by disorder or break-down at the highest level of nervous function.

Symptoms:

- (1) *Depression*, with or without adequate cause, out of proportion to the facts of the patient's life.
- (2) *Morbid feeling of fear*—free or attached—a dreadful feeling of fore-boding, as if something awful is going to happen, a feeling that he must have cancer or syphilis or heart disease.
- (3) *Negative complaints*—absence of normal sensation of pleasure in relation to work, play, food, drink, or tobacco; loss of sexual feeling and sense of mental fatigue.
- (4) *Bodily symptoms*: fatigue occurring easily, pains in the back and limbs, headache, flatulent dyspepsia, constipation, palpitation, and insomnia are the common symptoms. The

symptoms are subject to many variations, depending on the degree of the morbid feeling, and the personality and character of the individual. There may be conviction of ruin or guilt, or certainty of cancer or syphilis, or negative symptoms may predominate. The patient is totally apathetic and indifferent.

The illness has a gradual or sub-acute onset, a subsequent development by degrees to its maximum, a prolonged course and a gradual recovery.

Etiology: The family history supplies the main factor. Individual attacks may be precipitated by emotional stress, excessive fatigue (mental or physical), insomnia (e. g., from pain), or intoxication (e. g., bacterial—influenza; chemical—alcohol). But in 90 per cent of cases no physical or emotional cause is to be discovered. It is the fashion now-a-days to call the mild cases of this illness *psychoneurotic depression* or *anxiety neurosis*, and at some point or other to draw a line separating these from the severe cases, which are called psychotic, manic-depressive, or melancholic. The distinction is usually held to imply that the two groups are of different nature and causation. In this sense it is wrong.

Treatment.—

(1) Radical: There is no short cut to cure in this disease.

a. Weir Mitchell, assuming that the causation of the illness was exhaustion and undernutrition of the nervous system, treated these with rest in bed, isolation and superabundance of milk. The difficulty of assessing the value of any such theory or form of treatment arises from the fact that these patients almost always get well.

b. Exponents of intestinal stasis or focal sepsis had their day (not to mention endocrinologists and osteopaths).

c. Freud and his disciples argued that these patients deny an adequate cause for their depression or anxiety, which really exists in the unconscious mind. If, by means of a special method of psychological analysis, the patient can be made to bring his conflicts into consciousness, he would make a satisfactory emotional adjustment and will recover. This method of treatment is a failure: patients treated by the analytical method do not get well any more quickly than those treated by other methods, nor are they any more free from the liability to recurrence. On the whole, it has done more harm than good in that it has inflicted a good deal of unnecessary mental discomfort upon patients already miserable with their illness.

(2) Symptomatic: The first principle in the treatment is to tell the patient that he is suffering from a genuine illness from which he will get well, and give it a name such as neurasthenia, nervous depression, or nervous breakdown.

The next step is to assess the degree of the illness or resources of the patient. In a mild and early case, a complete and restful change for a week or ten days, with adequate hypnotics, is worth trying. In the case of a man keep him at his work if you care, watching his ability to carry the load. One man complains easily, another may go on to the point of suicidal despair without saying much about it. The former needs to be pushed, the latter spared. Increasing insomnia is an indication for lightening the burden.

The patient should be encouraged to talk to his doctor of any worries or troubles with which he is pre-occupied, and to accept advice and help in the management of current problems. Some near relative, husband, wife, parent or friend, should be interviewed, informed of the nature of the illness, and enlisted as ally. These patients need, and deserve, understanding and sympathy, though they may often need encouragement and persuasion as well. Nearly all these patients are better for some kind of physical treatment, be it medicine, helio-therapy, or massage. Changes in the form of physical treatment are obviously necessary from time to time. The treatment of insomnia is an important point, and *medinal* is the most useful. Of medicines, the old fashioned mixture of *bromide and strychnine* is valuable. A half-grain tablet of *luminal* as a remedy against intense waves of restless depression may have a physical as well as a psychological value. Alcohol as a refuge from depression is a danger; as an aid to digestion, it is useful. When sexual appetite is diminished or lost, the attempts at coitus are harmful, a fact of which both partners should be made aware.

(3) Preventive: When the liability is recognised much may be done in the way of preventive treatment. In childhood, the patient should learn to endure pain both physical and mental. So many boys are over-burdened at puberty with unnecessary fears. They believe that they are alone in their failures, that they have committed the unforgivable sin, that they are candidates for consumption or insanity. So much can be done for them by a frank exposition of the subject. On the physical side it should be remembered that a depressive illness may be precipitated by fatigue, infection, intoxication, or simple insomnia from physical pain or mental distress, and, in women, from childbirth. There is no doubt that, apart from the danger of habit formation, alcoholic excess may precipitate an attack.

K. L. N. R.

23 ACUTE INFECTIONS OF THE UPPER RESPIRATORY TRACT, MIDDLE-EAR AND MASTOID

J. R. PAGE and H. C. BALLENGER

Archives of Otolaryngology, Oct. 1934.

The authors have attempted to picture the correct pathological background underlying the irregular and apparently erratic course of acute infections of the upper respiratory tract and its appendages. Their observations are based on elaborate bacteriological investigations during acute infections of this region in over 300 cases, and collateral findings of other co-workers. Their investigations included (1) cultures in each case of material obtained from strategic points along the tract, grown on a divided blood plate containing a special medium of brain, veal-agar and brain-heart infusion broth with normal blood serum, using small applicators through sterilized specula to avoid contamination, and (2) blood cultures during the height of the disease. They found that *Streptococcus haemolyticus* was accountable for nearly 31 per cent of these infections, while *Pneumococcus* Type III accounted for nearly 17 per cent of the cases. *Haemolytic Streptococcus* and *Pneumococcus* Type III occur more frequently than any other organism in infections of the middle ear and mastoid, and this necessitates a guarded prognosis because of the insidious and virulent character of such infections, and because of the severe involvement of the mastoid. The degree of mastoid involvement, however, does not solely depend on the virulence of the infection, but on the

147296

patient's resistance as well, and on the anatomy of the particular mastoid which may retain or expel the infection. Too early an operation on a "green mastoid" is to be avoided; so also delay until the inner plate has softened. On the merits of myringotomy they found that almost three times as many operations on the mastoid were necessary in cases where the drum was allowed to rupture spontaneously as in those cases in which myringotomy was performed before rupture occurred.

The pathways of infection of the haemolytic *Streptococcus* from the nose and throat are—

- (1) Direct extension by continuity of tissue as exemplified by mastoiditis and sinusitis.
- (2) Extension by lymphatics as exemplified by cervical adenitis and retropharyngeal abscess.

- (3) Bacteraemia or a blood-borne infection from the upper respiratory tract.

This bacteraemia is difficult to demonstrate by the usual methods of blood culture. The difficulties encountered may be—

- a. The bacteria are not present in the quantity of blood with-drawn.
- b. The showers of bacteria may not be present in the blood at the time the cultures are taken.
- c. Improper technique, e. g., wrong culture media or too short a period of cultivation.
- d. The filtration system of the blood stream, that is, the liver, spleen, lungs, finer capillaries, or the phagocytic cells (particularly of the reticulo-endothelial system) may remove most or all of the bacteria before they reach the vein of the arm or neck from which blood is usually taken for culture. There is experimental proof to show that the liver and spleen remove enormous numbers of bacteria injected into the blood stream of animals.
- e. Recent studies in pleo-morphism and transmutation or dissociation of bacteria may explain the lack of positive blood cultures in many instances. If this is true, recognised strains may be in a filterable phase or may react unusually to the common staining methods.

The results of positive blood-cultures in various acute and chronic infections of the upper respiratory tract by different observers are not uniform on account of the above mentioned difficulties, but the importance of the positive results should not be under-estimated. If cultures had been made with blood from an artery the positive results would probably have been greater. Many infections of the kidneys, heart, brain and other vital organs, occurring during these upper respiratory tract infections, can be explained only by blood stream transmission. Admitting that the majority of these infections are due to haemolytic *Streptococcus*, and also admitting the probability of a bacteraemia from this source the following deductions can be made:—

1. With this apparent tendency of haemolytic streptococci to invade the blood stream or extend to the neighbouring regions during the acute stages of the infections, it is obvious that operations on the nose and throat should be deferred until a quiescent period is reached.
2. When, however, the infection extends to the middle ear, early paracentesis, as soon as bulging of the drum is seen, is indicated,

3. If mastoiditis develops, the question of operative intervention and especially when to do it, must be decided, each case being judged on its own merits. If one can wait three weeks from the onset of discharge with reasonable safety, it is desirable to do so, giving Nature a chance to throw up her protective barriers.

4. A transitory or an evanescent type of bacteraemia is fairly common, and is possibly a usual complication following an acute infection of the upper respiratory tract. Unexplained fever, out of all proportion to the degree of involvement of the throat, can be explained by this bacteraemia, the possible results of which are limited only by the distribution of the blood-stream.

M. S. R.

24 RADIATION IN THE TREATMENT OF CANCER

A. C. CHRISTIE

Journal of the American Medical Association, CIII, 1934, 985.

The rational basis for radiation therapy goes back to the early discovered fact that embryonal types of cells are more radiosensitive than adult forms. The shorter the life-cycle of the cell, the more sensitive it is to radiation, and *vice versa*. Among other factors that are equally important are the general condition of the patient and his reaction to radiation, the immediate tissue environment of the cancer (the cancer bed), and the relation of the growth to vital organs or important tissues that may suffer injury.

Except in superficial cancers in which it is possible to destroy the growth by the caustic effect of radiation, the single massive dose method has been practically abandoned. Some type of fractional dose is now in general use. The "saturation method" of Pfahler consists of administering a large initial dose and maintaining the radiation effect at a maximum by giving a small daily dose until the total dosage is attained. The Coutard fractional dose method of applying roentgen rays consists in giving a large total dose over a prolonged period (from 18 to 35 days), an equal amount being given daily until the total dose has been reached. Results have been obtained by this method which are superior to any previously thought possible. Holfelder has used a method that aims to take advantage of the large initial dose of the "saturation method" and subsequently to build up the dosage by the method of Coutard. The increased benefits from the fractionated dose method appear to arise from improvement in the ratio between the destructive effect on the neoplasm and the reaction in the surrounding normal tissues.

Progress has also taken place in methods of radium therapy especially in respect to interstitial radiation. The use of gold and platinum seeds was a great advance over glass seeds for the interstitial application of radon; the increased filtration lessens the danger of localized necrosis. The use of tubules, or needles from 3 to 6 cm. in length, with a very small amount of radon or radium element to the running centimeter, in the form of a grid, has secured a fairly uniform dosage. Further advance may be made in roentgen apparatus by securing

a much greater energy output by the use of tremendously higher milliamperage rather than by an increase in voltage.

Because of results obtained in large numbers of primarily inoperable and seemingly hopeless conditions and in recurrences after operation, and because of improvements in results when used in addition to surgery, the attitude towards radiation has radically changed during the past three or four years. Cancers of the skin have been treated by radiation methods almost from the time of the discovery of the roentgen ray. The basal-cell type, when it is small and does not involve cartilage or other important structures, is readily curable. The fractional method of Coutard constitutes an epochal advance in dealing with lesions involving cartilage. The primary treatment of squamous-cell epithelioma should be a thorough irradiation of the entire lymphatic drainage areas by the fractional method. The added danger of dissemination must be weighed when biopsy is contemplated; it should be postponed until about two weeks after roentgen irradiation to lessen the liability of spread. If there is any residuum of the primary lesion it can be destroyed by electrocoagulation or by implanting interstitially gold or platinum radon seeds or small needles of radium element. If excision is to be done at all, it should always be preceded by irradiation.

K. R. C. H.

25 ENDOCRINE EFFECTS OF CERTAIN OVARIAN TUMOURS

EMIL NOVAK

The American Journal of Medical Sciences, CLXXVII, 1934, 599

Certain tumours of the ovary, like tumours of other endocrine glands, are capable of highly developed endocrine functions. The two most clearly defined types are the granulosa-cell tumours and the so-called arrhenoblastomata. The latter belong to the group of testicular adenomata, and are most often of atypical variety.

The granulosa-cell tumours exert a feminizing effect, through the production of theelin by the tumour cells, so that in older women, perhaps far beyond the menopause, they produce most often a hyperplasia of the endometrium associated with periodic bleeding (pseudo-menstruation), together with increased size of the uterus. In the few cases seen in very young children, they have produced the syndrome of precocious puberty, with precocious menstruation.

The arrhenoblastomata, on the other hand, have a definitely masculinizing tendency, as might be expected from the fact that they apparently have their origin from undifferentiated epithelium in the region of the rete ovarii. Under conditions which are not clear, these cells, capable of development along either male or female lines, may assume definitely masculine tendencies. The relation of this endocrine cause of partial sex-reversal to the general problems of intersexuality, sex differentiation, and sex reversal is obvious, although our knowledge on these subjects is still far from complete. Patients suffering from arrhenoblastomata present in extreme instances such evidences of masculinization as a masculine type of hair distribution, deepening of the voice, and hypertrophy of the clitoris. Removal of the tumour brings about a regression of these symptoms.

K. R. C. H.

THE TREATMENT OF ANGINA PECTORIS

K. SHIRLEY SMITH

The British Medical Journal, 1935, i, 988.

Genuine and severe pain in the centre of the chest is most probably due to some form of angina pectoris, the pain being an expression of the myocardial ischaemia. This ischaemia may be temporary as during exertion when the heart muscle requires extra blood, when it is called *angina of effort*; this is by far the commonest form of angina. Or, it may be *paroxysmal* and arising independently of effort. Permanent arrest of blood supply (coronary thrombosis) leads to infarction of the heart muscle, and in this condition the pain is prolonged for hours or days.

The treatment of angina is more the ordering of a mode of life than a therapeutic system. Work, recreation, hours of rest, diet, habits and temperament—all need investigation.

Mode of life: Such activities as do not induce pain are allowed. The degree of attack-producing exertion is remarkably constant and an intelligent patient frequently learns this and keeps within the limit. Strenuous exercise is absolutely forbidden. Sources of worry, anxiety and fatigue should be removed, and adequate rest—a minimum of 10 hours—enjoined.

Diet: Moderation in dietary without irksome restrictions should be aimed at. Meals should be small in bulk and taken with a minimum of fluid. In the spare, active type of patient, a tablespoonful of medicinal glucose in powder form may be given with the principal meals. Alcohol is best avoided and exercise after meals should be strictly forbidden.

Rest: Angina, becoming increasingly frequent and severe, or having origin in a recent coronary thrombosis, or supplemented with night attacks, or associated with active myocardial disease, should mean confinement in bed for a period of 2 to 3 weeks.

Medicinal Treatment:

During an attack: A premonitory pain or sense of constriction during exertion should call for immediate cessation of the work while a tablet of 1/100 gr. trinitrin is taken; this may be all that is needed to bring relief. For more severe pain an amyl nitrate capsule (5 min.) is inhaled. If the attack persists for an hour or more in spite of nitrites, coronary thrombosis has probably occurred and the patient should be put to bed and an injection of morphine, gr. $\frac{1}{4}$, given.

To prevent attacks: Over-exertion is avoided. Attacks tending to occur at particular times of the day are combated by taking trinitrin a quarter of an hour previously. Theobromine sodium salicylate, in 10 gr. doses, thrice daily, may be given as a routine, while in highly strung patients a combination of theobromine, gr. v, with luminal-sodium, gr. $\frac{1}{2}$, may be useful.

For other symptoms: For a nervous and excitable patient a routine sedative such as potassium bromide, gr. x, with luminal gr. $\frac{1}{2}$, at bed time, or twice daily, may be given.

Insomnia may be met by a draught containing potassium bromide gr. xx, and chloral. gr. x, or chloral hydrate gr. xx, or 3 drachms of the syrup. For flatulence a pill containing calomel, gr. 1/6, menthol gr. 1/4, and powdered ginger, gr. 1/4, may be taken thrice daily after food. For constipation, which it is important to avoid, liquid paraffin (with or without phenolphthalein), cascara evacuant, or senna tea will be found useful.

Drugs to avoid are adrenaline, ephedrine, and thyroid extract. Smoking is unwise.

Treatment of associated disorders: (1) *Obesity* presents particular difficulty since both exercise and thyroid medication cannot be used to reduce it. Regulation of diet should be mainly depended on to reduce weight. *Flat-foot* and other minor foot troubles associated with obesity will throw strain on the heart and should be corrected. (2) *Septic Foci* should be eliminated. (3) For *high blood pressure*, the above regime would itself be adequate. If persistently high, venesection is indicated and may be repeated if it is beneficial. (4) *Angina* associated with thyrotoxicosis should be treated by total thyroidectomy, the resulting myxoedema being subsequently corrected by thyroid medication. (5) In *angina with diabetes*, the coronary syndrome should claim prior attention, and it is wisest not to interfere too much with the diabetic condition, the patient being allowed to pass slight sugar and be mildly hyperglycaemic, since any undue lowering of the blood sugar by insulin may interfere with the nutrition of the myocardium and may precipitate infarction and sudden death. (6) *Anaemia* of the pernicious type will need liver therapy; for the microcytic or nutritional form *341 Ferri*, gr. xxx t.i.d., or *Ferri et Ammon*, Cit. gr. xl, t.i.d., will be suitable.

Surgical Treatment is still largely in the experimental stage, although improvement has followed both paravertebral injection of alcohol and cervical or dorsal sympathectomy.

M. K.

CLINICAL MEMORANDA

TREATMENT OF SMALL-POX.

V. GOVINDAN NAIR, M.R.C.P (E.), F.R.F.P.S.,
King George Hospital, Vizagapatam.

I have found injectable liver extract useful in the treatment of small-pox. It modifies the course of the disease, shortens the duration, aborts the eruption at all stages and avoids the consequent disfiguration from deep scarring and pitting. It also controls the toxic symptoms, improves the general health and well-being of the patient and prevents the development of the usual complications of untreated cases of small-pox as boils, subcutaneous abscesses, toxic rash, phlebitis, thrombosis, etc.

The treatment proper consists of daily intramuscular (intragluteal) injections of liver extract till the eruptions desiccate and incrust. Five c. c. of the extract should be given for the first two or three days according to the severity of the case, followed up by two c. c. daily for the remaining period. The extracts used by me were (1) Campolon of Messrs. Bayer—Meister—Lucius and (2) Injectable liver extract of Messrs. Bengal Chemical and Pharmaceutical Works, Ltd.

For full particulars please refer to the article entitled "Treatment of small-pox with Liver" in the *Journal of the Indian Medical Association*, July 1935.

REVIEWS

MATERIA MEDICA OF PHARMACEUTICAL COMBINATIONS AND SPECIALITIES. By U. B. Narayanarao. Bombay: The Ryan Pharmacy, 1932. Pages xliv plus 249.

This volume is a compilation giving particulars of the various proprietary medicinal preparations on the market. While we do not deny that there is a need for such a publication to assist the prescribing physician in the choice of the particular proprietary medicine he would employ in any given case of disease, we consider that the present volume does not furnish the necessary information on which the physician could make a proper choice. For, the details given with regard to the preparations listed are only those furnished by the respective manufacturers, which in most instances are likely to be exaggerated, and no effort has been made to indicate how far in actual medical practice, as recorded in current medical literature, these preparations have proved the claims made for them. In the extensive medical literature of the present day, notes on, and references to, proprietary preparations are found in abundance, and their therapeutic indications and efficacy are discussed. We suggest that in listing proprietary medicinal preparations in a *Materia Medica* references be made to such notes and discussions in the literature so as to enable the medical practitioner to decide which preparation is best suited in any given case. The absence of such references detracts considerably from the usefulness of the present volume.

THE MADRAS MEDICAL DIRECTORY, 1935. Published by T. S. Lakshmipathy, Madras.

The second edition of the *Madras Medical Directory* published by Mr. T. S. Lakshmipathy is a distinct improvement on the first edition. The volume is much increased in size by the addition of matter which should be very useful to the public and the profession. There is great need for a dependable volume of reference where the addresses and other particulars of medical men scattered over all parts of the Presidency could be readily obtained. That would be the first step in the organization of the profession. We hope, therefore, that the medical practitioners and others concerned would help the publisher by furnishing him with the change of address and other particulars which may occur to their knowledge. We trust that this Directory would be published annually with all the necessary alterations and improvements.

CALCIUM THERAPY

Alembic Alcalzium Tablets (The Calcium Food) provides, in palatable form, a Scientific combination for use in all conditions requiring intensive Calcium Therapy.

Calcium Sodium Lactate has been chosen because:
1. It is highly assimilable and palatable. 2. It does not tend to increase the acidosis so commonly associated with conditions needing *Calcium Therapy*.

No *Calcium Salt*, however, is adequately assimilated except in conjunction with a sufficient supply of *Vitamin D*.

The presence of *Iron* accelerates the regulating of body-metabolism.

Alcalzium with Iron & Vitamin D is valuable for Growing Babies, Building strong bones and Teeth and maintaining proper Heart function.

During Pregnancy *Calcium* is required from the beginning not only for the mother but also for the growing baby.

It is also useful for Anaemias, Rickets, Deficient Growth, Disorders of Teething, Convalescence after Fevers.

Each Tablet of **Alembic Alcalzium** with *Iron & Vitamin D* contains:

CALCIUM LACTATE	1.75 Grains.
SODIUM LACTATE	1.25 "
DICALCIUM PHOSPHATE	2 "
FERRIC AMMON CITRAS	1 "
VITAMIN D.	100 Units.

Available at all Chemists in bottles of 50 Tablets.

Alcalzium
With Iron & Vitamin D.

ALEMBIC CHEMICALS

BOMBAY

BARODA

MADRAS

Facts WHICH SPEAK for THEMSELVES!



COW & GATE Milk Food is now supplied to practically all the hospitals and clinics in the United Kingdom, and the increasing demand in India is equally significant.

We give here a graph showing the growth of the COW & GATE sales during the last ten years. What is the reason for this extraordinary progress?

It is simply that the Medical and Nursing Professions all over the world have realised that COW & GATE Milk Food is the most satisfactory and reliable Infants' Food on the market to-day, and are recommending and prescribing it at every opportunity.

COW & GATE MILK FOOD

1923	1924	1925	1926	1927	1928	1929	1930	1931	1932

JOURNAL OF SOUTH INDIAN MEDICINE
JUNE, 1935

THE PROBLEM OF TUBERCULOSIS

India has many problems of public health to solve. From time to time some one or other of these engages the attention of the press and the public. Strenuous efforts are put forth towards the solution and a good beginning is made. In the meantime some other question comes to the fore, and the task well begun is half forgotten and gradually allowed to stagnate and get worse. Epidemics which set in with dramatic suddenness and result in fearful mortality arrest the attention and spur the Public and the Government on to take some measures of control. But in most cases these measures are temporary, and when the worst is over, things are again allowed to drift. But very little is heard of diseases like tuberculosis which set in insidiously and kill slowly. Medical practitioners are familiar with the increasing incidence of the disease. And they are very much distressed by their helplessness in the control of the condition.

A few decades ago there was some activity on the part of the medical and public health departments of this province over the control of tuberculosis. A few institutions were established where the disease could be carefully studied and the unfortunate patients efficiently advised. It was expected that such facilities would increase rapidly. At the end of twenty years one finds that the progress has been negligible. In the meantime the disease is spreading alarmingly and is carrying misery into many families. It is getting so common that medical men are not surprised to find it in all classes of the community. It is, therefore, more than ever necessary that concerted measures should be taken to check the spread of the disease and to minimise the death-roll caused by it. The magnitude of the task and the amount of money that is required to produce even fair results often frighten the Government into inactivity. The Government may be poor, and the economic condition of the country not favourable. But it is exactly under these conditions that the disease spreads wide and kills largely. Poverty of funds, therefore, instead of being used as an excuse for an attitude of fatal resignation, should stimulate the authorities to

devise means by which the work of amelioration could be carried out economically. Extensive sanatoria with imposing piles of buildings far away from thickly populated places are not certainly the most profitable way of spending limited resources. Nor private dwelling-houses improvised into wards and office-rooms the most suitable buildings for housing central Presidency Hospitals for dealing with tuberculous patients. Apart from sites and buildings, the staffing of the existing institutions is neither effective nor economical. The medical department in this country has long been used to deal with paid staff. Since the funds available have been limited, naturally the staff has been small and the sphere of activity limited. So long as its work in this limited field has been satisfactorily done, the department has been willing to let things alone. When any expansion of its activities was called for, it was always planned within its capacity to employ additional staff. This narrow outlook has been responsible to a large extent for many of the defects in medical relief in this country. If, instead of relying solely on the revenues of the country, the Government would earnestly seek the co-operation of the Public, and the Medical Profession particularly, we are sure a vast deal more could be done to expand rapidly medical relief in all its aspects in this Presidency. Those who may be reluctant to believe in the fruitfulness of this co-operation of the medical profession should study the reports on the administration of the medical department and the civil hospitals in the Presidency presented by Major-General Sprawson while he was in Madras.

THE VIZAGAPATAM MEDICAL COLLEGE

The Vizagapatam Medical College has been engaging the attention of the Government and others for some time. This College has not yet been recognized by the Indian Medical Council as efficient enough. We learn from the papers that there were a series of conferences over this subject. We are glad to find that the Government have agreed to find the necessary funds to improve the conditions in the college so that it might be soon given the hall-mark of recognition by the Indian Medical Council. But we must sound a note of caution in this connection. There is a common impression that the efficiency of an institution increases in proportion to the amount of money spent on it. But experience has generally been otherwise. Institutions where much of the available funds has been lavished on buildings and additional staff often show a poor out-turn compared to those whose buildings are modest and staff humble. Many of the missionary institutions in this presidency prove this fact: what tells is

efficient equipment and zeal on the part of the staff. We do not suggest that any teacher should be paid less than his due. But we object to unnecessary multiplication of jobs or ostentatious raising of so-called status of the teacher. The status of the teacher depends on his standing in the professional world and his contribution to the enrichment of the subject he deals with and on nothing else. We fully hope that the Government will bear this in mind when they actually plan for the expenditure of their money on the Vizagapatam Medical College.

Colonel E. W. C. BRADFIELD, O.B.E., C.I.E., I.M.S.

Surgeon-General with the Government of Bombay.

We are very glad to learn that Col. E. W. C. BRADFIELD, O.B.E., C.I.E., I.M.S., has been appointed Surgeon-General with the Government of Bombay. We, in Madras, feel highly gratified that an officer who has been with us, and whom we can with justifiable pride call ours, has been given this high appointment. Col. BRADFIELD came to Madras in 1912 and, except during the Great War, in which he served with distinction, he remained in the Province till he reverted to the Army early in 1931. He has been for several years a Surgeon in the General Hospital, Madras, and has held appointments as Professor of Anatomy, Professor of Biology, and Professor of Surgery in the Madras Medical College. During the last 5 years of his service here he was Superintendent of the General Hospital, in which capacity he distinguished himself as a capable administrator. He was mainly responsible for the many improvements which have been made in the General Hospital in recent years, and in recognition of his services (both as Surgeon and Superintendent) the new surgical block of the Hospital has been named THE BRADFIELD BLOCK after him. He has also acted as Surgeon-General with the Government of Madras. He goes to Bombay, as an experienced Medical Educationist and Administrator, and we are sure the destinies of the Medical Department of Bombay will find able guidance at his hands. We congratulate Col. BRADFIELD on his appointment, which he fully merits.

RETRENCHMENT IN HOSPITAL EXPENDITURE

We learn from the papers that the Government are appointing a special officer to investigate into the possibilities of economizing. As usual in retrenchment procedures, they are beginning at the wrong end. The money spent on diet and drugs is not much more than a fifth of the total

expenditure on civil hospitals and dispensaries. About half the expenditure goes towards the Medical Staff. To think of appointing a special officer to advise on the possibilities of saving a moiety of the fifth part while leaving alone the major portion would strike any one except the Medical Department and the Government as absurd. And yet, here is the proposal. We do not know anything about Col. Dyer who is expected to be the special officer. But if he comes with the idea that very much could be done in the way of retrenchment in drugs, diet and menial establishment, he is likely to be disappointed. It is more probable that a careful enquiry into the working of the hospitals would reveal the necessity for increase in the menial staff and more adequate supply of newer drugs and appliances. It is not uncommon to find hospitals which claim to be first class and model institutions condemned to carry on their work of treatment and investigation of diseases with rickety microscopes and inefficient instruments. The Government, if it means business, should look for real economy elsewhere. We wonder if the Government has taken the advice of the head of the department before venturing on their policy.

MEDICAL PERIODICALS

THE INDIAN MEDICAL GAZETTE

VOL. LXX. NO. 4. APRIL, 1935.

The Incidence of Pneumococcal Types in Pneumonia in Assam, L. Everard Napier and Dharmendra, *p. 181.*

"The sputa from 65 cases of pneumonia were investigated with reference to the species and type of streptococcus that predominated. Direct typing, mouse inoculation and cultural methods were used. Five strains were identified as hæmolytic streptococci and 60 strains as pneumococci. Of the 60 pneumococcal pneumonias 15 were broncho-pneumonia and 45 lobar. Of these 60 strains, 8 belonged to the 'fixed' types, 24 to other American types, and 29 were not identifiable with any sera we had in our possession; by the preparation of high-titre sera, 22 of the unidentified group were shown to belong to one of two types; these we have provisionally called A and B. Our type A was not associated with any mortality and appears not to be a virulent type. Type B was responsible for three deaths in 7 cases of lobar pneumonia whose fate was known; on the other hand 4 out of the 15 strains from cases of broncho-pneumonia and 3 out of 12 strains from normal throats were type B. It is evident that there is little 'fixed' type pneumonia in this district and therefore commercial therapeutic serum is not likely to be of much use in treatment. It is considered that type B is probably a strain with virulent potentialities that has a wide-spread distribution in the area in which we were working. If further experience shows that our type B pneumococcus is common in other districts in Assam and in other years, the question of making serum against or vaccines from this organism should be considered."

Preliminary Observations on a New Soluble Atebrin Compound, John R. Blaze and A. T. W. Simeons, *p. 185.*

"(1) The intramuscular injection of atebrin musonate in therapeutic doses does not give rise to toxic or unpleasant symptoms. (2) While a single injection of 0.375 gramme of atebrin musonate has in some cases a remarkable effect on the clinical picture, a recrudescence usually occurs within a few days. Heavy malignant tertian infections are only slightly influenced. (3) Two injections of 0.375 gramme given on successive days are sufficient to control within 48 hours the temperature, and in a maximum of 4 days all forms of benign tertian parasites, and the ring forms of malignant tertian. Crescents appear to be in no way influenced, and must be destroyed by plasmochin. In some cases there is a reappearance of parasites in the blood after a few days, but they disappear spontaneously. In no case hitherto observed has there been a return of fever after 2 injections. We are dealing with unusually heavy infections in a poor and mostly underfed class of patients. (4) The intravenous route though harmless is not satisfactory for routine treatment."

(5) The theoretical considerations already set out are supported by the result obtained. Atebrin musonate is a drug worthy of further investigation. (6) The immediate results of treatment with atebrin musonate appeared to be good enough to justify our making this preliminary report. But the period of observation has not been long enough to enable us to compare the relapse rate after atebrin musonate with that after oral atebrin or quinine. Observations upon this important question are being made."

Habitual use of Barbituric Acid Derivatives in India, R. N. Chopra and G. S. Chopra. p. 188.

"(1) The use of barbituric acid derivatives has greatly increased in India during the last few years. These compounds have powerful hypnotic and pain-relieving properties being only second to opiates in this respect. (2) The fact that this group may produce toxic effects even in therapeutic doses and are cumulative in their action has not been sufficiently appreciated by the medical profession. (3) Their repeated use is harmful and dangerous, definite pathological changes being produced in vital organs. No tolerance is developed and any increase of dose may give rise to dangerous symptoms and even fatal results. The margin between toxic and therapeutic doses is very narrow and certain individuals are particularly susceptible to the action of these drugs. Caution is therefore recommended in their use. (4) Habit formation has occurred with these drugs in Indian subjects but the desire for the repetition of the dose is more for relief of symptoms than anything else. They do not appear to produce euphoria like opiates, hemp drugs and alcohol. Abstinence symptoms are not produced and habit can be easily broken. (5) The need for bringing this group of drugs under control, similar to that exercised in Great Britain, is emphasized."

Bernhardt's Syndrome, E. Muir and S. N. Chatterji, p. 192.

The Reticulo-Endothelial System in Malarial Haemoglobinuria of Monkeys. Part I. K. V. Krishnan, p. 193.

"1. The reticulo-endothelial system probably plays an important part in the aetiology and prevention of malarial haemoglobinuria. 2. When the system is depressed, as is evidenced by a low reticulo-endothelial cell count and inability to take up large quantities of vital dyes, sudden severe haemolysis results in haemoglobinuria. 3. When the system is active and functioning properly, as is evidenced by a high reticulo-endothelial cell count and ability to engulf large amounts of vital dyes, sudden severe haemolysis does not result in haemoglobinuria."

The Reticulo-Endothelial System in Malarial Haemoglobinuria of Monkeys. Part II. K. V. Krishnan, and B. M. Ghosh, p. 197.

(1) Removal of the spleen in monkeys infected with *P. knowlesi* intensifies their infection and increases the incidence of haemoglobinuria. (2) Treatment of monkeys showing haemoglobinuria with quinine is unsatisfactory. It is even less effective in splenectomized animals. (3) The results recorded in this paper stress

the importance of susceptibility in haemoglobinuria and suggest that the spleen and the reticulo-endothelial system play an important part in this susceptibility.

Preliminary Note on the Cultivation of Vaccinia Virus on the Chorioallantoic Membrane of Chick Embryo, K. S. Shah, p. 200.

Hygienic Method of Composting Refuse with Night-Soil, J. J. Rao, p. 202.

The Value of Phrenic Exairexis in the Treatment of Pulmonary Tuberculosis, P. V. Benjamin, p. 207.

"The paper reviews 275 cases of phrenicectomy done at the Union Mission Tuberculosis Sanatorium, Arogyavaram, near Madanapalle."

"The analysis of the results of treatment with phrenicectomy show that the operation is of considerable value, but mostly so when it is used to aid the collapse and healing in partial artificial pneumothorax. While its effect when used alone in cases with fever is not marked, it does cause a considerable increase in the percentage of cases losing bacilli from the sputum."

JOURNAL OF THE INDIAN MEDICAL ASSOCIATION

VOL. IV. NO. 8. APRIL, 1935.

On The Possible Role of Inanition in the Aetiology of Beri-Beri Etc. D. D. Chatterjee, p. 309.

(1) Animals fed on a diet deficient in Vitamin B starve in the midst of plenty. (2) The structural and functional changes which take place in the intestines of fasted animals and of animals presented with a diet deficient in Vitamin B are similar. (3) It is concluded that the symptoms which follow on a deficiency of Vitamin B are due to two processes: (a) A starvation process — this is a constant process and is possibly due to a duodenitis depriving the animal of hunger sensation. There may also be changes in the capacity to smell. (b) An infective process — this is probably variable, and may or may not be added to the former according to the environment.

The Role of Malaria in the Causation of Cirrhosis of the Liver—A Preliminary Note. T. S. Tirumurti and M. V. Radhakrishna Rao, p. 315.

The common association of malaria and cirrhosis of the liver in India has given rise to divergence of opinion regarding the role of the former in the causation of the latter. Brief reference is made to the views of some of the workers in India and the importance of the problem is pointed out. Judging from personal experience it is concluded that malaria per se is not a direct cause of cirrhosis of the liver.

Preliminary Report on a New (?) Strain Isolated From C. S. F. of Epidemic Meningitis Cases 1934. Benode Behari Sen, J. N. Naha, S. C. Bhattacharyya and S. C. Acharjya, p. 318.

Puerperal Sepsis, C. L. Houlton, p. 321.

The Reticulocyte Count in Malaria and Kala-Azar and its Significance. T. Bhaskara Menon, T. K. Krishnaswamy, and D. R. Annamalai. *p.* 359.

- (1) Studies on untreated cases of malaria have shown that the reticulocyte count in such cases is slightly higher than the normal. The count is generally higher in chronic relapsing malaria than in acute attacks. (2) No definite relation could be made out in untreated cases between the reticulocyte count and the count of the red blood cells. (3) Some evidence is brought forward to show that cases of untreated kala-azar give on an average, higher counts than acute malaria. (4) The reticulocyte behaviour in malaria and kala-azar is regarded as due to the action of two differing mechanisms: in the former a haemolytic anaemia is followed by a reaction of active haemopoietic tissue, while, in the latter, there seems to be an irritation of the displaced and compressed erythrocytic tissue by parasitic invasion of the reticulo-endothelium.

Studies on the Biochemistry of Cerebrospinal Fluids of Epidemic Meningitis in India (1933-34). S. Hazra. *p.* 364.

The Operative treatment of Vesico-Vaginal Fistula. W. C. Paton. *p.* 368.

Insuring Against Sickness. Benoy Kumar Sarkar. *p.* 371.

Ready for Sale!

Ready for Sale!!

Only Limited Copies are Available

MADRAS MEDICAL DIRECTORY

(REGD.)

Revised and enlarged second edition including over 5000 addresses of Hospitals, Chemists and opticians with Medical Practitioners of Madras Presidency.

Over 300 Pages with 26 illustrations neatly finished.

Price Rs. 2-8-0 Postage extra.

(Readers of this Journal Postage Free)


Apply to:

MATHEWS & CO.,

16, General Muthia Mudaly St.,

SOWCARPET :: MADRAS

Sirop "THIO-CALCIN"



A Super calcium Tonic for Expectant and Nursing mothers. An effective Lung tonic for Tuberculosis, Troublesome Coughs and low fevers. Useful in Rickety and weak children.

An ounce Represents 20 grains Pot sulpho Guaiacol, 30 grain assimilable calcium, Glycerine Orange Juices, P. S. Dose—age one to 2 Teaspoonful 3 or 4 times a day.

Sold at all Chemists:

6 oz., 16 oz., and one gallon Bottles.

Madras Agents:

DADHA & Co., MADRAS.

For Literature and Samples Apply to:

THIO-CALCIN Co.,

16, VENKATESAPURAM, MYLAPORE, MADRAS.

HEMOGONON

(SYRUP)

"MADE IN FRANCE"

WHAT IS IT?

Hemoglobin, Fresh Liver Extract, and Iron Glycerophosphates, scientifically blended into a palatable syrup.

WHEN SHOULD IT BE PRESCRIBED?

In Anaemia (Primary & Secondary) General Debility, Convalescence, during Pregnancy and Lactation, etc.

WHY SHOULD IT BE PREFERRED?

It increases the Erythrocyte count and Hemoglobin values promptly and lastingly. It causes no Diarrhoea or Constipation. Patients relish it for its very palatable taste, and

ITS PRICE IS REASONABLE

Sold in glass bottles of 10½ fl. ozs.

THE Research Laboratory,

Harris Road,
MADRAS.

Microscopical, cultural and bacteriological examinations of Blood, Sputum, Urine, Faeces, etc., done.

Widal, Wassermann and Kahn tests and Complement Fixation test for Tuberculosis, etc., done.

Stock and Autogenous Vaccines prepared.

Sterile Solutions for injection always in stock.

Please apply for further particulars to:

MANAGER.

For difficult Infant Feeding Cases

The difficulties and trouble of feeding infants and others suffering from gastric weakness are now overcome by this highly scientific preparation. Peptalac is a dry stable powder. The simple addition of hot water, fruit, cream, milk, and predigested wheat is available to infants, but containing as it does a proportion of predigested wheat flour, it will serve as an ideal food to weak infants taking more solid foods at 8-9 months.

PEPTALAC

This is a perfect milk in powder form, made in three strengths, namely: Lactac Separated (10 Fat), Lactac Half Cream (27 Fat), and Lactac Full Cream (37 Fat). It is proving of great value in the feeding of infants recovering from gastroenteritis, broncho-pneumonia, and other diseases, and eliminates the difficulties and disadvantages of administering liquid milk treated with lactase and as proposed by Marriott.

LACIDAC

This is a Full Cream milk to which has been added Iron, Ascorbic Acid, Calcium B.P., and Vitamin B. It was the food recommended by the Food Hygiene Commission in the 1940s, and was used by the Ministry of Health on the 10th of April 1940. It was also recommended by the Medical Research Council in their Report, Special Series No. 157, in 1941. It is a perfect milk for infants and for premature babies.

HÈMOLAC

This has been prepared at the request of a leading medical authority for the treatment of infants with jaundice. It is characterised by its high iron content and is so constituted that it is easily absorbed and assimilated by the infant's system. It is a perfect milk for infants and for premature babies.

ALLERGILAC

Brestol is an excellent food for infants and others suffering from gastric weakness. It is a dry stable powder. The simple addition of hot water, fruit, cream, milk, and predigested wheat is available to infants, but containing as it does a proportion of predigested wheat flour, it will serve as an ideal food to weak infants taking more solid foods at 8-9 months.

BRESTOL

This was prepared originally for the use of infants recovering from gastroenteritis, broncho-pneumonia, and other diseases, and eliminates the difficulties and disadvantages of administering liquid milk treated with lactase and as proposed by Marriott.

FRAILAC

These Foods are indicated for all cases of intolerance of milk, and are particularly useful in the treatment of infants with jaundice, and in the treatment of infants with gastroenteritis, broncho-pneumonia, and other diseases.

SEPARATED & HALF CREAM

Pro-lac is a dry stable powder. The simple addition of hot water, fruit, cream, milk, and predigested wheat is available to infants, but containing as it does a proportion of predigested wheat flour, it will serve as an ideal food to weak infants taking more solid foods at 8-9 months.

PROLAC

COW & GATE LTD.
GUILDFORD, SURREY

ASSOCIATION NOTES

THE TRICHINOPOLY DISTRICT MEDICAL ASSOCIATION (REGISTERED)

A monthly meeting of the above association was held on Saturday the 18th May '35 at 5 P. M. in the E. R. High School, Trichinopoly, when about 30 members were present. Dr. C. Somasundaram of Puvaloor was 'At Home' and Dr. R. Kalamegham, M.B.B.S., presided.

Dr. T. S. S. Rajan showed a small boy aged 10 whom he had operated upon for Hydronephrosis. The removed matter resembled a big bag with the kidney substance embedded in its wall. He demonstrated the old case of Stricture of Oesophagus shown in the previous meeting and which showed some improvement. He also demonstrated a case of Multiple Arthritis in a girl of 10 sent over for X-ray and demonstration by Dr. V. Iravatham of Thirukkattappally.

A case of Electric Burns treated with cod-liver-oil ointment in the Government Head Quarters Hospital, Trichy, was also demonstrated.

After demonstration there was discussion in which many members took part.

Dr. K. S. Narayanan of Madura then delivered an interesting lecture on 'Peroral Endoscopy' explaining the technique, indications, and its usefulness for diagnosis and treatment. He also demonstrated a number of latest instruments and explained their use.

With a vote of thanks proposed by the Secretary to the Lecturer, Chairman of the evening, the Host, and the Manager of the premises, the meeting terminated.

THE TINNEVELLY DISTRICT MEDICAL ASSOCIATION, PALAMCOTTAH.

The Tinnevelly District Medical Association had its monthly meeting at Tiruchendur, one of the most sacred and salubrious seaside resorts of the Presidency, on Sunday the 26th May 1935. Members numbering 38 including some lady doctors gathered from all parts of the District. Some of the local officials and visitors were also present. Refreshments were served to all

present at 5 P. M. One of the local residents gave sweet music on the harmonium.

The meeting began at 6-30 P.M. under the Presidentship of Lieut. Col. T. S. Shastry, I.M.S. The Secretary read the minutes of the last meeting which were passed.

The following resolution was moved from the chair which was carried unanimously:—

The last Sunday of every month will be the meeting day of the Association hereafter in view of the Surgeon-General's suggestion to hold it on Government holidays.

The President moved that the following be made Honorary members of the Association and they were elected unanimously:—

1. Lieut. Col. T. W. Harley, I.M.S., (Retired).
2. Dr. B. Rama Baliga, B.A., M.B. & C.M., (Retired D. M. O.)
3. Dr. D. Krishnayyah, B.A., M.B. & C.M., (Now D. M. O. Chittoor).

A small booklet entitled "Notes on Cholera and its Treatment" compiled by Lieut. Col. T. S. Shastry, I.M.S., was distributed to every one of the members. Lieut. Col. T. S. Shastry, I.M.S., explained the use of the notes for every medical practitioner as a pocket guide for the treatment of Cholera.

Dr. S. Kolappa Pillai, L.M. & S., B.S.Sc., District Health Officer, Tinnevely, presented certain posters on Malaria, Tuberculosis, etc., to the Association for which the President thanked him.

Dr. N. Isaac, L.M.P., read interesting clinical notes on two cases of Caesarean Section performed by Lieut. Col. T. S. Shastry, I.M.S. at the Government Women and Children Hospital, Vannarpet. Then Lieut. Col. T. S. Shastry, I.M.S., delivered an interesting lecture on Caesarean Section.

Dr. K. S. Narayanan, L.C.P.S., (Bombay), L.D.Sc. (Cal.) Z.I.O. (Vienna), gave an interesting demonstration and lecture on Haslinger's Endoscope which was very much appreciated. The President offered cordial thanks to Dr. Srinivasa Prabhu, L.M.P., and his co-workers for their kind and generous hospitality and for organising the meeting.

Then a very enjoyable sea-bath and a grand dinner brought the modest function to a close late at night.

ANDHRA MEDICAL ASSOCIATION

Annual Session

The annual session of the Andhra Medical Association for the year 1934-35 was held at the Town Hall, Vizagapatam, along with the conference which met on the 7th, 8th and 9th June, under the presidency of Dr. A. Lakshmipathi, B.A., M.B. & C.M., of Madras.

Among the amendments made in the draft rules of the Association was one to the effect that 10 per cent of the collections of the branches by subscription for the membership of each branch shall be paid annually to the Central Council.

The Association passed a number of resolutions the following being some of them:—

(1) The Association resented the Government attitude requiring a second opinion from Government Medical Officers on certificates issued by private practitioners, as this procedure is against the dignity of the independent medical profession. (2) It resolved that doctors in Government service should not be allowed private practice unless it be of a consulting nature, and (3) protested against the non-recognition of the Andhra Medical degrees by the Indian Medical Council and appointed a committee to investigate and agitate about the non-recognition of the Andhra Medical degrees. (4) It thanked Sir S. Radhakrishnan and Major Edden for their sincere efforts to improve the Vizagapatam Medical College and Hospital. (5) It recommended the use of Khaddar and Indian made articles and medicines as far as possible by its members. It passed resolutions, (6) urging that appointments in the Vizagapatam Medical College should be given to Andhras as far as possible, (7) urging institution of M. D., and M. S. degrees in the Andhra University, (8) recommending the starting of a central pharmaceutical laboratory in Andhradesa by the Association either by private efforts or through the agency of the Government, (9) urging the need for paying a minimum of at least Rs. 600 per year to rural medical practitioners, and (10) requesting the Government to ameliorate the services of medical men in the Local Boards and Municipalities by allowing them the same scale of pay that is granted to men in Government service.

Office-bearers for the coming year were then elected with Dr. D. S. Ramachandra Rao as President, Dr. S. V. Ramarao as Vice-president and Drs. M. V. Krishna Rao and C. Suryanarayanamurti as Secretaries, the latter being also the Treasurer.

An editorial advisory board consisting of Dr. Pattabhi Seetharamayya, Dr. D. S. Ramachandra Rao and others, was constituted to bring out a journal.

Dr. Lakshmipathi, in his presidential address, remarked that he was pleased to find that the bridge between Allopathic and Ayurvedic practitioners in the Andhra country was quickly closing up, that the untouchability of the Ayurvedic physician was removed by the Government opening the School of Indian Medicine under its own guidance and that the progress of medicine in the country depended upon the young and enthusiastic graduates, who should study the subjects from all points of view. He added that, though he was a strong supporter of Ayurvedic system, he was not against the introduction of the western scientific methods.

Continuing, he said that he was struck by the remarkable spirit of co-operation and friendliness shown by the students, the graduates and the Professors of the Vizagapatam Medical College in conducting the session. He exhorted the Andhra medical practitioners to organise themselves strongly by the establishment of branches in all the District and Taluk headquarters and by helping the progress of independent medical practitioners.

EYE, EAR, NOSE, & THROAT HOSPITAL FOR CHETTINAD. *New Buildings Opened.*

Before a crowded audience of Officials and non-Officials besides several influential and prominent Nagarathars of the District, Mr. M. P. Pai, M. A., I. C. S., Sub-Collector, Devakottai, declared open last evening the newly constructed extensions for the Eye, Ear, Nose & Throat Hospital of Dr. Shetty at Chettinad in Ramnad District.

The function was held at the decorated pandal erected in front of the Hospital premises. Despite inclement weather, there was a large gathering and prominent among those present were Messrs. the Kumara Raja of Chettinad and Mayor of Madras, Rao Bahadur K. V. AL. RM. Ramanathan Chettiar, Rao Bahadur O. A. PR. M. AR. Arunachalam Chettiar C. V. C.T. V. Venkatachalam Chettiar and the Commissioner of Karaikudi and several members of the Medical profession.

Dr. Shetty, who is specially qualified in Vienna for Eye, Ear, Nose & Throat treatment, welcomed the Collector and the audience and made a short speech, tracing the successful working of the Hospital for the past



A man aged 45, suffering from very severe pains in the knee joints, came to us one day with high fever and severe pains in the body. The peculiarity of the case was that there was no pain in the elbow joints, though it was unbearable in the knee joints.

For two days the patient was kept on Sodi Salicylas mixture which made him feel a little better. After a few days again he began to complain of pain in the joints. He was, therefore, given intravenous injections of Sodi-Iodide and Sodi-Salicylas (A.F.D.) for three successive days. The patient began to show a marked change as regards his temperature chart as well as his gait. The fourth injection was not administered because he began to perspire after the third injection.

The patient was kept on a salt free diet. After three or four days the patient once more began to complain of pain in the knee joints. Closer enquiry revealed that the patient was formerly suffering from Gonorrhœa, which confirmed that his present arthritis was of the gonorrheal type. He was given Rheantine Pills (A. F. D.) 2 pills, four times a day to attack the traces of gonorrheal infection left. On the fourth day he felt much better than when taking only Sodi-Iodide and Sodi-Salicylas mixture. So it was decided that the combined treatment of both this mixture and Rheantine pills should be continued. Two further injections of Sodi-Salicylas and Sodi-Iodide were also administered.

Within a fortnight the patient felt much better. He is now able to walk about with much ease. To combat his anaemic condition he has been put on Fer-Ma-Pho (A. F. D.) tonic. The patient is still taking the aforesaid mixture.

From the Diary of a Medical Practitioner.

ADVT.

CONOSAN

the mild and effective balsamic for

CONORRHOEA

Dosage: 2 capsules 2 to 3 times a day during or after meals.

Supplied:
in glasses of 40 capsules of 4½ grains each.



NOCTAL

the hypnotic in all cases of

INSOMNIA

Dosage: 2 tablets half an hour before retiring preferably in some warm liquid.

Supplied:
in tubes of 10 tablets of 1½ grains each.

Agent: T. RAMACHANDRA RAO, 153, Broadway, MADRAS.

JOURNAL OF SOUTH INDIAN MEDICINE

NOTICE

CONTRIBUTIONS of medical interest—original articles, case notes, clinical memoranda—are invited. They should be addressed to the EDITOR, JOURNAL OF SOUTH INDIAN MEDICINE, EGMORE, MADRAS, and should be accompanied by a stamped addressed envelope if their return is desired in case of non-publication.

Articles are accepted for publication on condition that they are contributed solely to this journal.

Matter appearing in the JOURNAL OF SOUTH INDIAN MEDICINE is covered by copyright. Reproduction elsewhere without permission is not allowed.

MANUSCRIPTS must be written clearly or preferably type-written, with double spacing, on one side of the paper only.

Books and pamphlets for review should be sent to the Editor, who will be pleased to receive also reprints of which abstracts will be of interest to the profession. Journals for exchange or review also should be addressed to the Editor.

two years. He said that the institution has been fully equipped with the most up-to-date Electro Therapeutic Apparatus, such as Diathermy, Pantostat, Ultraviolet etc., besides apparatus for specialist department of Eye, Ear, Nose and Throat and for general surgery. The Hospital, he added, had provision for about 20 beds with modern conveniences. Dr. Shetty, the founder of the Hospital, was glad to announce that it had been always the principle of the institution not to refuse any case when he felt he could do some good, however poor the patient might be, and assured them all that the Hospital would cater to the needs of the poor patients free of all charges in the special department if such deserving patients would bring a note from their Doctors.

The new buildings of the Hospital had been constructed in up-to-date fashion under the skilled supervision of Mr. J. F. Pulford, Chartered Architect of South India. Messages wishing the institution every success received from the Rajah of Chettinad, Dr. U. Rama Rao, Dr. T. S. S. Rajan, M. L. A., and M. S. M. M. Meyyappa Chettiar, Municipal Chairman, Karaikudi, and several others were read.

In declaring open the new buildings, Mr. Pai congratulated Mr. Shetty on his private enterprise in successfully working the institution for the past two years. He was glad to note that 40 poor out-patients and 4 in-patients were daily being treated free and that the people were increasingly realising the usefulness and importance of this private Hospital for special treatment. The Institution, the speaker observed, stood for the very high ideal of its profession during its past, viz., giving free treatment to the poor, and any institution with such ideals was bound to succeed. He wished the institution on a bright future.


With a vote of thanks by Dr. Shetty, the function terminated. This was followed by a well attended dinner arranged by Dr. Shetty at the "Chettinad Aerodrome Grounds."

XXXVI

JOURNAL OF SOUTH INDIAN MEDICINE
MEDICAL EXAMINATION RESULTS
UNIVERSITY OF MADRAS.

The following candidates were successful in the Final examination for the Degree of M. B. B. S., held in April 1935:

Audikesavalu, M., Bhimaneshbushan Chattopadhyay, Muktha Bai, B., Bhupendranath Mukhopadhyay, Chakrapani Menon, M., Saraswathi, M. A., Dwaraka Bai, G. S., Krishnamurti, V., Dalal, I. C., Venkatesvara Rao, T., Venkatalakshamma, K., Padmarathi, K. K., Kokila Ramabrahma Paranjoti, Prakash, D., Gopala Pillai, P. P., Balakrishnan, V. K., Cheriyar Mary P., Daniel, P. H., George, P. V., Grant, May E., Jagannathan, N., Janardhanan Poti, A., Koman, Pramella, Krishnamurti Rao, B., Kurien P. V., Lakshmikantham, M., Martinus, G. F., Marhai, P. M., Mather, Achy., Nagaswami, S., Narayanan, G., Rajaram, S. R., Shankara, Advanthava, N., Sitarama Rao, Thomas, P. G., Valisvaran, S., Vasudeva Rao, G., Venkata Jagannatha Rao, P., Venkatasubrahmanyam, C. S., Yegnanarayanan, M. S.



Please fill the Order Card below.
No postage necessary, we pay postage on delivery!

GONEAL
MADE BY
(I.O.N.)
(ITALY)

VENEREAL SPECIALIST
WRITES ON
'GONEAL'

* "We tried a new Preparation called "GONEAL" (A product of the National Opoterapic Institute, Italy, samples of which were supplied by Messrs. GOPAL & CO., Madras) on several cases of GONORRHEA. The preparation is a Colloidal Solution of Silver and Calcium Caseinate put up in 2 c.c. ampoules. The drug is given intramuscularly on alternate days. The REACTION both focal and local is MINIMAL. The therapeutic response in COMPLICATIONS of GONORRHEA was satisfactory.

(*Extract from "ANTISEPTIC" May Issue, 1934, Page 310.)

Usually ONE BOX

is MORE THAN SUFFICIENT for one course.

Rs. 9 per Box of 5 amps. of 2 c.c. for Intramuscular use.

An introductory offer:—

Postage and packing charges FREE for orders from Doctors only.
SPECIAL INDIAN PACKING.

TELEGRAMS:

"GOPAC" MADRAS.

Sole Agents for India & Ceylon:—


GOPAL & CO.

No. 10, T. P. Koil Street, Triplicane, Madras.

GONEAL
MADE BY
(I.O.N.)
(ITALY)

T. V. Lander and ...	th (G. 3)	...	83
Christie, A. C.—Radiation in the Treatment of Cancer (G. 24)	...	...	253
Chronic Amoebiasis (G. 7)	...	...	121
Chronic Appendicitis and the Right Lower Quadrant Syndrome, Charles W. Lester (G. 6)	...	...	120
Clinical Memoranda, Treatment of Small-pox, V. Govindan Nair	...	...	257
Collier, James, Epilepsy and Its Treatment (G. 21)	...	...	248
CONNOR, SIR F. P.	...	...	1
Foreword	...	...	6

An Important Therapeutic Conquest.



"ERGAL"

(Injectable Extract of
SUPRARENAL MEDULLA)

This Preparation which
MUST NOT BE CONFUSED
with the ordinary solution of
ORGANIC ADRENALIN has
shown itself capable of evolving
in the organism in the mildest but
more lasting and complete form

than adrenalin, the multiple harmonic and pharmacodynamic
actions of the medullary portion of the suprarenal capsule.

Highly potent, "ERGAL" can be used with greater benefit,
in conditions where the commercial adrenalin is indicated.

The "Ergal with Atropine" finds its precise indication in
treatment of Bronchial Asthma.

Issued plain in combination with Atropine and with
Strychnine separately.

Rs. 6-4-0 a Box of 6 amps.

INDEX

Note: (O)—Original Articles; (G)—Gleanings from Medical Literature.

Acute Infections of the Upper Respiratory Tract, Middle-Ear and Mastoid, J. R. Page and H. C. Ballenger (G. 23) ...	251
Alcohol and Automobile Accidents (G. 8) ...	123
ALEXANDER, R. D. A Note on a Case of Tropical Typhus occurring in Madras (O) ...	89
All-India Ophthalmological Society, Fourth Conference, The ...	195
Arnheim, E. E. and Newhof, H. A Lowered Morality in Acute Appendicitis (G. 5) ...	119
Artificial Pneumothorax in the Treatment of Pneumonia, Lewis J. Moorman, (G. 18) ...	213
BACILLUS COLI INFECTIONS (O) ...	41
T. Satakopan ...	
BALASUBRAMANIA IYER, T. S. The Problem of Birth Control (O) ...	152
BHASKARA MENON, T. A Search for New Types of Tropical Splenomegaly (O) ...	227
Bradfield, Col. E. W. C., O.B.E., C.I.E., I.M.S., Editorial ...	261
Budget Speeches, Editorial, ...	167
Building Craze, The, Editorial ...	86
Carcinoma of the Male Breast, Gilbert Judson (G. 2) ...	82
Case of Evipan Paralysis, A T. V. Lander and Mahomed Sallah (G. 3) ...	83
Christie, A. C.—Radiation in the Treatment of Cancer (G. 24) ...	253
Chronic Amoebiasis (G. 7) ...	121
Chronic Appendicitis and the Right Lower Quadrant Syndrome, Charles W. Lester (G. 6) ...	120
Clinical Memoranda, Treatment of Small-pox, V. Govindan Nair ...	257
Collier, James, Epilepsy and Its Treatment (G. 21) ...	248
CONNOR, SIR F. P. Foreword ...	1

Control of Cancer, <i>Editorial</i> ...	218
DEVELOPMENT OF THE HEART, NORMAL AND ABNORMAL (O), P. K. Koshy ...	236
DIAGNOSIS OF KALA-AZAR, THE (O) K. C. Paul and V. Govindarajulu ...	129
Diagnosis and Treatment of Intestinal Obstruction, David H. Patey (G. 20) ...	215
Douglas, Symmers Status Lymphaticus (G. 2) ...	81
DURAI SWAMI, T. S. Headache: Its Differentiation and Significance (O) ...	104
EDITORIALS	
Bradfield, E. W. C., Col. I.M.S., Budget Speeches ...	261
Building Craze, The ...	167
Control of Cancer ...	86
Election to the Indian Medical Council ...	218
Krishna Menon, The Late Dr. T. Problem of Tuberculosis, The ...	217
Problems on Filariasis ...	37
Research and Record ...	259
Retrenchment in Hospital Expenditure ...	127
Retrospect ...	125
Rural Medical Relief ...	261
Rural Medical Scheme ...	35
Vizagapatam Medical College, The ...	85
Elmslie, R. E. (and others) — Fibrocystic Disease of Bone (G. 11) ...	219
Endocrine Effects of Certain Ovarian Tumours, Emil Novak (G. 25) ...	260
Epidural and Perineural Injections in the Treatment of Peripheral Pain, E. A. Joseph (G. 14) ...	160
Epilepsy and its Treatment, James Collier (G. 21) ...	254
Faxon, H. N.—Treatment of Varicosities, (G. 19) ...	162
Fibrocystic Disease of Bone, R. E. Elmslie, R. Davis-Colley, H. Taylor (G. 11) ...	248
FOREWORD ...	215
Sir F. P. Connor ...	160
GASTRIC NEUROSES (O) ...	1
T. Satakopan ...	97

GLEANINGS from Medical Literature—

Acute Infections of the Upper Respiratory Tract, Middle-Ear and Mastoid (23) ...	251
Alcohol and Automobile Accidents (8) ...	123
Artificial Pneumothorax in the Treatment of Pneumonia (18) ...	213
Carcinoma of the Male Breast (2) ...	82
Case of Evipan Paralysis (3) ...	83
Chronic Amoebiasis (7) ...	121
Chronic Appendicitis and the Right Lower Quadrant Syndrome (6) ...	120
Diagnosis and Treatment of Intestinal Obstruction (20) ...	215
Endocrine effects of Certain Ovarian Tumours (25) ...	254
Epidural and Perineural Injections in the Treatment of Peripheral Pain (14) ...	162
Epilepsy and its Treatment (21) ...	248
Fibrocystic Disease of Bone (11) ...	160
Lowered Mortality in Acute Appendicitis, A (5) ...	119
Management of Urologic Complications in Injuries to the Spine, The (13) ...	161
Nervous Breakdown (22) ...	249
Oral Urography (15) ...	163
Pathology and Treatment of Lymphadenoma (12) ...	161
Radiation in the Treatment of Cancer (14) ...	253
Recurrent Dislocation of the Shoulder (16) ...	163
Spinal Anaesthesia: Nervous System Sequelae (4) ...	118
Status Lymphaticus (1) ...	81
Strychnine and the Barbiturates: Mutual Antidotes (17) ...	213
Studies on Reticulocytes (9) ...	123
Treatment of Angina Pectoris, The (26) ...	255
Treatment of Fracture of Spinal Column (10) ...	124
Treatment of Varicosities (19) ...	215
Gordon, M. H. and Levitt, W. M. Pathology and Treatment of Lymphadenoma (G. 12) ...	161
GOVINDAN NAIR, V. Treatment of Small-pox (Clinical Memoranda) ...	257
GOVINDA RAU, P. Ligature of Femoral Vein in Diabetic and Senile Gangrene (O) ...	179
HEADACHE: ITS DIFFERENTIATION AND SIGNIFICANCE, (O) T. S. Duraiswami ...	104

270	JOURNAL OF SOUTH INDIAN MEDICINE	
Hyslop, George H., Spinal Anaesthesia; Nervous System Sequelae (G. 4)	118	
<i>Indian Journal of Venereal Diseases</i> , REVIEW	165	
Jimeno-Vidal, F.		
Treatment of Fracture of Spinal Column (G. 10) ...	124	
Joseph, E. A.		
Epidural and Perineural Injections in the Treatment of Peripheral Pain (G. 14)	162	
Judson, Gilbert.		
Carcinoma of the Male Breast (G. 2)	82	
KOSHY, P. K.		
Development of the Heart, Normal and Abnormal (O)	236	
T. Krishna Menon		
Obituary	39	
KRISHNA MENON, T.		
Some Common Drugs of Cardiac Practice (O)	17	
The Late Dr. T. Krishna Menon, <i>Editorial</i>	37	
Lander, T. V. and Sallah, Mahomed,		
A Case of Evipan Paralysis (G. 3)	83	
Lester, Charles, W.		
Chronic Appendicitis and the Right Lower Quadrant Syndrome (G. 6)	120	
LEUKAEMIAS, THE (O), S. Thambiah	145	
LIGATURE OF THE FEMORAL VEIN IN DIABETIC AND SENILE GANGRENE (O), P. Govinda Rau	179	
Lowered Mortality in Acute Appendicitis, A E. E. Arnheim and H. Newhof (G. 5)	119	
McROBERT, G. R.		
A Note on Tropical Typhus Fever (O)	93	
Madras Medical Council, News and Notes	88	
<i>Madras Medical Directory, The</i> , REVIEW	258	
MAHADEVAN, V.		
Surgery of the Breast (O)	3	
Mallayya, B. S.		
Obituary	88	
Management of Urologic Complications in Injuries to the Spine, The T. F. Connors and I. E. Nash (G. 13)	161	

INDEX	271
<i>Materia Medica of Pharmaceutical Combinations and Specialities</i>	258
REVIEW	88
Medical Miscellany, A, News and Notes	169, 221, 263
MEDICAL PERIODICALS	
Moorman, Lewis J.,	213
Artificial Pneumothorax in the Treatment of Pneumonia (G. 18)	
NEED FOR A NATIONAL ANTI-CANCER CAMPAIGN	183
IN INDIA, THE (O), P. Rama Rau	249
Nervous Breakdown, C. P. Symonds (G. 22)	
Nicola, Toufic.	163
Recurrent Dislocation of the Shoulder (G. 16)	
NOTE ON A CASE OF POPLITEAL ANEURYSM TREATED BY HUNTER'S METHOD (O), B. M. Sundaravadanan	233
NOTE ON A CASE OF TROPICAL TYPHUS OCCURRING IN MADRAS, A, (O), R. D. Alexander	89
NOTE ON TROPICAL TYPHUS FEVER, A, (O) G. R. McRobert	93
Novak, Emil	
Endocrine Effects of Certain Ovarian Tumours (G. 25)	254
Oral Urography	163
Harold Swanberg (G. 15)	
Page, J. R. and Ballenger, H. C.	
Acute Infections of the Upper Respiratory Tract, Middle-Ear and Mastoid (G. 20)	251
Parasuram, A. R.	88
Obituary	
Pathology and Treatment of Lymphadenoma M. H. Gordon and W. M. Levitt (G. 12)	161
Patey, David H., Diagnosis and Treatment of Intestinal Obstruction (G. 20)	215
PAUL, K. C. and GOVINDARAJULU, V.	129
The Diagnosis of Kala-azar (O)	
PROBLEM OF BIRTH CONTROL, THE, (O) T. S. Balasubramania Iyer,	152
Problem of Tuberculosis, The, <i>Editorial</i> Problems on Filariasis, <i>Editorial</i>	172

272	JOURNAL OF SOUTH INDIAN MEDICINE	
	<i>Proceedings of the All-India Ophthalmological Society, Vol. III</i>	
	REVIEW	
	Radiation in the Treatment of Cancer, ...	165
	A. C. Christie (G. 24) ...	253
	RADIUM THERAPY, (O), ...	68
	P. Rama Rau ...	200
	Ramachandra Rao, D. S. ...	58
	Presidential Address (Rural Practitioners Conference) ...	68
	RAMAN PILLAI, G. ...	183
	Streptococcal Disease (O) ...	163
	RAMA RAU, P. ...	125
	Radium Therapy (O) ...	35
	RAMA RAU, P. ...	165
	The Need for a National Anti-Cancer Campaign in India (O) ...	258
	Recurrent Dislocation of the Shoulder ...	165
	Toufic Nicola (G. 16) ...	209
	Research and Record, Editorial ...	85
	Retrospect, Editorial ...	219
	REVIEWS—	
	<i>Indian Journal of Venereal Diseases</i> ...	97
	<i>Madras Medical Directory, The</i> ...	227
	<i>Materia Medica of Pharmaceutical Combinations and Specialties</i> ...	255
	<i>Proceedings of the All-India Ophthalmological Society, Vol. III</i> ...	17
	Rural Medical Practitioners Fifth Provincial Conference ...	23
	Rural Medical Relief, Editorial ...	118
	Rural Medical Scheme, Editorial ...	
	SATAKOPAN, T. ...	
	Gastric Neuroses (O) ...	
	SEARCH FOR NEW TYPES OF TROPICAL SPLENOME- GALY, A. (O)—T. Bhaskara Menon ...	
	Smith, K. Shirley, ...	
	The Treatment of Angina Pectoris (G. 26) ...	
	SOME COMMON DRUGS OF CARDIAC PRACTICE (O) ...	
	T. Krishna Menon ...	
	SOME RECENT ADVANCES IN MEDICINE (O) ...	
	R. Viswanathan ...	
	Spinal Anaesthesia; Nervous System Sequelæ ...	
	George H. Hyslop (G. 4.) ...	

INDEX	273
Srinivasan, E. V.	
Welcome Address, Fourth Conference, All-India Ophthal- mological Society ...	195
Status Lymphaticus,	
Symmers Douglas (G. 1.) ...	81
STREPTOCOCCAL DISEASE, (O)	
G. Raman Pillai ...	58
Strychnine and the Barbiturates: Mutual Antidotes (G. 17)	213
Studies on Reticulocytes, (G. 9)	
P. Brahmachari ...	123
SUNDARAVADANAN, B. M.	
Note on a Case of Popliteal Aneurysm Treated by Hunter's Method (O) ...	233
SURGERY OF THE BREAST (O)	
V. Mahadevan ...	3
Symonds, C. P.	
Nervous Breakdown (G. 22) ...	249
Swanberg, Harold,	
Oral Urography, (G. 15) ...	163
THAMBIAH, S.	
The Leukæmias, (O) ...	145
Treatment of Angina Pectoris, The	
K. Shirley Smith (G. 26) ...	255
Treatment of Fracture of Spinal Column	
F. Jimeno-Vidal (G. 10) ...	124
Treatment of Small-pox,	
V. Govindan Nair (Clinical Memoranda) ...	257
Treatment of Varicosities,	
H. H. Faxon ...	215
Usman, Sir Mahomed	
Opening Speech, All-India Ophthalmological Society, Fourth Conference ...	195
VISWANATHAN, R.	
Some Recent Advances in Medicine (O) ...	23
Wright, R. E.	
Presidential Address—All-India Ophthalmological Society, Fourth Conference ...	202

CALCIUM THAT IS ABSORBED

THE daily adult requirement of calcium amounts, according to highest authority, to 0.45 gram (7 grains) per day. It can be predicted that the intake of a very large proportion of patients is far below this requirement. Our great interest in cod-liver oil, violet rays, vegetable diets, etc., should not blind us to the fact that after all they contain little or no calcium. The satisfactory use of lime requires a number of considerations, such as (1) acidity of the upper bowel, (2) alkaline metabolic end products, (3) a balanced administration of phosphorus, magnesium and the alkaline bases. These considerations are fully met in

CALCIONATES

Three teaspoonfuls the lime of 1 pint of milk
Calcionates is supplied in 4 ounce bottles.

Manufacturers:

THE UPJOHN COMPANY
U. S. A.



Further particulars from:

T. M. THAKORE & CO.,
43, Church-Gate Street,
FORT, BOMBAY.

Owned by its Policyholders

The INDIAN MUTUAL is a policyholders' institution. The policyholders' interest is the sole and the paramount interest in the working of the concern.

.....

Founded on ideal lines under an experienced management and public spirited Directors, the Association is growing stronger every day. Its progress has been phenomenal. Its constitution, methods of working and judicious husbanding of resources have attracted the attention of all thoughtful insurers. Join us to-day either as a member or as a representative.

INDIAN MUTUAL LIFE ASSOCIATION LIMITED,
69, BIG STREET, TRIPPLICANE, MADRAS.

TELEPHONE NO. 3777

CONTENTS

	PAGE
ORIGINAL ARTICLES	
1. A SEARCH FOR NEW TYPES OF TROPICAL SPLENOMEGALY By T. Bhaskara Menon, M.D.	227
2. NOTE ON A CASE OF POPLITEAL ANEURYSM TREATED BY HUNTER'S METHOD By B. M. Sundaravadanai, B.A., M.B.B.S.	233
3. DEVELOPMENT OF THE HEART, NORMAL AND ABNORMAL By P. K. KOSHY, B.A., M.B.B.S.	246
GLEANINGS FROM MEDICAL LITERATURE	248
CLINICAL MEMORANDA	257
REVIEWS	258
EDITORIALS	
1. THE PROBLEM OF TUBERCULOSIS	259
2. THE VIZAGAPATAM MEDICAL COLLEGE	260
3. COLONEL E. W. C. BRADFELD, O.B.E., C.I.E., F.R.C.S.	261
4. RETRENCHMENT IN HOSPITAL EXPENDITURE	261
MEDICAL PERIODICALS	263
ASSOCIATION NOTES & NOTICES	
INDEX FOR VOLUME ONE	

Printed by S. SESHAGIRI RAO, at the LODHRA PRESS
Published by Dr. K. C. PAUL, General Secretary
SOUTH INDIAN MEDICAL UNION, Egmore, Madras