


12

The Madras Agricultural
Journal

1941 - Jan



JL
Jm 21, N13
N 41.29.1
94540.

Regd. No. 1155.

THE MADRAS AGRICULTURAL JOURNAL

XXIX.

January 1941.

No. 1

CONTENTS

Editorial

ORIGINAL ARTICLES

1. Studies in the Barnyard Millet—
Echinochloa colona var. *frumentacea* C. E. C. Fischer 3
By G. N. Rangaswami Ayyangar
2. The Cultivation of Betel Vine (*Piper betel*)
in Poonamallee Village 12
By R. H. Krishnan and M. Kanti Raj
3. *Phaseolus sublobatus* Roxb. A New Green
Manure and forage plant 17
By K. Cherian Jacob
4. Better Methods of Virginia Tobacco Cultivation in
Guntur District 19
By M. Narasimham

SELECTED ARTICLE:

- The Man and the Plant 28
By H. Martin Leake
- Abstracts 31
Gleanings 34
Correspondence 34
Coffee Control Scheme 37
Crop and Trade Reports 38
College News and Notes 42
Moffussil News 44
Weather Review 45
Departmental Notifications 47
Additions to the Library 48

Published by

THE MADRAS AGRICULTURAL STUDENTS' UNION,
Agricultural College and Research Institute,
Coimbatore, S. India.

Annual Subscription Inland Rs. 4-0.
" " Foreign Rs. 4-8.

Single Copy
As. 6.

GREWALS NURSERY

BISHANPURA ESTATE.

P. O. PAYAL, Punjab.

Growers of all Imported and best local Varieties of Citrus Fruit plants. Winners of Silver cups, Gold medals and Certificates of Merit in Various fruit Shows. Approved in the Reliable Nurseries' list of Government of Punjab and the Horticultural Department, Patiala State.

Is one of the Biggest Citrus Nurseries in Northern India and it has produced the greatest number of Citrus plants of all varieties both imported and Country. Results obtained from these plants have satisfied thousands of people all over India.

Patiala,

15th April, 1940.

Dear Sirdar Sahib,

I am writing this to say that His Highness Shri 108 Maharajadhiraj Mohindar Bahadur was impressed by what he saw during his recent visit to Bisanpura and that His Highness greatly appreciates what you have done for the development of fruit culture in this part of the country. Keenly interested as His Highness is in the promotion of Horticulture and fruit cultivation, he was, indeed, very pleased to visit your flourishing and carefully planned and nurtured fruit farms and nurseries which should serve to give a fillip to others in the neighbouring territories to give to fruit Cultivation the attention it deserves. His Highness trusts that you will continue your efforts to popularize fruit culture in that part of the State.

Yours sincerely,

Private Secretary to

His Highness—Maharajadhiraj of Patiala.

Y. W. C. A., Asoka Road,

New Delhi.

November 20, 1940

"I have pleasure in stating that Her Excellency the Marchioness of Linlithgow in a visit to the Horticultural Fete on the 9th November in the compound of Y. W. C. A., Asoka Road, New Delhi inspected the Citrus fruit stall of Grewals Nursery, Bishanpura Estate (Patiala State) - one of the biggest Citrus Nurseries in northern India - and was very much impressed to see a very large collection of Citrus Varieties of Oranges, Grape fruits Lemons etc. Her Excellency expressed her full satisfaction over the quality of Grapefruit exhibits. The Fruit Marketing Officer who also acted as a judge explained to Her Excellency the individual characteristics and commercial importance of each variety.....

A. Tafts,

General Secretary.

Please apply for a free and Descriptive catalogue of citrus and other fruit plants to the Managing Director with the mention of this journal.

When answering our advertisers, please mention 'The Madras Agricultural Journal'

FOR QUALITY INSIST ON STANES

Well balanced Manure Mixtures for all crops. We mix only the best ingredients, and our manures are sold on guarantee of analysis. Our War-time prices are very competitive and within the reach of every Ryot.—Suppliers to The Agricultural Departments of Madras, Travancore, Cochin etc., etc.

WE SOLICIT A TRIAL ORDER FROM YOU

STANES POULTRY FOODS

For more eggs give STANES LAYING MASH.

STANES CHICK FOODS, are well balanced, stimulate growth, and make the Chicks healthy.

STANES CONDITION MIXTURE for Dogs.

STANES Fine Fishmeal, Shell-grit and Bone Grit are specialities.

STANES MINERAL MIXTURE FOR CATTLE


The only Tonic at low prices with high tonic value. Prevents diseases due to deficiency in CALCIUM and PHOSPHORUS. Largely used by all the Agricultural Research Stations, Dairy Farms, Live-Stock Stations in Madras, besides by the Travancore Royal Stables.

Sold in tins of 14 lbs., 28 lbs. and 56 lbs.

Agricultural Sprayers, Hose and Spraying Materials.

Calix No. 4. self-acting Knapsack Sprayer of 1¼ gallon capacity, made of best brass. Suitable for Copper and Sulphur mixtures, also for Health Departments.

Price very moderate.





For further particulars please apply to:—

MESSRS T. STANES & CO., LTD.,

RYOTS SALES DEPARTMENT,

COIMBATORE

When answering our advertisers, please mention 'The Madras Agricultural Journal.'



Kerala Shaving Soap



FOR ADORABLE SHAVES

*The finest gardens
in India grow on*


**POCHA'S
SEEDS!**

PESTONJEE P. POCHA & SONS,
8, NAPIER ROAD, POONA

Catalogue of Vegetable & Flower Seeds FREE.

When answering our advertisers, please mention 'The Madras Agricultural Journal.'

SPRAYERS FOR ESTATE USE Powerful and Efficient.



Manufactured by
THE STANDARD FURNITURE CO., LTD.,
KALLAI, Malabar.

THE JOURNAL OF THE
Mysore Agricultural and Experimental Union,
BANGALORE
Published Quarterly by an Expert Board of Editors.
Chief Editor : M. J. Narasimhan, B. A.
SUBSCRIPTION : RS. 3 PER ANNUM

The Madras Journal of Co-operation
The Premier Monthly Co-operative Journal in India.
Voices forth the co-operative opinion of the Province

ANNUAL SUBSCRIPTION

Inland (including postage)	Rs. 4-0
Foreign " "	" 4-12

Please apply to: **The Madras Provincial Co-operative Union,**
Royapetta, MADRAS

When answering our advertisers, please mention 'The Madras Agricultural Journal.'

DO YOU KNOW?

That Bees can increase the yield of your Fruit trees?

And that they can better the quality of your fruits?

All progressive Modern Fruit Growers in the West find it profitable to keep bees in their orchards. Keep bees and be up-to-date.

Bee-keeping is a fascinating hobby.

The Indian Bee Journal will tell you all about this extremely fascinating subject. Yearly Rs. 3.

INDIAN BEE JOURNAL.

JEOLIKOTE DIST. NAINITAL, U. P.

Kerala Soap Institute, Calicut

Stockists in Coimbatore.

1. S. Ottajee Branch, Big Bazaar.
2. S. Murugiah Gowder and Sons, Big Bazaar.
3. N. Dasai Gownder and Co., Mill Road.
4. D. S. V. and Sons.
5. P. P. Raman Nambiar.
6. Coimbatore Swadeshi Ltd.
7. Noor Mohamed Latif Sait.
8. P. Sultan Branch.
9. N. B. Krishnamurthi Chettiar, Avanashi Road.
10. Sri. Kannikaparameswari Vilas Stores,
Sullivan Street.

Agent : **R. C. Dalal, Calicut.**

When answering our advertisers, please mention 'The Madras Agricultural Journal.'

TATA'S MANURE MIXTURES

These Manure Mixtures are a combination of organic and inorganic matter balanced with considerable care. The ratio of the Nitrogen, Potash and Phosphoric Acid content is adjusted according to the purpose to which the manure is to be put. Special mixtures are available for Tea, Coffee, Rice, Tobacco, Sugarcane, Coconut, Garden Vegetable, etc. Others can be prepared on request.

Tata Manure Mixtures invigorate growth, provide resistance to disease, and increase both the quantity and quality of the yield.

It will pay you to secure full particulars of these Mixtures. Please write to:

THE TATA OIL MILLS COMPANY LTD.,

TATAPURAM

When answering our advertisers please mention 'The Madras Agricultural Journal'

The Madras Agricultural Students' Union.

This is an association of the students, past and present, of the Madras Agricultural College, and of the officers of the Madras Agricultural Department, the Principal of the Agricultural College being the President, *ex-officio*. The Union conducts this Journal and celebrates annually the College Day on which occasion it convenes an Agricultural Conference. The Association has Members, Associates, Honorary Members and Patrons. For detailed particulars apply to the Secretary.

The Madras Agricultural Journal

THE PREMIER AGRICULTURAL MONTHLY IN INDIA

EDITORIAL BOARD

K. M. Thomas, B. A., M. Sc., D. I. C., Editor.
M. C. Cherian, B.A., B.Sc. (Edin.) D.I.C., T. Nataraj, B.A. B. Sc. Ag.
M. Kantiraj, M.A., B. Sc. (Edin) K. Ramaswami, B. Sc. Ag.
N. Krishnaswami, B.Sc., Ph. D. R. Narayanamurthi, (Student)
N. Baskara Reddy, Secretary, Students' Club (*ex-officio*.)

A monthly Journal in English devoted to the study and development of Agricultural Science; contains articles of scientific as well as of general interest written in popular language. Reviews of agricultural publications and extracts from various Journals--both Indian and foreign--a special feature. Commands a wide circulation and is an excellent medium, therefore, for advertisements.

Rates of Subscription.

Journal supplied <i>free</i> to Members, Associates and Patrons of the Union				
Annual subscription rates for others:				
		Rs.	A.	P.
	India and Ceylon	4	0	0
	Foreign	4	8	0
Single number	...	0	6	0

Rates of Advertisement.

For single insertion				
	Rs.	A.	P.	
For full page	...	6	0	0
For half page	...	3	8	0
For quarter page	...	2	0	0

Liberal discounts are allowed for standing matter provided space is booked ahead and pre-payment made.

Inside cover pages 25 per cent. extra.

Outside cover pages 50 per cent. extra.

Preparation of blocks, unless supplied by advertisers, charged extra.

N. B. All advertisements are subject to the approval of the M. A. S. U. Managing Committee before insertion. The charges are strictly payable in advance.

Contributions to the Journal.

Contributions on subjects of agricultural interest or pertaining to rural economy and co-operation will be accepted for publication, subject to approval of the Editorial Board. Correspondence and queries on agricultural and allied subjects are invited. Manuscripts should be sent clearly written or typed on one side of the paper only.

Twenty-five copies of reprints of contributions will be presented to authors. Extra copies and printed covers for reprints, if required, can be supplied at actual cost if intimation is given when the manuscript is presented.

N. B. The Editorial Board does not hold itself responsible for the opinion expressed by the contributors.

All communications should be addressed to:

THE SECRETARY, M. A. S. UNION,
Agricultural College, Coimbatore, Lawley Road P. O.

The Madras Agricultural Journal.

(ORGAN OF THE M. A. S. UNION)

Vol. XXIX]

JANUARY 1941

[No. 1.

EDITORIAL

The New Year. The year 1941 has dawned none too bright on an unhappy world. Three out of five continents are witnessing the ravages of a devastating war. Destruction of life and property continues unabated. Agriculture and trade are in the throes of a depression. Science and Industry, save those that are of direct consequence to the prosecution of the war, are waning ruinously. Restricted food distribution has created slumps in producing centres and distress in consuming areas. Despite these maladjustments inevitable in a period of crisis, the outlook for 1941 is brighter and more cheery than what it was in 1940. The *Madras Agricultural Journal* enters on its twentieth year in an atmosphere of hope and faith. Increases in the cost of production and distribution of the *Journal* have been counter-balanced by increased patronage from our *clientele*. We thank our members, subscribers, contributors, advertisers and readers for their continued support and wish them all a happy and prosperous New Year.

Groundnut Trade. We had occasion to advert in a previous issue of this Journal to the sad plight of the groundnut growers in our Province and the groundnut trade as affecting India. As a result of the active interest taken by the Hon. Sir A. Ramaswami Mudaliar, the Commerce Member, the Government of India in consultation with the Food Ministry in England fixed the minimum price of groundnuts at £ 10 per ton f. o. b. Indian ports which works at Rs. 27-8-0 per candy of 531 lb. This has however not so far benefitted the growers as reports received from different parts indicate the existence of a great disparity in the prices now prevailing in the Indian markets and the minimum price offered by the Food Ministry. Thus the distress of the agriculturist and the small producer is profound and remains unsolved. It is now learnt that an arrangement has been reached between the Food Ministry in London and the Government of India for creating a fund for the benefit of the agriculturists from the proceeds of a rebate of 25 shillings per ton from South Indian shipments. This fund is proposed to be distributed to the growers on an acreage basis. We daresay that some such action to relieve the distress of the groundnut growers is urgently necessary to ease the feeling of despondency prevailing in the villages. We are encouraged by the recent press reports that the Government of Madras and some other Provincial and State Governments have agreed to foster the manufacture of products such as vegetable ghee, varnishes, lubricants and soaps from ground-nut oil. It is also understood that the Government of India are in correspondence with the Provincial

Governments in this connection. It is well known that groundnuts cannot be stored long without deterioration and it would greatly benefit our growers if this scheme is put on hand expeditiously.

We would urge that the question of utilizing groundnuts in India itself is tackled on another front also. That the groundnut is a cheap source of protein and a valuable article of food is beyond doubt. How far and in what manner it can be included usefully in South Indian diets needs a careful study at the hands of our nutrition experts. Such an investigation coupled with effective propaganda will no doubt stimulate the consumption of groundnuts within our own country. Further the increased crushing of groundnuts would release large quantities of groundnut cake which is a very rich source of proteins for our cattle, besides its being a valuable manure particularly for sugarcane and paddy crops. The price of nitrogenous fertilizers has shot up prohibitively since the outbreak of the war and imports have dwindled. Oil cakes could well replace them now at a much cheaper rate.

We are glad that the recent conference of representatives of groundnut interests at Madras has focussed the attention of all concerned to the pitiable plight of our ryots and we hope that their sustained efforts would ere long afford the agriculturists the much needed relief.

An Agricultural Pests and Diseases Act for Bombay. The decision of the Bombay Government to enact a legislation on the lines of the Madras Agricultural Pests and Diseases Act of 1919 is a move which has not emanated too soon. Adequate control of insect pests and diseases is a matter of vital importance to the cultivator in India as elsewhere. In the matter of pests and diseases, the policy of Agricultural departments in India has been to bring to the door of the peasant cultivator the necessary knowledge about methods of control. With many pests and diseases this is all what is needed, since the adoption of the measures gives the cultivator relief, while failure to follow up demonstrable remedies does harm only to himself. But there is another important category of pests and diseases where the range of spread is not confined to the limited area of an individual holding. In such cases, the negligence on the part of the few would not only do harm to themselves but would render the efforts of their neighbours futile. The negligent and the unbelieving would thus become a danger to society so that the protection of the crops belonging to the many by compelling the few to fall in line with the rest, becomes a paramount duty of the state. The enactment of a piece of legislation making it incumbent on all cultivators in a specified tract to adopt a uniform procedure, is the only means of mitigating this evil. Failure to carry out the measures would authorise a state agency to enter the holding, carry out the operations and recover the cost from the delinquent. Wilful infringements of the regulations would also render him liable for prosecution. Legislation on these lines has worked successfully in Madras and her neighbouring states for well over two decades and we trust that Bombay will be no exception in deriving benefit from the contemplated enactment.

Studies in the Barnyard Millet—*Echinochloa colona* var. *frumentacea*, C. E. C. Fischer.

By G. N. RANGASWAMI AYYANGAR, F. N. I., I. A. S.,
Millets Specialist, Agricultural Research Institute,
and

U. L. SRINIVASA RAO,
Sub-Assistant, Millets Breeding Station, Coimbatore.

The Barnyard millet, known in Tamil as *Kuthiraivali* (=horse-tail), is one of the less important millets of the Madras Presidency. Except in certain restricted tracts its importance as a food crop in India is not much. In view of the fact that it forms a good famine crop and also comes to the rescue of very poor cultivators, some of its characters were studied at the Millets Breeding Station and are presented below:—

This plant is considered to have been first brought into cultivation in India. That it has been grown all over India since very remote times is seen by the fact that it has a name in Sanskrit, as well as in every one of the other Indian languages.

Werth (1937) mentions this plant occurring as a weed of cultivation in the temperate and tropical zones of both hemispheres, particularly the Northern and that it is cultivated as a millet (as a poor man's millet) in India, China, Japan, Dutch India and in smaller quantities in Africa. In China and Japan particularly it is said to be used as a substitute crop when paddy fails.

Its composition is given by Church (1886) as follows:—

		In 100 parts.
Water	...	12.0
Albuminoids	...	8.4
Starch	...	72.5
Oil	...	3.0
Fibre	...	2.2
Ash	...	1.9

The nutrient ratio here is 1 : 9.5 and the nutrient value is 88.

This millet is used in India either boiled in water like rice, or parched or boiled with milk and sugar.

As a fodder it seems to have attracted greater attention especially in the U. S. A. Bressman and Fry (1932) consider this plant as the best late season feed, taking the place of maize in certain parts, which are unsuitable for maize. Thatcher (1900) gives the following composition of this plant as a forage crop.

Composition of feeding stuff at different stages of growth.

Feeding stuffs	Water	Protein	Albuminoids	Ether extracts	Nitrogen free extracts	Crude fibre	Ash
Millet: Heads just appearing	10.24	8.41	5.79	2.54	32.03	35.86	10.92
Headed out seeds near ripe.	10.47	6.12	4.49	1.52	43.33	29.10	9.41

He concludes that here the protein decreases rapidly while heading out and to obtain a fodder having as narrow a ratio of flesh-forming to fat-forming foods as possible the crop should be cut at as early a stage as it can be well cured. As a roughage it can be allowed to grow till seeds are formed.

Lindsey (1900) gives the following coefficients of digestibility of this plant: green, as hay and as silage with soybean, obtained with sheep.

Coefficients of digestibility of millet, millet hay etc.

	Dry matter	Protein	Fat	Nitrogen free extracts	Crude fibre	Ash.
	%	%	%	%	%	%
Green millet, early to late bloom	71	69	63	72	73	64
Millet hay, full bloom	56.5	47.5	48	53.5	62	43.5
Millet and Soy bean silage	59	57	72	59	69	—
Corn and soy bean silage	69	65	82	75	65	—

He points out that when harvested early, in blossom, the fodder contains less nitrogen-free extract matter, more fibre or woody matter and rather more ash than corn fodder and so it must be cut when in blossom, to secure it in the most desirable condition for feeding.

On well-manured soils it gave 11,297 lbs. of straw and 66.7 bu. per acre and 12–15 tons per acre of green forage. Wood (1928) gives the yield of straw as 2,000 lbs. per acre, Mukerji (1915) records 800 lbs. and Mollison (1901) 1,500 lb. Watt (1889) states that the straw is used much in the Madras Presidency and Mysore as cattle fodder, though considered inferior to ragi as well as to paddy straw.

Kuthiravali is sown mostly as a rainfed crop. In the drier districts, it is grown as a subordinate crop to sorghum (Duthie and Fuller 1882) or maize (Watt l. c.) The plant can be grown either on light sandy soils (Duthie and Fuller, l. c.; Mollison, l. c.) with fair rainfall or in water-logged areas (Wood, l. c.; Watt, l. c.), such as lowlands or banks of rivers that often get submerged. Mukerji (l. c.) reports that rough jungle land could be used. It withstands transplanting (Mollison l. c.) The crop requires little or no manuring.

The seed rate is generally 8–10 lbs. per acre and when sown with a drill about 6–8 lbs. per acre. Wood (l. c.) quotes 35 lbs. per acre. The July sown crop is harvested in October.

In Punjab this plant is said to be ploughed into the soil as a green manure.

Systematics. Hooker (1897) brings this plant under the sub-section *Echinochloa* of the genus *Panicum*. He recognises two very closely allied species viz. *Panicum crusgalli*, Linn. and *Panicum colonum*, Linn. The cultivated species is regarded as a variety of *P. crusgalli*. In his species of *P. crusgalli*, Linn. he finds innumerable forms so that it became "impossible to find characters constant enough for their limitation". He finds that *P. colonum* Linn. exhibits a gradual transition into *P. crusgalli* and vice versa. The cultivated variety *frumentaceum* has been assigned to both. Cosson and Thwaites are reported having regarded *colonum* as a form of *crusgalli*.

Cook (1908) mentions that the *Panicum crusgalli* of Linnaeus does not occur in the Bombay Presidency. He considers the cultivated form as a variety of *P. stagnium*, Retz.—*P. stagnium*, Retz., var. *frumentacea*, Trin. The species *crusgalli* has no ligule. *P. frumentaceum*, Roxb. and *Oplismenus frumentaceus* Dalz. and Gibs. are given as synonyms of *P. stagnium* var. *frumentacea*. He remarks that *P. colonum* is very similar to var. *frumentaceum*, Roxb. but a more slender plant with smaller spikelets.

Chevalier (1922) quotes the following synonyms for this plant: *P. frumentaceum* Roseb., *P. grossum*, L. *P. segetale*, Roxb., *Echinochloa frumentacea*, Link., *Oplismenus frumentaceus*, Kunth. *P. frumentaceum* is said to be a variety of *P. crusgalli*, L. It is considered by most authors that *P. frumentaceum* possesses characters intermediate between those of *P. crusgalli*, L. and *P. colonum*, L.

Bressman and Fry (1932) distinguish the weed *crusgalli* from the cultivated millet by its greater hairiness of the glumes, and the longish seeds. The millet is less hairy and has broader seeds.

Fischer in Gamble (1934) removes the group from *Panicum* and gives the plants a separate generic stand, viz., *Echinochloa*. Under this genus he enumerates 3 species:—

1. *Echinochloa colona*, Link. = *P. colonum*, L.
2. *Echinochloa crusgalli*, Beauv. = *P. crusgalli*, L.
3. *Echinochloa stagnina*, Beauv. = *P. crusgalli*, L.

The cultivated species is treated as a variety of *colona* viz. *E. colona* var. *frumentacea*, C. E. C. Fischer = *P. crusgalli*, Linn. var. *frumentaceum*, Hook.

The following is the description of the species as given by Gamble:—

Echinochloa, Beauv. Annual or perennial, often tall herbs. Leaves narrow. Inflorescence of crowded panicles of loosely arranged, secund, spiciform branches bearing spikelets from the base or near it; rachis triquetrous. Spikelets ovate

to elliptic or lanceolate-oblong, 2-nate or clustered, articulated on and falling entire from the pedicels.

Glumes membranous, unequal; the lower much the shorter, mucronate, cuspidate or awned, the upper coincident in outline with the spikelet, acute, cuspidate or shortly awned. Lemmas: Dissimilar; the lower equalling the upper glume (excluding cusp or awn) its palea 2-keeled, empty or containing a male floret; the upper subcoriaceous or crustaceous, ovate, to elliptic-oblong, obtuse or apiculate, polished, very convex on the back, its pales as long, with rounded sides and flaps, containing a bi-sexual floret. Lodicules 2. Stamens 3. Styles free. Grain broadly elliptic, plane-convex.

Racemes simple, rather distant 3-1.25 in. long; lower glumes and upper glumes about equal, obtuse or cuspidate. Annual, up to 2 ft. high; leaves 2-8 in long, 1-4.5 in. wide, ligule 0; spikelets ovoid 1-1.2 in. long; lower glume .04-.05 in. long; upper .09-.11 in. long; lemmas .08-.1 in. long lower with male floret.....*colona*.

Racemes usually more or less branched 8-2 in. long; lower glume and upper lemma cuspidate or awned, the latter the longer; lower lemma often awned:

Annual, up to 3 ft. high leaves 3-21 in. long; 2-5 in. wide, ligule 0, junction of blade and sheath glabrous, usually marked by a brown zone; spikelets 15-18 in. long; lower glume .07-.12 in. long, upper .15-.17 in. long; lower lemma empty, .14-.17 in. long, upper .12-.15 in. long; awn of lower lemma, up to 2 in. long.....*crusgalli*.

Usually perennial, up to 6 ft. high; culm rooting and often branching from submerged nodes; leaves 3-18 in. long, 2-4 in. wide, ligule a fringe of stiff hairs, sometimes absent on the uppermost leaf; lower lemma empty or with a male floret. Otherwise as in the last species.....*stagnina*.

Echinochloa colona (Link.) var. *frumentacea*, C. E. C. Fischer: a taller and more robust plant with dense, sometimes corymbose panicle, cultivated.

Observations on flower pollination. Hildebrand (in Knuth 1909) reports that only self-pollination is possible in this species, owing to the simultaneous protrusion of stigma and anthers, but crossing may be effected when the anthers have fallen, as the stigmas are persistent.

Youngman and Roy (1923) found the flowers opening between 7-30 and 8-30 a. m. They observed that the stigmas and anthers emerge simultaneously. The stigmas spread out immediately on emerging, while the anthers dehiscence only after about 1-1½ minutes. The glumes are observed to close back after half an hour.

Observations made at the Millets Breeding Station, Coimbatore on two types of panicle shapes, viz., open and compact, gave the following results. The plants were grown under irrigated conditions. The observations were done on three plants in each type. The plants commenced to flower in about two months after sowing. The tip of the leaf subtending the panicle (flag leaf) appears first and takes about 10 to 14 days for the complete emergence of the flag. The appearance of the panicle is almost simultaneous with that of the flag, but its emergence is gradual, taking from 9 to 13 days.

The flowers begin to open in the same order as they emerge out of the sheath. The first flowers open as soon as the panicle tip emerges out. The order of flowering is thus from the tip of the panicle to the base. But in the

individual spikes the spikelets along the two margins open earlier than those at the middle. The flowering period is 19-22 days in the open panicles and 2 or 3 days more in the compact ones. The largest number of flowers open during the sixth to eighth day from the commencement of flowering. The opening of the individual flowers is between 5-10 a. m. The maximum number of flowers open between 6 and 7 a. m.

The glumes open out very gradually and at the mouth of the gaping glumes the two stigmas and the anthers stand out like a column. Two to four minutes later the stigmas emerge out of the lemma in a column and spread out on either side. The stamens thus come to occupy the central position. The filaments begin to elongate gradually in about 1 to 8 minutes after the spreading of the stigmas. The dehiscence occurs only when the filaments have elongated to their maximum. The dehiscence is by lateral sutures. It begins at both ends and meets in the centre. The glumes close again in about 5 to 10 minutes after the dehiscence. The stamens and stigma remain outside the closed glumes. The whole process, from the opening of the glumes to their complete closing, takes about 24 minutes.

Inheritance of characters.

I. *Anthocyanin pigmentation.* The only anthocyanin pigmentation met in this plant is purple. When the plants do not show any trace of the pigmentation they are designated as 'green through-out' (abb. GT.) The purple pigment normally manifests itself in the following regions of the plant:

(a) *Vegetative parts*—nodes, internodes, as two bands on the upper and lower sides of the nodes, leaf margins, midrib, sheath, panicle-rachis, glumes.

(b) *Reproductive parts*—lemmas, anthers and stigma

Three grades of pigmentation designated P₃, P₂, P₁, in descending order of their intensities, are met with. The anthers and stigmas show various colorations on drying corresponding to the grade of the plant-pigmentation. The following gives the chief differences between the various types:

Class of pigmentation.	Vegetative Parts.	Character and incidence of pigmentation			
		Reproductive parts			
		fresh	dry	fresh	dry
P ₃	Deepest purple of all types	deep purple	deep blue	deep purple	black
P ₂	Pigmentation less than in P ₃	purple	a blue ring around the sutures	purple less than in P ₃	dark brown
P ₁	Pigmentation less than in P ₂	yellowish brown	brown	light purple	brown
GT	No purple pigmentation, all green	yellow	brown	colourless	pale brown

A certain amount of fluctuation in the depth of pigmentation occurs in each of the three types. In the following segregations it was found that P.3 and P.2 groups were often rather difficult of separation and in certain cases the P.1 almost approached P.2.

(a) *Purple and green segregations:—*

Two kinds of segregations were met with, viz., a monofactorial and a bifactorial one.

The bifactorial segregations gave the following F_2 proportions.

Female parent = Green throughout

F_1 = P.2

F_2 :—

	P.3	P.2	P.1	GT
Observed	266	80	88	44
Expected on 9:3:3:1	268.9	89.6	89.6	29.9
	$X^2 = 7.73$	$P > .05$		

In the monofactorial segregations the following F_2 proportions were obtained :—

Female parent = green throughout.

F_1 = P.1

F_2 :—

	P.2	P.1	GT
Observed	29	71	33
Expected on 1:2:1	33.25	66.5	33.25
	$X^2 = 0.84$	$P > 0.50$	

The same proportions were met with in the further progeny of the hybrid. (Total of 9 families gave P.2 = 330; P.1 = 628; GT = 329. calculated P.2 = 321.75; P.1 = 643.5; GT = 321.75). This clearly indicates a 1:2:1 ratio. According to expectations all the P.1 selections proved to be heterozygous.

(b) In segregations between the three pigmentation groups, the 1:2:1 proportions were again met with :—

	P.1	P.2	P.3
Total of 20 lots	751	1540	890

The P.2 group is smaller and the P.3 group larger than they should be for an exact 1:2:1 ratio, because their separation is not quite easy, the obvious P.2 alone going into the middle group. Subsequent selections however showed that the P.2 were all heterozygous and the P.3 and P.1 selections bred true.

It is evident from these that there are only two factors responsible for pigmentation in this plant. Further, it is evident that pigmentation differences between the heterozygotes and the pure ones are easier to be noted in the monofactorial segregations than in the two factor ones.

The factors may be designated as follows :—

$P_1 P_1 P_1 P_1$	P.3	$P_2 P_2 P_1 P_1$	P.1
$P_1 P_1 P_2 P_2$	P.2	$P_2 P_2 P_2 P_2$	GT




Fig. 1. (Top) Inheritance in ear-head shape. (Left) Compact head; (Right) Open head; (Middle) F_1 ; Single branches:— Top—cross section; Middle—rear and bottom—front view.

Fig. 2. (Bottom) Striped plant.

II. *The Panicle* (Plate I, Fig. 1). The panicle is conical in appearance. The number of panicles on a plant depends on its branching and tillering capacity. Each panicle is peduncled.

The peduncle is cylindrical and is continuous with the rachis, which however is angular. The spikes arise in whorls of 2 or more often 3—4, or sometimes 5. Each whorl thus constitutes more or less a node. The disposition of the whorls is more distant at the base and becomes less distinct and more congested towards the apex. The spikes arise on the face and not on the angle itself. They alternate in successive whorls so as to give a $\frac{1}{2}$ taxis. Thus the panicle shows a tetraquetrous arrangement. Consequent on the congestion towards the apex the whorls may become disturbed but still the tetraquetrous nature is kept up. The length of the spikes reduces gradually from the base towards the apex and the ultimate apex ends in a spike. This arrangement is responsible for the conical shape of the panicle.

The spikes are more or less ascending. The base of the rachis is somewhat pulvinate. The node at the base of the pulvinus is pubescent. Hairs, isolated or in tufts of 2—3 are found on the rachis, at the base of the spikelets. Groups of spikelets are arranged distichously along the dorsal side of the rachis. These groups consist of 3—4 spikelets distichously arranged on a short rachilla. The spikelets are all more or less equal in size. In general design they appear to be arranged in horizontal rows of 4 to 5 spikelets. The rachis attenuates towards the apex. The number of spikelet groups, however, remains the same and consequently the spikes tend to curve on their ventral side. This becomes more pronounced as the grains begin to mature. The spike ends may be seen almost spirally twisted and bent towards one side to avoid overlapping the upper one, so that the panicle gets a plaited appearance.

Two distinct types in head-shape could be distinguished, viz., Open and Compact.

In an open panicle the spikes are more or less horizontal and may or may not be curved towards the tip. The interval between the whorls is greater than in a compact panicle. The length of the spike is also greater and the spikelet groups are more spread out.

The compact panicle on the other hand shows the spikes much plaited. The spikes are much shorter and spikelets are very close together.

Further, the spikes in the 'opens' have a greater number of spikelet groups with usually 2—3, often 4, and rarely 5 spikelets per group. The 'compacts' on the other hand have usually only 3-4 spikelets per group, often 2 and very rarely 5.

An intermediate condition between the lax-ness of the 'opens' and the plaited appearance of the 'compacts' is met with. This is termed the 'semi-compact' head. In this the spikes do not completely overlap one another so that the main rachis becomes visible.

An analysis of typical heads gave the following results:—

Panicle type	Average number of spikelets	Average length of spike	Average number of spikelets per cm.
Open	26	2.8	9
Compact	22	1.9	12

Thus it becomes apparent that the headshapes are distinguished by the difference in the length of the spike and secondly the denseness of packing of the spikelets, i. e. the number of spikelets per cm. A similar experience is met with in the *E. coracana* also. (Ayyangar et. al. 1932).

III. *Sterility*. A case of male sterility was noted in a purple-pigmented plant. The emerging anthers instead of being purple were seen to be yellow and shrivelled up later without dehiscing. The anther sacs were devoid of free, healthy, pollen. This character was found to behave as a monogenic recessive to the normal condition.

IV. *Striped plant* (Plate I, Fig. 2). A single case of albino-striping was met with. This plant produced 31 tillers. The first was half albino and half green. The next six were all green and 4 white. Then again were formed seven green and the rest 10 white. Three tillers developing from the first were found to be green, though the parental one was half green and half albino. This is probably due to a chimaera, the mutation having taken place very early, probably even in the seed itself and affecting one half of the embryonal growing point, so that about half of the tillers is striped or white and the other half is green.

Seeds were collected from 27 tillers individually and progenies raised. The seed-setting in the heads from the green tillers was better (average 293 grains per head) than in those from the white ones (average 159 grains per head). The viability of the seeds was good (88% germination in seeds from green and 84% from those of white tillers). The progeny in either case (i. e., from the heads from green portion and that from white portion) gave green and pale seedlings. The proportions of green to pale seedlings were widely different:—

i. Progeny from green tillers

	Green	Pale	Ratio
Total of 7 tillers	1074	350	3:1
" 3 "	511	57	15:1 (approx)
" 1 "	144	102	9:7
" 3 "	254	327	7:9

ii. Progeny from white tillers

	Green	Pale	Ratio
Total of 4 tillers	414	168	3:1
" 5 "	658	46	15:1
" 1 "	58	50	9:7
" 2 "	95	119	7:9

iii. Progeny from the half green and half white tiller

Green	Pale	Ratio
103	206	1:3 (approx)

On the total the progenies of this plant gave 3311 green seedlings and 1425 pale seedlings. Some of the pale seedlings that were allowed to

grow further put forth only green leaves. Neither the green nor the pale seedlings produced tillers with striping or albino.

Cytology:—Hector (1936) gives the following resume. "The chromosome number was first reported as ca. 48 (2n). According to Church (1929) the haploid number is 21. Avdulov (1931) found 54 (2n). Hunter (1934) 36. As the basic number of the Paniceae is 9, Avdulov's and Hunter's figures would appear to indicate polyploidy."

Echinochloa colona, Link. is a close relation of this plant which is cultivated as a millet in several parts of India and occurs wild all over the country.

E. colona appears to hybridize spontaneously with *E. colona*, var. *frumentacea*. The two plants can be distinguished easily by means of their panicle characters:

	<i>E. colona</i> ,	var. <i>frumentacea</i>
Rachis	Flat, triquetrous	Tetraquetrous
Arrangement of spikes	Bilateral, alternating	Whorled to spiral
Disposition of the spikes	Almost vertical and adpressed to rachis	Horizontal to ascending, divergent
Spikelets	Solitary or in twos, pedicelled	Always in groups of 3-5, sub-sessile.

The *colona* species itself sometimes shows a tendency to whorling. The weaker and later-formed panicles of *frumentacea* tend to be less whorled. The progenies of some natural hybrids gave in the F₂ a large number of intermediates with regard to the number of spikes and also the manner of arrangement.

Summary. *Echinochloa colona* var. *frumentacea*, originally classified under Paniceae (*Panicum crusgalli* var. *frumentaceum*), though one of the less important millets of India forms a good famine time fodder and grain plant.

It can be fed green or as hay, or as silage. The plant has the advantage that it can grow in poor soils, as also under water-logged conditions.

Three pigmentation types P. 3, P. 2, P. 1 and one non-pigmented type GT. have been described. These show a two factor difference.

The panicle-shapes are of three types—Open, semi-compact, and compact. These are considered to be due to differences in the length of spikes and density of the spikelets. The relationships of these three types are yet not fully clear.

A type of male-sterility, simple recessive to the normal condition has been described. Albino-striping was met with and its progeny has given seedlings with green and pale colours in various proportions.

Literature cited.

- Ayyangar, G. N. R. et al. 1932. *Ind. J. Agri. Sci.* 2:254.
Bressman, E. N. and E. S. Fry. 1932. *Jour. Amer. Soc. Agr.* 24.
Chevalier. 1922. *Int. Rev. Sci. and Practice. Agri. N. S.* 1:671.

leaves which find a ready market. The *agathi* crop has to be irrigated from time to time and in about three months the standards will be about four to five feet high.

(iv) *Planting cuttings.* The next operation is the preparation of the flat surface of the ridges, to plant the vine cuttings. The surface is levelled and the edges are slightly raised to hold water. Cuttings with at least three nodes are planted in a slanting position, so that two of the three nodes are well covered by soil.

(v) *After cultivation.* These newly planted cuttings have to be irrigated (usually by splashing water stored in the trenches) four or five times a day during the first fortnight. During the third and the fourth weeks, watering is done twice a day and afterwards once a day. The cuttings strike root after about 30 to 40 days, and fresh shoots appear from about the 60th day onwards. About four or five months after the planting of cuttings, long bamboo poles (*agathi* stems from old gardens can be got at a cheaper rate) are tied horizontally to the *agathi* standards about six feet from the ground level. Out of the four rows of *agathi* standing on the ridges, two from one ridge and two from the next are bent and tied together just above the middle of the trench in the form of an arch. The growing shoots are generally trailed on the standards once in ten days or a fortnight. In about six months from the date of planting, the vines would have grown up to a height of about seven feet and the first picking of leaves can be done. If the vines are allowed to grow tall, gathering of leaves will become difficult. So, once in the first year and twice in the succeeding years, the vines that have grown above seven feet are brought down, the lower four feet coiled and covered up with earth. The distal ends of the shoots are trailed again on the 'standard'.

(vi) *Harvesting.* The first picking of the leaves can be done at the end of the sixth month after planting. In the first and second years the leaves are gathered once in a fortnight, while in the third year, the interval between two successive pickings is reduced to ten days and the quality of leaves obtained is slightly inferior. Usually the cultivators themselves and the members of their family attend to picking of leaves. Occasionally, outside labour is engaged, and each labourer is paid at the rate of one to one and a half annas for a basket full of leaves (6,400) gathered. Apart from the betel leaves, the growers gather tender *agathi* shoots at intervals of about one month and sell them in the market. There is a great demand for it as 'greens' and for feeding milch cattle. From the plantain suckers, about 175 bunches are got by the end of the first year. After harvesting the bunches, the mother plants are cut off, leaving one side-sucker in each stool. Thus, every year, about 175 bunches of plantains are also obtained; besides these, plantain leaves are also cut and sold. In addition, vegetables like brinjals, chillies, etc., are also grown (as a catch crop) here and there in the gardens. They are mostly used for domestic consumption and rarely sold.

Cost of cultivation per acre.

Details.	1st year.	2nd year.	3rd year.
(a) <i>Expenditure.</i>	Rs. a. p.	Rs. a. p.	Rs. a. p.
1. Digging trenches, 2 ft. broad and 1½ ft. to 2 ft. deep, 4 ft. apart, @ 2½ annas for 50 running feet of the trench. ...	20 0 0		
2. Marking lines on the ridges making holes 6" apart on the lines and sowing <i>agathi</i> seeds 12 men at 4 annas each. ...	3 0 0		
3. Cost of <i>agathi</i> and <i>thagarai</i> seeds (24 M. M. at Rs. 2-8-0 per M. M.*) ...	60 0 0		
4. Cost of 200 plantain suckers at 1 anna each. ...	12 8 0		
5. Cost of labour for planting the suckers 8 men at 4 annas each. ...	2 0 0		
6. Cost of watering <i>agathi</i> on alternate days from the time of sowing to the time of planting the betel vine cuttings (3 months) at Re. 1-4-0 per watering. ...	56 4 0		
7. Taking mud from the trenches, plastering the sides of the ridges and rectifying the ridges just at the time of planting the cuttings and subsequently once in a month (9 times in the first year and 12 times in the succeeding years) 30 men for each operation at 4 annas each. ...	67 8 0	90 0 0	90 0 0
8. Cost of betel vine cuttings 135 bundles of 220 cuttings each at 3 bundles for a <i>Varahan</i> (Rs. 3-8-0). ...	157 0 0		
9. Cost of labour for planting the cuttings 20 men at 4 annas each. ...	5 0 0		
10. Cost of irrigation during the first fortnight (splashing water 4 or 5 times daily) at Rs. 2 per day. ...	30 0 0		
11. Cost of irrigation during the second fortnight (splashing water twice a day) at Rs. 1-8-0 per day. ...	22 8 0		
12. Subsequent irrigations (splashing water from the trenches) once a day at Re. 1-4-0 per watering—about 240 irrigations in the first year and 300 irrigations in the succeeding years. ...	300 0 0	375 0 0	375 0 0
13. Cost of tying up the cuttings to the <i>agathi</i> stems once in a fortnight from the 6th month onwards (12 times in the first year and 24 times in the second and third year) 10 men at 4 annas each for each operation. ...	30 0 0	60 0 0	60 0 0
14. Cost of 3,500 bamboos for tying the <i>agathi</i> standards at Rs. 5 for 120 bamboos. ...	137 8 0		
15. Cost of ropes required for tying the above. ...	15 0 0		
16. Labour required for the above item, 40 men at 4 annas each. ...	10 0 0		
17. Cost of labour for picking leaves at 1½ annas for basketful of leaves (6,400)—four pickings in the first year, 24 in the second and 36 in the third year. In the first and second years 100 baskets per picking and in the third year only 50 baskets per picking. ...	31 4 0	187 8 0	140 10 0

* M. M. = Madras Measure (roughly three poun's).

Details.	1st year.	2nd year.	3rd year.
	Rs. a. p.	Rs. a. p.	Rs. a. p.
18. Cost of labour for gathering <i>agathi</i> leaves twice in the first year, 10 times in the second year and 6 times in the third year.	3 0 0	15 0 0	9 0 0
19. Manuring, twice in first year 4 times in the second and third years at 25 cartloads each time at Re. 1-8-0 per cartload.	75 0 0	150 0 0	150 0 0
20. Cost of labour for bringing down the grown up vines making coils of the lower portions, putting soil over them and trailing the young shoots to the standards—twice in the second year and twice in the third year, 80 men at 4 annas each for each operation.		40 0 0	40 0 0
21. Lease amount (Land).	100 0 0	100 0 0	100 0 0
22. Cost of levelling the fields and bringing to the original condition.			14 14 0
Total.	1.138 0 0	1017 8 0	979 8 0

Grand Total. ... Rs. 3135/-

(b) Receipts.

1. By sale of betel leaves at Rs. 2 per basket of 6,400 leaves—1st year, 4 pickings, 100 baskets for each picking. 2nd year, 24 pickings, 100 baskets each time, third year 36 pickings, 50 baskets each time.	800 0 0	4,800 0 0	3,600 0 0
2. By sale of <i>agathi</i> leaves at 2 annas a bundle—400 bundles in 2 cuttings in the first year, 2,000 bundles in 10 cuttings in the second year, and 900 bundles in 6 cuttings in the third year.	50 0 0	250 0 0	112 8 0
3. By sale of plantain bunches (175) at 6 annas each.	65 10 0	65 10 0	65 10 0
4. By sale of plantain leaves, vegetables grown in the garden, etc.	10 0 0	50 0 0	30 0 0
Total.	925 10 0	5,165 10 0	3,308 2 0

Grand Total.	Rs. 9,899 6 0
Gross income for three years	.. 9,899 6 0
Gross expense for three years	.. 3,135 0 0
Net profit for three years	.. 6,764 6 0
Net profit per annum	.. 2,254 12 8 or Rs. 2255/-

This amount has to be shared by about 15 cultivators, so, the net gain per cultivator per annum, will be about Rs. 150.

(vii). *Marketing.* The chief market for the leaves, is the city of Madras. Every day about 3 p. m. all the betel vine growers of the village assemble in a common place with the gathered leaves, cleanly washed, counted and arranged in baskets. The unit of sale is one *Kavuli* or hundred leaves. Merchants from the city come to the spot, the price for the day is fixed, and sales are effected.

Some years back there was considerable demand for these leaves from Northern India, but now-a-days those far off markets, have been reported to be captured by the betel vine growers of Sankaridrug (Salem District) and other places.

Pests and Diseases. The betel vines of these parts do not suffer from any insect pest, but the wilt disease is causing considerable damage. About 20 years ago each garden is reported to have fared well up to six years. Gradually, due to wilt disease, the life has been reduced and the gardener must now consider himself lucky if it survives for full three years. Recently, cases have been noticed where the vines have begun to wither even in the second year. The main cause of wilt disease is reported to be due to (i) cultivation of vine in the same land without adopting any rotation, or (ii) keeping the trenches always moist, a condition much favourable to the growth of the fungus. The ryots abandon the gardens when the disease has advanced to a fair extent.

The *agathi* 'standards' have been found to suffer from two insect pests, (a) *Agathi* stem borer, (*Azygophleps scalarids*) and (b) *Agathi* weevil (*Alcides bubo*). Generally these appear only as minor pests, but some times the attack may be severe. When the *ogathi* plants are very young, if these pests appear on a large scale, the plants never grow above two feet, and in such cases also, the ryots abandon the garden and start afresh.

Phaseolus sublobatus Roxb. A new green manure and forage plant.

By K. CHERIAN JACOB, L. Ag., F. L. S.

Agricultural Research Institute, Coimbatore.

A specimen of a leguminous plant (*Phaseolus sublobatus* Roxb.) known in Tamil as *Karum payar* was received from Sri. C. S. Seshagiri Ayyar, Agricultural Demonstrator, Perambalur, Trichinopoly District, for identification and on requisition he has given the following information regarding its cultivation in the Trichinopoly taluq.

An area of about 10,000 acres is grown under this plant in the Trichinopoly taluq as a mixture with irrigated *Cumbu* (*Pennisetum typhoides* Stapf and Hubbard) from April-May to July-August. After the harvest of the *cumbu* crop, the *Karum payar* crop is allowed to be grazed by cattle or used as green manure by puddling *in situ* for the next paddy crop. The plants which are grazed by cattle will soon shoot up and give sufficient quantity of green manure. The crop is also raised as a mixture in dry lands with cholam (*Sorghum* sp.) red gram (*Cajanus Cajan* (Linn) Millsp.), etc., and from this crop seeds are gathered for sowing in the next season. Seeds are not collected when it is grown in the wetlands. This green manure crop is preferred by the ryots first because it does not smother *cumbu* like

Daincha (*Sesbania bispinosa* (Jacq.) Fawcett and Rendle) and secondly it readily decomposes when puddled unlike *Daincha* which develops much wood if the paddy sowing season is delayed.

Phaseolus sublobatus Roxb. (Tam: *Karum payar*; *Mukani*, *Mataki*, Bombay; *Ghora Mung*, (Assam) is very closely allied to *Phaseolus aureus* Roxb. (green gram), of which it may be the wild form. It is also allied to *Pillipesara* (*Phaseolus trilobus* Ait.) the famous green manure plant of the Telugu area. *Phaseolus sublobatus* Roxb. is more robust than *Pillipesara*.

It is found in the hills of Deccan and Western Ghats up to 6,000 feet altitude. It also occurs in the Konkan, Bihar, Bengal and Ceylon.

The seed is rich in protein and is largely used as food in the Deccan during famine periods.

William Roxburgh in *Flora Indica* describes it as an annual while J. D. Hooker in *The Flora of British India* and T. Cooke in *The Flora of the Presidency of Bombay*

describe this plant as a perennial herb. The following description of the plant will be helpful for its identification. Stems twining when meeting support, sometimes suberect, slender, clothed with spreading or deflexed reddish-brown hairs. Leaves three-foliolate; petioles 6-9.5 cm. long, channelled, very hairy; stipules 8 mm.-1.5 cm. long, ovate-oblong, hairy, slightly ciliate, attached a little below the middle. Leaflets, the terminal the larger and ovate or rhomboid-ovate, equal-sided, with cuneate base, 4.5-8.5 cm. long, usually as broad; the lateral ovate-acute, inequilateral, with rounded or truncate base, 3-8 cm. long, usually as broad, acute, silky-hairy on both sides, occasionally somewhat three-lobed, conspicuously three-nerved from the base; petioles, middle one, 1.8-2.2 cm. long, the lateral ones 3-5 mm. long, hairy; stipels linear-subulate. Flowers in short close 6-12 flowered racemes with swollen nodes; peduncles 1-3 cm. long, hairy; pedicels very

Phaseolus sublobatus, Roxb. short, 2 mm. long; bracts ovate, nearly 2 mm. long acute, ciliate, deciduous; bracteoles 3 mm. long, linear-subulate, ciliate, prominently nerved, deciduous. Calyx 3 mm. long, glabrous, shortly ciliate; teeth, shorter than the tube, deltoid.



Corolla 8 mm.-1.2 cm. long, yellow. Pods 5.5 cm. long, slightly compressed straight, clothed with short reddish-brown hairs. Seeds 8-12, oblong with truncate ends, 2-3 mm. long, dark-brown and mottled.

This promises to be a very useful green manure plant suitable to be grown both in dry and wet lands. Cattle readily graze this even though it is beset with hairs. It therefore deserves to be tried under diverse conditions of soil and climate.

Bibliography.

- Roxburgh, W. (1832). *Flora Indica*, Vol. III, p. 288.
 Watt, G. (1892). *Dictionary of the Economic Products of India*, Vol. VI, part I, p. 194.
 Hooker, J. D. (1879). *Flora of British India*, 2: 203.
 Cooke, T. (1903). *Flora of the Presidency of Bombay*, Taylor and Francis, Red Lion Court, Fleet Street, London, 1: 377.
 Gamble, J. S. (1918). *The Flora of the Presidency of Madras*, Adlard and Son, Ltd., London, Part II, p. 363.
 Sampson, H. C. (1936). *Bulletin of Miscellaneous Information*, Additional Series 12: 138.

Better Methods of Virginia Tobacco Cultivation in Guntur District.

By M. NARASIMHAM, B. Sc. (Ag.),
 Agricultural Demonstrator, Guntur.

Introduction. Guntur district has nearly 50 per cent of the total area under tobacco in the Madras Presidency. During the past decade, as a result of persistent efforts of the India Leaf Tobacco Development Company, ryots have taken to the cultivation of Virginia tobacco and its curing in barns. The present area extends over 1,00,000 acres, fetching a return of about one and a half to two crores of rupees to the District. The country tobacco (*Natu*) extends over 50,000 acres and it may fetch another half a crore of rupees every year. Virginia tobacco produced in the Guntur, Kistna and Godavari districts forms the bulk of cigarette tobacco produced in and exported from India. With proper methods of cultivation, curing and marketing, it is possible in course of time to find a steady market in foreign countries especially the United Kingdom for a fairly large quantity of high class leaf. Tobacco is at present the main 'money' crop in the Guntur district. Unless proper steps are taken to increase the yield and improve the quality of leaf and rectify some defects in other respects, the trade is likely to suffer in a very short time.

Importance of quality in Virginia Tobacco. Unlike other agricultural commodities, and other varieties of tobacco, the range of variation in the market price of Virginia tobacco, even on the same day, may be anywhere between Rs. 10 to 350 per candy of 500 lb. This indicates the extent to which quality is important in the crop. The aim of the Virginia tobacco grower should therefore be to produce as much of the high grade leaf as

possible to obtain better prices. There is always a good demand for high grade leaf for which remunerative prices are offered. Steps are being taken in several parts of India to introduce and extend Virginia tobacco cultivation. Naturally Guntur District is not likely to enjoy its present monopoly of trade in the commodity for long. It has also to be mentioned that of late Virginia tobacco in the district has been showing considerable deterioration in quality and yield. On the other hand, it is behaving very well in some of the new areas along the Godavari and Kistna rivers. There is no doubt that Guntur District is one of the few places where good quality Virginia tobacco can be grown. Hence Guntur can still hold the monopoly if the ryots adopt judicious methods of cultivation and curing.

Factors affecting quality in Virginia tobacco. The quality of cured leaf obtained is determined by a number of factors and they can be classified as follows:—

(a) Field factors affecting the development of proper colour of the leaf.

1. Rainfall before and after-planting.
2. Nature of previous crop.
3. Manuring given to previous and present crop.
4. Time of planting.
5. Nature of seeds and seedling used.
6. Season during the growth of the crop.
7. Prevalence of pests and diseases.

(b) Those affecting quality of cured leaf during the process of harvest and curing:—

1. Method of harvest and handling.
2. Method of curing and handling.

A good quality leaf from the barn is obtained from the crop that matures uniformly and produces the right type of yellowish green colour in the leaf. Factors such as rainfall are beyond control while the occurrence of pests and diseases are to some extent within human control. Proper methods of rotation and manuring, timely planting and use of pure seed can, however, be followed and they go a long way in producing the proper type of crop.

To produce good colour in the green leaf, Virginia tobacco should not be grown on the same field more often than once in three years. It should not be preceded by a pulse crop or groundnut or by a heavily manured crop like chillies. Cereal crops have been found to be the best to precede it. If for any reason, cereals could not be sown in the preceding *pyru* season, the land should be manured liberally and a fodder crop (e.g. maize) raised from June to August. Tobacco can be planted in October–November following. In brief, the preceding crop should not be one that will enrich the soil in its nitrogen content, a factor that will adversely affect proper colour development in the green leaf. Raising tobacco too frequently on the same land will favour the spread of pests and diseases. Early planting (September end or early October) is good, wherever it is possible. Planting later than 15th December is undesirable as the resulting crop will not ordinarily develop sufficiently good colour in the green

leaf. Late planting is however, inevitable in some of the low-lying Badava lands. It is inadvisable to attempt planting in heavy deep clays, where good drainage facilities do not exist, before the cessation of the North East monsoon.

The most important factors adversely affecting the quality of leaf during harvest and curing are the following.

- i. Entrusting harvest and subsequent processes on a contract basis to a gang of coolies who pick over-ripe, ripe and un-ripe leaf together, and cause mechanical injury to the leaf while handling it.
- ii. The absence of sufficient attention paid to curing methods by curers who do not understand the technique of curing.
- iii. Curing over-ripe, ripe and under-ripe leaves together.
- iv. Over loading the barn.

Contract work is usually cheap but the results are usually disappointing. If the grower exercises sufficient supervision in the field and at the barn, several defects of the system can be remedied. As bad curing results in loss of 25 to 50 percent of his profits, the grower would do well to learn the art and do the barn curing himself. If he cannot do it, he should at least procure the services of a good curer. It does not pay the cost of coal to cure the over-ripe, ripe and under-ripe leaves together.

Since two to four leaves are picked from each plant at a time, the stage of maturity will be different in each case. Experience has shown that it is desirable to sort out the green leaf after picking, into three grades with reference to their colour, and string them separately. The darker green leaf should be arranged on the top tiers the light green on the middle tiers and yellowish leaf on the lower tiers. The higher temperature near the flue pipes gives the over-ripe leaf at the bottom a chance to cure without sponging and the lower temperature and higher humidity at the top allows the dark green leaf to turn yellow quicker. The usual blemishes noticed in the cured leaf such as 'sponging', scalding, etc. are the result of over loading the barn with too many sticks, a habit which is almost universal in the district at present. By overloading, the leaves are kept so closely packed that some of them will not have the full benefit of the changes in temperature, aeration, and humidity to the right extent and at the right time, with the result, that they fail to cure well. The more thinly the barn is loaded the greater the possibility for the uniform distribution of temperature, aeration and humidity in it. What is gained in fuel consumption by loading the barn with too many sticks is many times lost in the quality of the cured leaf produced. It is known by experience that about 600 sticks in the small barn (16' x 16') and about 1,000 sticks in the big barn (20' x 20') form the optimum load.

How best to manure a Virginia tobacco crop:—To maintain yields at a high level and produce the right quality of leaf, it is necessary to manure the land in the right manner. Cattle manure in sufficient quantities is not available in the tract as nearly 15 to 20 acres are cultivated with a pair of

cattle. Heavy application of cattle manure induces rank growth resulting in dark-coloured coarse leaf which is totally unfit for barn curing. The use of artificials to supplement the natural resources of the soil, is therefore inevitable.

Nitrogen, besides assisting in the vigour and growth of the plant, has a marked effect on the quality of leaf. If a soil is poor particularly in its nitrogen content, light coloured leaf is produced. If heavy doses are applied, big coarse, dark green leaf with prominent veins is produced. Cattle manure, oil cakes, ammonium sulphate and nitrate of soda are the usual nitrogenous manures now available in the market. Black soils are generally rich in total potash. But as tobacco is a heavy potash feeder (about 100--150 lb. K_2O per acre per year) and as potash influences the quality of leaf, the available potash in the soil will not fully meet the needs of the crop. Potash manuring is, therefore quite essential for tobacco. A crop suffering from potash starvation will have in the leaves yellow mottling, dead specks, hobby-surface, and downward incurving tips and margins when green. Such leaves become harsh, dry, short and non-elastic when dry. Potash helps to improve the elasticity and fire-holding capacity of the leaf. The sugar content of the leaf is also improved by the supply of potash. Potash in the soil is rendered available by the application of sodium nitrate; hence the suggestion to apply half the quantity of nitrogen to the crop in the form of sodium nitrate. But there is the inevitable danger of leaching out of the nitrate during the heavy rains of September--October. Too much of potash turns the ash dark in colour. Phosphoric acid assists in an early, healthy development of the crop, regulates the rank growth resulting from too much nitrogenous manuring, hastens maturity and aids in the general colour development of the leaf. Lime is not required in Guntur soils which are rich in lime. If the burning quality (closer burn) is bad and snow-white ash does not result, about 30 lb. of magnesia (MgO) with half of it at least in a soluble form, may be applied. Tobacco does not ordinarily require any other manure to improve its yield or quality. Under Guntur conditions manuring with about 10 to 15 lb. of nitrogen (N), 50 lb. of potash (K_2O) and 75 lb. of phosphoric acid (P_2O_5) in addition to 5 to 10 cart loads of cattle manure per acre appears to be the best dose. Some fields do grow good tobacco without manuring or with moderate quantities of cattle manure in certain seasons, but to ensure good crops of tobacco over a length of time, systematic manuring with artificials is inevitable.

Nitrogen is best applied as ammonium sulphate and sodium nitrate in equal proportions (25 to 40 lb. ammonium sulphate and 32 to 48 lb. sodium nitrate per acre), potash as sulphate (100 lb. potassium sulphate per acre), and phosphoric acid as superphosphate (450 lb. super-phosphate per acre.) Undue proportion of chlorides should not be included in the mixture. The above suggestions are based on the general conclusions drawn from the elaborate manurial trials carried out over a number of years and on various soil types by the Indian Leaf Tobacco Development Company. Cattle manure

is best ploughed in, in June--July and artificials drilled deep into the soil or spread over the soil and ploughed in whenever there is a dry spell of weather in August, i.e., about two months before planting. As tobacco is grown dry in the district, the full benefit of the manure applied cannot always be realised by the crop unless it is applied sufficiently early and well worked in.

Use of pure seed. The purity of the Virginia tobacco crop in the district is poor. With a mixture of indifferent types the cured leaf will be far from satisfactory. The use of pure seed (strain) of proved merit results in a pure crop that matures uniformly. This will naturally result in the production of a uniform quality of cured leaf. In no other crop does purity of seed play such an important part. Harrison special No. 9 has been found to meet the needs of the district very well. It grows well, has an ideal type of leaf and develops good colour wherever conditions are favourable. Growers have reported favourably on it. As compared to the local bulk crop, it fetches them an additional net profit of at least Rs. 25 per candy of produce. As natural crossing is very common in this crop, it is not always the most vigorous plant in the field that is to be selected for seed purposes. Till a uniformly good quality crop is raised by the ryots themselves, they are advised to purchase their seed requirements from the Agricultural Department which is multiplying the pure strain and doing its best to supply as much seed as is wanted by growers. The price of pure seed required for an acre of planted crop is about 2 annas.

Raising nurseries. Under the present system of cultivation in the District, though a few big ryots raise their own nurseries nearly 75 per cent of the ryot population entirely depend upon the nurseries raised by professional nurserymen. This system has come into vogue for the following two reasons:—

- i. Nurseries have of late been failing in black soils due to 'damping off'.
- ii. It is easier to raise nurseries with success in the sandy soils and light loams, and seedlings from these areas establish much better on planting than those from the black soils.

In Guntur virginia tobacco cultivation has, assumed such industrial proportions that the man that raises the nursery is very often different from the man that grows the crop, the grower different from the barn owner and the barn owner different from the grader and ultimate exporter. As purity in the seedling stage cannot be easily determined, nursery men take little care to use good seed and the man that grows the crop pays the penalty. It is therefore highly desirable that this system vanishes soon. The next question is how best to raise tobacco nurseries in black soils. Before the I. L. T. D. company thought of raising nurseries in the Chirala sands, ryots were raising their nurseries in the black soils with success. Similarly, on the Agricultural Research Station, Guntur representing a typical black soil area, good nurseries are being raised with success from year to year. It

should therefore be possible to raise nurseries in the black soil areas. To achieve success, the present methods of raising the nurseries have to be modified and any amount of attention bestowed on them will not be too much.

The following are some of the precautions to be taken while raising a tobacco nursery in heavy black soils:—

- i. Select a fertile high level sloping bit of land with a good water source such as a tank, pond or well near at hand. The water table should be so low that there is absolutely no danger of inundation or bad drainage even in the heavy rainy season.
- ii. Make the soil permanently lighter in physical texture by heavy application of sand, tank silt, organic matter and green manures, to facilitate free movement of soil moisture.
- iii. If chillies are also raised, tobacco and chilli nurseries should be rotated. Otherwise two plots may be set apart for tobacco and alternately put under tobacco nurseries.
- iv. Plough the land deep after the North-east monsoon and apply liberal quantities of cattle manure, in December or January, and work it in well with the plough.
- v. Keep the land ploughed till the end of July, then lay it out into a number of long beds, 4 to 5 feet wide with intervening drainage channels $2\frac{1}{2}$ feet wide and about 1 foot deep ranged along the slope, to facilitate quick drainage. The length of the bed may be adjusted with reference to the length of the field and general slope of the land. Periodical deepening of the drains and earthing up of the beds are necessary.
- vi. About 3 or 4 days before the proposed date of sowing the seed beds, a fairly thick layer of cotton or red gram stalks, rejected cattle fodder, organic rubbish from the manure heap, casuarina twig, groundnut husk or any other type of inflammable materials may be spread evenly over the prepared nursery beds, and set fire to. This is best done when the air is still. The beds should not be burnt when they are too wet. As soon as the bed cools down, the ashes may be worked in lightly with hand hoes to prevent the ashes being blown off by wind. The beds are then ready for sowing.

Sowing of nursery beds. The beds should be raked and moderately compacted. The seed should be mixed with plenty of ashes or fine sand and sown evenly, distributing it twice across at the rate of about an ounce for a bed measuring 5' x 150'. An ounce of tobacco seed contains about 3 lakhs of seed. The seed should then be pressed down with the palm of the hand or a flat board after sprinkling a small quantity of burnt soil on the sown beds. The practise of applying powdered cattle or sheep manure to cover the seed or just before sowing seed should be avoided as the manure may infect the soil with the fungus responsible for "damping off."

Watering should be done with a fine rose-can till the plants are about three weeks old. Covering seed beds temporarily with casuarina leaf in the stages as adopted by ryots at Bapalla and Chirala is a practice worth copying.

After-care of nursery beds. Burning is done to kill the soil organisms that cause 'damping off' 'blackleg' etc. With favorable weather conditions, they attack the seedlings causing rotting and death. To avoid these diseases the beds should be thinned out wherever they are too thick. The thinnings may be planted where the beds are patchy. Such transplanting can be done when the leaves are the size of an eight-anna piece. If the seedlings are very weak or fail to do reasonable progress, sulphate of ammonia may be applied at about 5 to 20 ounces for a bed of 250 feet length. Similarly when the rate of growth of the seedlings has to be accelerated after a batch of seedlings are removed from the bed, the bed may be manured at 12 to 25 ounces ammonium sulphate for a bed of 250 feet length. The manure should be well mixed with sand or dry earth and evenly broadcast over the beds in the evenings. A heavy watering should be immediately given. As the plants usually become soft after manuring, at least a week should elapse before the seedlings so manured are ready for planting out. Even when manure is not applied, the seedlings should similarly harden off for about a week before being pulled out for planting. If proper attention is paid to weeding, control of insect and fungus attacks, an acre of nursery is expected to supply seedlings sufficient to plant at least 100 acres with sufficient seedlings left over for subsequent filling up of gaps. In black soils, many more seedlings can be expected from a unit area of nursery. At present ryots use at least five times the above seed rate to start with and sow more seed in the same beds whenever the nursery looks thin according to their own standard of judgment. In the majority of cases it is this thick sowing that encourages 'damping off'. In addition to the precautions mentioned above a weekly spraying with a suitable spray mixture containing a fungicide and insecticide is found useful to protect the tiny seedlings from diseases and pests. The first spraying should be done when the leaves of seedlings have attained the size of one's finger nail. The following mixture is recommended for the first two sprayings:—

Bouisol (Colloidal copper)	1 oz.
Lead arsenate	$\frac{1}{2}$ oz.
Agral (spreader)	$\frac{1}{8}$ oz.
Water	1 gallon.

If the above preparations are difficult to obtain, the following spray mixture is recommended.

Copper sulphate	1 lb.
Quick lime	1 lb.
Lead arsenate	4 oz.
Water	$12\frac{1}{2}$ gallons.

The copper sulphate is dissolved in half the quantity of water. The quick lime is first slaked and then diluted in the remaining quantity of water. Copper sulphate solution is added to the lime solution slowly and the

SELECTED ARTICLE

The Man and the Plant.

By H. MARTIN LEAKE, M. A. Sc. D.

Cotton is an agricultural product, the product of a plant. Its successful cultivation, apart from the financial aspect, must conform, therefore, to two requirements. It must provide both the essential conditions which the nature of the plant as a living organism demands, and the amenities which man, the producer, considers to be his due in return for his labour. But there is no question as to which of the two is the more fundamental. The plant is a sentient, but unreasoning, organism and can only adapt itself to a certain limited range of conditions. The plants in a greenhouse will die if left unwatered for a few days, and the fact that the absence which led to the neglect was unavoidable makes no difference to the result. With man the vital limits are wider, but he is reasoning and not only can, but does, voice his protests in strikes and rioting long before those limits are reached. This is a simple proposition, but one which is frequently over-looked. The human requirements, for the above reasons, assume primary importance, and the protest of the plant, because silent, is ignored.

The cotton plant, with plants in general, will only grow and mature its crop if the conditions, and the environment, fall within somewhat narrow limits. If rainfall, air-humidity temperature and so on transgress those limits failure will result, though a time-factor is here involved, the extent of the injury being dependent on the duration of the exposure to the adverse factors. These are physical factors, but they are not the only ones concerned; there are the factors concerned with the soil, also physical factors, and there are biological factors which influence the resultant crop, weeds, pests and diseases. It is the function of the successful farmer so to alter the conditions existing in nature that the plant throughout its growth is never subjected to such adverse conditions. It is an objective rarely, if ever, attained, for man has only a partial control over the environment. Herein lies the fundamental difference between agriculture and industry; in the latter the conditions are under the full control of man.

There is here implied the dependence of success on a detailed knowledge of the particular plant in relation to the environment, and it is a knowledge which is of two kinds. There is the knowledge which concerns the effect of those rapidly changing factors of the environment, seasonal and daily changes of temperature and rainfall. These require immediate action, and decisions must be taken by the cultivator himself. It is knowledge which has been acquired by farmers of all races in greater or less degree as the result of the accumulated experience of generations. Then there is the knowledge of those more subtle influences, nature of the soil, availability of plant food, the life-histories of the various pests and diseases and so on the prerequisite to control. These are technical matters which require and technical investigation to acquire and technical control to apply.

It is possible, in view of the above, to define agriculture as the adjustment of the environment to the needs of the particular crop plant—perhaps the most comprehensive definition of agriculture there is. To achieve this objective three conditions must be fulfilled: skill on the part of the farmer, technical investigation, and technical control. But there is a further aspect. Where the conditions of growth cannot be sufficiently altered to suit the plant's requirements, it is often possible by careful search to find a variety which is capable of

growing under the conditions better than any variety hitherto available. It is a process which has been adopted instinctively through the ages and now forms a standard operation with all crops and in most countries. But recent advances in the technique of plant breeding have gone further and rendered it possible within reasonable limits to manufacture varieties having the desired adaptability. This alteration of the plant to suit the environment is a matter for technical investigation.

The human aspect demands requisites of a different nature. Primitive man is readily satisfied with little more than the absolute essentials to maintain life—food and a minimum of clothing and shelter—but this simple state is everywhere passing. With the spread of education, news and facilities generally, he is coming to recognize that there is a more desirable standard of life only to be obtained by the retention of a larger fraction of the return for his labour. It is inevitable that this urge on his part should bring him into conflict with those who handle his product and are naturally unwilling to reduce their share of the return. As long as demand exceeded supply, as was the case until early in the present century, this adjustment could be made within reason at the expense of the ultimate consumer. But latterly, contracted demand and vast increases in supply have made adjustment by such a method impossible. These forces have created conditions under which not merely has it become impossible to maintain the progressive advance in labour's material welfare, but actual reduction has taken place in some cases. If the labour force engaged in tropical agricultural production has progressed in its views as to what constitutes an adequate standard of life, it has not yet learned to appreciate the influences at work which restrict its capacity for earning. Herein lies the stimulus to the unrest which has characterized recent years not only in the tropical colonies but in general. Being capable of action labour has protested, and often with violence, and the strength of the movement of unrest has forced the human needs to the front, not infrequently to the neglect of the more fundamental needs of the silent plant. In practical life it is no answer to say, if the facts do not fit the theory, so much the worse for the facts. Facts have an unpleasant way of asserting themselves, and a solution to the present problem can only be sought through a recognition of its dual nature. The solution will not be found through argument on *a priori* grounds; the best approach will be made through a review of the various systems found in action and of the response in each to the changed conditions.

During the early history of colonial development a purely nationalistic view prevailed: development in the interests of the mother country, as witness the Navigation Laws and the slave trade. Under it arose a system of plantations worked by a labour force composed originally of slaves, and later of wage-earners often bound by indentures. With the passing of slavery a change of sentiment took place which slowly crystallized out at the end of last century in the dictum of Chamberlain: the trust for civilization. Under it, the insistent demand for the products of tropical agriculture was to be met by extended cultivation by the readiest means then available. The plantation system, and the results are to be traced in the encouragement of settlement in Kenya. It was assumed rather than argued that from such a development benefits must automatically accrue to the native populations. Another quarter of a century elapsed and a further change of sentiment took place which crystallizes out in Article 22 of the Covenant, the trust of the native populations. It was a sentiment which Lord Lugard, with a prescience in advance of his time, anticipated by a quarter of a century. The readiest means were discarded and the more leisurely method through a system of peasant proprietors was adopted, and this course was facilitated by the lessening urgency of the demand owing to the large extension in the sources of supply. It takes note of the human aspect and again, it was

assumed rather than argued that the needs of the plant would be adequately met. These two systems, existing as they do in a vast variety of forms and admixture, have supplied in the past practically the whole of the agricultural produce which comes on to the markets from the tropics. The test of their efficiency to supply both requisites is found in the reactions during the recent years of depression.

The plantation system, in its highest development, that of the large company-owned estate, is in a position to provide the requirements of the plant, financial considerations alone determining whether adequate provision will be made or not. Irrigation works will be provided if the extra return is calculated to cover the cost and processing plants to handle the product can be erected, technical control and even investigation can be organized, while much of the details of cultivation will be directed by a system of supervision which leaves little or nothing to the initiative of the manual worker. The latter is paid a fixed wage, and his interest in the success of the undertaking is limited to the chances of loss or continuance of employment. The small plantation has a lesser capacity owing to the limited financial status of the owner, but the position of the manual worker is the same in both cases.

The demand of labour here takes the form of a claim for an increased wage, which carrying as it does no incentive to greater energy, is naturally resisted as far as possible. The outcome will depend on the relative bargaining power, and where labour has an alternative field of employment, it is the employer who must yield; where no such alternative exists, it is labour that must do so. But the matter does not end there. The general reaction, especially where labour is in the stronger bargaining position, is for a resort to be made to mechanization with higher wages to a restricted labour force. It is a solution which has answered well where there is a shortage of labour as in Hawaii, but one which has had bad repercussions where there is an excess of labour as in some West Indian Islands, and notably in Cuba and Puerto Rico. There is no general solution to the problem here.

The peasant system appears to accord more closely with the human requirements, for the peasant is his own master and works for his own benefit. But agriculture among a tropical peasantry is not founded on accumulated experience and there are not the financial resources for technical control and, still less for technical investigation. Processing plants, which may be necessary if the product is to be acceptable in a competitive market, cannot be erected, and the regulation of a seed supply, so important an item in the case of a naturally cross-fertilized crop raised annually from seed such as cotton, is impossible. Some outside agency has to provide these, and in most cases that agency is Government. There has thus arisen in place of the independent employer—between whom and labour Government could, in the last resort, arbitrate—an authoritarian control against which there is, and can be, no appeal, for the final arbiter, Government, is one partner. It is only necessary to read the cotton Ordinances, Rules and Regulations of Uganda or the like enactments of Zanzibar for cloves and copra to appreciate the extent of the arbitrary power which Government has been forced to take to itself. This is the inevitable result of the failure to realize that an uncontrolled peasantry cannot "deliver the goods". Herein lies the cause of the unrest which has in recent years characterized countries with a peasant system no less than those with a plantation system. Government, it is true has drifted rather than deliberately entered into this position, and is seeking a way out through co-operation. On the question of co-operation it can here only be said that, for success on the scale needed a standard of education is required which can hardly be attained in the time which the urgency of the problem demands.

It is not possible in the brief space at command to argue the intermediate positions of a mixed peasantry and plantation in all their varieties. Such a case is found in Mauritius, and unrest has not passed by that island. It is difficult, therefore, to avoid the conclusion that the dual requirements are provided neither by the plantation nor the peasant system alone or in combination. The remedy must be sought elsewhere.


There are fortunately in operation two developments which promise to provide the solution, one in Fiji and one in the Sudan and if the first is here taken as an example, it is because it gives in many respects a clearer picture of the principles and is less well known to cotton interests. The Fijian sugar industry in the past was worked on the plantation system with labour under indenture from India. Unrest was rife and kept within bounds only because settlement on the extensive tracts of waste fertile land offered an alternative field of securing a living, and because fresh supplies of labour were readily recruited in India. These conditions were brought to an abrupt end in 1917 when the indenture system was abolished, and an acute labour shortage followed. The solution of the deadlock was found in the subdivision of the large estates into small holdings leased to the tenantry, and the organization of a technical supervisory service of which the guiding spirit is co-operation with the tenantry. The result has been so successful from the industry's side that it places its sugar on the world's market with profit, and from the labourers' side in that an intense demand for tenancies has arisen and the whole of the area controlled with the exception of certain small portions retained for trial purposes, is now so subdivided. Here, both requisites are met and it is along such lines that the general solution of the present troubles must be sought. *The Empire Cotton Growing Review*, (16: 180-185.)

ABSTRACTS.

Soya Beans.—Arnold, H. C.—*Rhodesian Agricultural Journal* 38, (1940). 588—606.

The introduction of strains of soybeans which yield good crops of seed and also retain their seed in their pods long enough to allow a reasonable period for reaping, appear to make the cultivation of this crop economically possible. The choice of a variety suited to the district is of the utmost importance. Potchefstroom No. 184 is the best yellow-seeded and the Jubiltan strains are the best fodder varieties, but seed supplies of these are at present very limited. Rainfall and soil conditions suited to maize are the best for soya beans also. Seed requirements are 40 to 50 lb. of the large seeded and 15 to 20 lb. of the small-seeded kinds, per acre. Twenty-four inches between rows and 3 to 4 inches in the rows may be considered as standard spacing for the yellow-seeded kinds, but the distance between rows may vary from 8 inches to 30 inches if compensating differences are made in the spacings between plants in the rows. Depth of planting should be between one and two inches and on no account exceed three inches. Bacterial inoculation of the seed is advisable when the land is sown for the first time. Early and thorough weeding is imperative. Harvesting may commence when the seed reaches the hard dough stage. The stalks should be tied in bundles and stooked as soon as they are dry enough. Three to four bags per acre could be expected from fairly well attended farm crops, though twice that amount has been obtained from well manured land.

Its high protein content makes it especially valuable for balancing the starchy nature, and it can be used as a substitute for animal protein in the diet of both Europeans and natives. The valuable nutrients, protein and oil, can be transported more economically as soya beans than is possible in any form of either maize, wheat, groundnuts or beef. Although the costs of growing would be higher, the marketing costs would be lower than those for maize.



Liebig's Mineral Theory of Plant Nutrition—Acharya C. N. Indian Farming 1: (1940).

It is now exactly a hundred years since Liebig propounded his mineral theory of plant nutrition, which stated that plants need be supplied with minerals only in order to provide for normal growth. It synchronised with the change-over from an era of natural or organic manures to a new era, which believes in the potency of artificial fertilizers, properly used. In the early stages of the enthusiasm for artificials, the case for natural organic manures which were in use through thousands of years was overlooked—with disastrous results in several cases by way of a rapid deterioration in the physical condition and fertility level of the soil. The manurial pendulum has now swung to a middle position between the age-long practice of using organic manures alone and the immediate post-Liebig belief in the sufficiency of artificials alone. Present-day practice aims at combining the advantages of both systems by returning to the soil as much of farmyard manure, composts and other natural manures as can be obtained and using artificials as supplements in order to obtain peak yields. The relative amounts of organic and inorganic manures that can be used with advantage for any crop will depend on a number of factors such as the nature of the soil, the nature of the crop and relative prices of the two classes of manure on the farm. Recent work, both on the agronomic and on the scientific sides, has shown the essential limitations of the purely mineral hypothesis put forward by Liebig for the nutrition of plants. Controlled experiments in solution cultures, containing only mineral salts, have shown that though plants can be grown for one or two generations in such media, they deteriorate rapidly if grown through a number of generations with seed obtained from the previous mineral solution cultures. This has been found to be due to the lack of essential phyto-hormones which are carried in small amounts in the normal seed grown in the soil and are found in abundance in organic manures and also in soil organic matter. In the temperate and colder climates, where the soil contains a good percentage of organic matter (5 to 10 per cent) and the rate of oxidation is slow, the return of the root systems of crops may be sufficient to maintain the soil in good physical condition; and continuously high yields may be obtained for decades together by the application of artificials alone, as evidenced by the results of the permanent wheat plots at Broadbalk (Rothamsted). But the case is different in the tropics, where the soil contains barely 1 to 2% of organic matter and the rate of oxidation of added organic matter is high. The return to the soil of the root-system alone is insufficient, under tropical conditions, to maintain the soil in proper condition. For this purpose additional quantities of bulky organic manures in the form of farmyard manure, composts or green manures, have to be added almost every year. Already, the system of using 'straight' artificials is yielding place to an alternation or conjoint use of organic and inorganic manures.

M. K.

Feeding of Milking Stock. Kartha K. P. Indian Farming 1: (1940).

Cattle improvement in India is as much a matter of feeding and management as breeding. While better breeding may produce an improvement of 10 to 12 per cent in milk yield from generation to generation, better feeding and management alone can be relied upon to produce an immediate increase of 50% in ordinary Indian cattle. Feeding of milch cows in villages is done haphazard without any regard to the requirements of the animals. In many cases, cattle are seriously underfed while in others there is considerable overfeeding, both of which affect milk efficiency. The object of giving food to milch animals is to provide the energy required for performing the various functions. In the feeding of animals, therefore, care should be taken to see that (1) you provide sufficient food for the animal to perform its vital functions viz. the 'maintenance ration', (2) you provide extra food for the production of milk viz. 'the production ration'.

and (3) that the quality of the food is such that it contains enough proteins, minerals and vitamin viz. 'a balanced ration'. The quantity for maintenance will depend upon the live weight of an animal and that for production on the amount of milk produced. The actual quantity will also depend upon the quality of the feed. If good fodder is available, a mixture of green and dry fodder fed *ad lib* should provide the requirements for maintenance. If fodder is of poor quality, the maintenance ration should contain in addition 5 lbs. concentrates in the case of buffaloes and 3 lbs. in the case of cows. Production ration is supplied in the form of concentrates like gram, oil-cakes and bran and the optimum requirement is one lb. of a suitable mixture of concentrates for every 3 lbs. of milk yielded.

The following are a few typical mixtures found suitable for milch cattle.

Mixture.	Bran.	Crushed gram.	Crushed barley.	Crushed cotton seed.	Crushed groundnut cake.
1	4	1	1	—	2
2	4	—	2	—	2
3	4	1	1	2	—
4	5	1	—	—	2
5	4	2	—	2	—
6	4	1	2	—	1
7	3	—	2	3	—
8	3	1	2	2	—
9	5	—	1	—	2

M. K.

The control of Nut grass in the Sudan Gezira. F. W. Andrews. Empire Journal of Experimental Agriculture.

Nut grass (*Cyperus rotundus* L.) is reported to be a troublesome weed in the irrigated cotton land of Gezira—a flat plain lying between the Blue and the White Niles. It was found that the eradication of this weed was fairly accomplished if a root cutting implement (corresponding to blade harrow or *Guntaka*) was drawn through the soil at a depth below that of the lowest tuber and the disturbed soil containing the tubers left in a dry condition. A good many of the tubers were about six to nine inches below the ground surface. Repeated cultivation of the land may eventually destroy this weed.

M. K. R.

Rice Straw and Molasses for Range Cattle R. C. Macahilig Philip. Agriculturist, (1939), 28, 561.

In view of the efforts to widen the basis of an industry hitherto largely dependent on monoculture and to find an outlet for excess molasses these experiments on the use of rice straw and molasses as supplements for grazing cattle will be of interest. During the dry season (112 days) the average increase in weight was 29.43 kg for cattle with supplementary feed against 1.43 kg for those fed on pasture alone. This was definitely economic. During the wet season (196 days) the respective increases were 19.93 and 16.45 kg., a difference which was uneconomic. *International Sugar Journal*, 42 (1940) 256.

Manufacture of "Leesa" Soft Sugar in India. S. D. Agnihotri Proceedings of the 6th Convention of the Sugar Technologists' Association of India, pp. 435-486. There are several factories in Bombay producing a soft sugar, e. g., *Leesa* made by crushing refined beet or Java special crystals and mixing it with 9 to 15 per cent of 5A corn glucose (35° Be.) of the Corn Products Co. (India). *Leesa* so made forms hard lumps on keeping, and invert sugar gives a product which is better in feel and in keeping quality. Japanese *Leesa* appears to contain a fat, which the author was unable to identify. No modern sugar factory in India has undertaken to produce *Leesa*, though from the prices ruling for that imported from Japan the margin is quite alluring. (*International Sugar Journal*, 42: (1940) 253).

Gleanings.

Healthier Plants. From the Horticultural Department of the American Chemical Paint Company comes news of the commercial production of a vitamin hormone stimulant, *Transplantone*, for plants, that not only invigorates old roots but also multiplies the production of new ones, reduces the loss which frequently occurs with transplanting operations, and reduces wilting. It is applied to rooted plants to add to existing root growth and to force their general growth. *Transplantone* is a water-soluble powder impregnated with vitamin B₁ and other parts of vitamin B₁ fraction, plus root-promoting hormones. The hormone initiates root growth and plant physiologists assert that the Vitamin B chemicals are necessary for the maintenance of their growth. That it is quite concentrated is obvious for it requires only one level teaspoonful to a gallon of water to make a stock solution which is then further diluted. Seedlings may be lightly sprinkled weekly, or it may be applied to plants set out in the soil, whether they be trees, shrubs, vines, annuals, or perennials. In the case of plants which are set out without a ball of earth, the manufacturer recommends that the roots be soaked in the stock solution for an hour. Treatment usually results in vigorous and extensive root growth and this, in turn, requires more frequent watering than is ordinarily necessary. The manufacturer further claims that, owing to frequent clipping, grass is unable to produce enough vitamin and hormone naturally for the roots and that watering with an ounce of stock solution to three quarts of water will improve turf quality. Sod's similarly treated before being set in place will also readily form new roots. (*Science and Culture* 6: 219.)

Indication of a New Chick Growth Factor in Rice. Hogan and his co-workers (*Jour. Biol. Chem.*, 64, 113, 1925) suggested that polished rice contains a chick growth factor not present in any significant amount in yeast. MacFarlane et al (*Jour. Nutrition*, 4, 331, 1931) supported this observation. Stokstad and Manning (*Poultry Science*, 18, 413, 1939) studied this problem and found that polished rice has a growth-promoting effect which cannot be provided by levels of dried yeast up to 15 per cent of the ration. Very recently Almquist, Stokstad, Mecchi and Manning (*Jour. Biol. Chem.*, 134, 213, 1940) have found that glycine is required in the diet for optimum growth of the chick. When adequate glycine is present in the diet chondroitin has a growth-promoting action on the chick. A combined effect of glycine and chondroitin can replace the "rice factor".

(*Science and Culture*, 6: 239.)

Correspondence.

To The Editor, Madras Agricultural Journal.

One-cent paddy nurseries.

Sir,

I enclose two photographs of one cent paddy nurseries the value of which has been demonstrated to ryots and which was a feature of the landscape on the main bus route in the Mayavaram to Tiruvarur Road between June and September 1940. These are representative of the 4574 one cent nursery plots laid in 5676 acres of nursery raised during the period.

After the last ploughing the nursery fields were levelled by a levelling board drawn by a pair of bullocks, the fields marked out into plots of 8 feet breadth and 54 feet length with channels one and half feet wide all round. The plots thus marked out were again levelled with a hand levelling board, the coolies



ONE-CENT PADDY NURSERIES

Top. A view of adjacent plots showing the channels on all sides.

Bottom. Another view showing the uniformly grown seedlings ready for transplanting.

working from the intervening channels. Water was then let in to about an inch depth and allowed to stagnate in the beds for a few hours. The sprouted seed was then measured out at 1 Madras measure per plot for *samba*, $1\frac{1}{2}$ M. M.* for *Kuruvai* and 2 M. M. for *Udukuruvai* and handed over to the sowing coolies who went round the plot twice or thrice along the intervening channels scattering the seeds uniformly.

This method of sowing ensured uniform and good germination of seeds and uniform growth of seedlings as there was good drainage. The method also prevented the formation of foot print pockets caused by labourers walking over the beds which is unavoidable in the local method. The seedlings were transplanted in singles 4 inches apart in *Kuruvai* and 4 to 5 inches apart in *Samba*. The *Kuruvai* yields have given an average increased yield of about 10% over the bunch planting apart from a saving of 50 per cent in seed.

Agricultural Demonstrator's Office,
Mayavaram.

Yours etc.,
M. J. David.

To The Editor, Madras Agricultural Journal.

A Novel device for pushing sale of Groundnuts.


Sir,

Groundnut has long been recognised as a very palatable and cheap "nut" though people seldom realised its high nutritive value. Roasted groundnuts are often sold exposed in shallow bamboo trays at street corners or in the vicinity of toddy shops, in most of the towns, by old women, but are patronised only by labourers or poorer class of people chiefly because they are sold under dirty conditions, or in uninviting surroundings. Other people go in for this tasty and cheap "munch" only at places of public resort, like parks, sea-beach, or theatres when it is sold by decent-looking mobile vendors. A new method of selling groundnuts now prevalent in the West Coast towns and which has come to my notice recently, is described below with the hope that it would be of interest to others particularly at this time when much is talked about the restriction of area under groundnuts and stimulation of internal consumption to meet the situation brought about by the closure of foreign markets and the consequent lack of export trade.

The new device is an adaptation of a contrivance for a game of chance to the sale of groundnuts, which easily provokes the crowd that gathers round it to try their luck. It consists of a wooden plank about $4\frac{1}{2}$ feet long and 6 inches broad, one edge of which is divided into a number of small spaces each marked with a number in the following order, viz., 25, 25, 40, 28, 31, 28, 24, 32, 28, 24, 26, 29, 27, 28, 32, 28, 50, 25, 34, 23, 29, 25, 28, 24, 32, 26, 25, 23, 31, 27, 28, 100. (Vide illustration). Over this plank a small toy car is flicked, the movement being confined to a straight path by providing at each end of the car, a vertical metal piece having a longitudinal slit through which passes a galvanised wire. The wire is drawn tight and fixed to two vertical struts at the two extreme ends of the plank. Three inches from each end two more struts are fixed and a narrow strip of rubber is tightly tied round them to act as springs for throwing back the toy car when it is flicked backwards or forwards. The toy car is provided with a pointer towards the side of the plank on which the numbers have been marked. The set is made locally and costs only about a rupee.

On receipt of the token money which is only a pie in this case, the vendor—generally a boy—who sits on the road side with a bag of roasted groundnut pods gives a flick to the toy car. The car moves forward, strikes against the strip of rubber, recoils back and finally comes to rest, its pointer standing over some

* M. M. means Madras measure = 3 lb.



A novel device for pushing sale of Groundnuts.

number painted on the plank. This number of roasted groundnuts is then counted out to the man who has had his chance. The idea of trial of luck being contagious spreads through the whole crowd that gathers round this novel contrivance. The vendor by this simple method is able to drive a brisk sale up to about Rs. 1/8 to Rs. 3 a day, and earn a profit of annas two for every rupee worth of groundnut sold. In a small panchayat like Pattambi in South Malabar about half a dozen such sales-boys are said to operate during the season. It has also been gathered that this method of selling groundnuts is prevalent in many other small towns in the West Coast and has spread even to the States of Cochin and Travancore. The method proves attractive and may be introduced elsewhere as well.

Agricultural Research Institute,
Lawley Road, P. O., Coimbatore,
Dated the 8th November 1940.

Yours &c.


C. M. John.

To The Editor, Madras Agricultural Journal.

A newly designed bullock-drawn levelling-board.

Sir,

In wet lands levelling is usually done for paddy nurseries to get good, even germination of paddy seeds. It is not generally done for transplant fields. It is specially so in Tanjore Delta and in all other districts where irrigation water is supplied to wet lands from big projects. Wherever there is a plentiful supply of irrigation water, ryots do not bother about levelling wet lands. Even if land should be uneven from one end to the other, they can let in water in plenty to reach the other end. But the case is different with dry districts like Salem, North Arcot etc., where water is baled out from deep wells to raise paddy crops. Ryots in these districts have necessarily to keep their land perfectly level so that they can easily, quickly and economically irrigate their crops. The levelling of wet land further helps even growth of plants, even flowering and maturing of grains, which must result in greater yield. For securing such levelling in wet land, a new levelling board was designed to be worked by cattle.



The implement (see illustration) consists of an ordinary level plank 5 feet long at the bottom, 4 feet at the top, 1 1/2 feet deep, and 1/2 to 3/4 inch thick, to which a shaft-pole 7 feet long is fitted about 3 inches from the top edge and at right angles to the plank at the centre. The two long edges of the plank are made straight and even. It is provided also with a wooden handle about 3 1/2 feet in height. The shaft-pole passes through the plank and through the handle at the back and firmly secured behind by a wooden peg. Two iron brackets are

provided binding the shaft-pole to the plank in order to give steadiness and even pull to the levelling board. Four holes about 1 1/2 inches in diameter are symmetrically-made on both sides of the plank about 3 to 4 inches from the top edge. This is to provide an outlet for excess water that may be collecting over the puddle when the implement is working as there is no need to drag the water also. It will give some relief to the cattle by lessening the strain.

The implement is to be worked in wet lands as the last operation before planting paddy and when the field is in a well-puddled condition. Wherever there is uneven ground which can be easily spotted out by having 1 to 2 inches of water all over the field, the implement is lightly pressed by the handle when it is working. The excess puddle will collect in front and is carried forward and where there is depression the implement is lifted by the handle gently and gradually and the mud will be dropped evenly in the depression as the implement moves. Where the land is already level the implement is allowed to move on without any strain on the handle. One pair of animals can easily cover upto 2 to 3 acres in half day according to the amount of levelling to be done.

If the levelling board is worked every year one can easily dispense with an annual operation in wet land which is called *Kulivattu* over which large sums of money are spent at least once in four or five years over the same piece of land. If the soil should be sandy or very sandy loam it is better the levelling is done with the implement previous to the last ploughing and planting. The idea is such soils may be hardened and it may prevent young roots from easily taking root and the succeeding ploughing will rectify and loosen the soil. This levelling board was tried by several ryots and found to do excellent work. There are now 30 to 40 of them working successfully in Shiyali Taluk.

Agricultural Demonstrator's Office }
Shiyali 6-11-40.

Yours etc.

M. Gopala Chetty.

Coffee Control Scheme.

We publish below a Press Note issued by the Government of Madras regarding the Coffee Control Scheme. Ed. M. A. J.

The coffee industry which is concentrated in Madras, Coorg, Mysore, Travancore and Cochin has lost a considerable proportion of its export market as a result of the war. At a meeting of the representatives of all coffee interests, held at Madras in September last, which was attended by the representatives of the Governments and States concerned and the Hon'ble the Commerce Member of the Government of India, an agreed scheme for dealing with the situation was arrived at.

The Central Government have decided to introduce that scheme, as an experimental measure, for the season 1940-41. The Coffee Market Expansion Ordinance, 1940, has been promulgated to give effect to that decision. It comes into effect from December 21, 1940. Sub-section 2 (1) of Section 14 of the Ordinance requires every person owning land planted with coffee plants aggregating not less than 25 acres, whether such land is comprised in one estate or in more than one estate and whether it is wholly or only partly in British India, to apply to the registering officer appointed in this behalf by the Provincial Government to be registered as an owner and in respect of each estate owned by him. Such application has to be made before the expiration of one month from 21st December 1940, the date of commencement of the Ordinance or before the expiration of one month from the date on which after the commencement of the Ordinance he becomes subject to the provisions of sub-section (1) of Section 14 of the Ordinance. The Ordinance empowers the Provincial Government to appoint the registering officer.

The Government have accordingly appointed the Collectors of districts in which the estates are situated as the registering officers in this Province. Every owner of a coffee estate to whom the Ordinance applies should apply for registration to the Collector of the district in which his estate or one of his estates is situated within the time limit prescribed in the Ordinance.

Crop and Trade Reports.

Statistics—Cotton—1940-41—Third forecast report The average of the areas under cotton in the Madras Province during the five years ending 1938-39 has represented 9.7 per cent of the total area under cotton in India.

2. The area under cotton up to 25th November 1940 is estimated at 1,874,300 acres. When compared with the area of 1,730,300 acres estimated for the corresponding period of last year it reveals an increase of 53 per cent.

The area is the same as that of last year in Tanjore and South Kanara. A decrease in area is revealed in Vizagapatnam, West Godavari, Kurnool, Nellore, South Arcot, North Arcot, Ramnad and Tinnevely and an increase in area in the other districts of the Province. The variations are marked in Kurnool (-34,000 acres), Bellary (+25,000 acres), Coimbatore (+69,300 acres), Madura (+32,500 acres), Ramnad (-22,200 acres) and Tinnevely (-17,500 acres).

The area under irrigated cotton, mainly Cambodia, is estimated at 226,300 acres as against 153,800 acres in the corresponding period of last year, thereby representing an increase of 47.1 per cent.

3. Pickings of the *mungari* or early sown crop in parts of the Deccan are in progress and the yield is expected to be slightly below normal.

Normal yields are expected in all the districts outside Tinnevely. The seasonal factor for the Province as a whole works out to 99 per cent of the average as against 100 per cent in the corresponding period in the previous year. On this basis, the total yield is estimated as 410,400 bales of 400 lb. lint as against 366,800 bales of last year, thereby representing an increase of 11.9 per cent. The crop is young and it is too early to estimate the yield with accuracy.

The estimated area and yield according to varieties are given below:—

(Area in hundreds of acres, i. e., 00 being omitted; yield in hundreds of bales of 400 lb. lint, i. e., 00 being omitted.)

Variety.	Area from 1st April to 25th November		Corresponding yield.	
	1940	1939	1940	1939
1	2	3	4	5
	Acres.	Acres.	Bales.	Bales.
Irrigated Cambodia	212.6	142.8	132.5	89.2
Dry Cambodia	214.7	157.8	46.0	33.6
Total Cambodia	427.3	300.6	178.5	122.8
Uppam in the Central Districts	12.3	20.0	1.9	3.9
Nadam and Bourbon	29.8	22.1	1.5	1.2
Total, Salems	42.1	42.1	3.4	5.1
Tinnevelles*	382.7	429.0	94.0	107.3
White and red Northernns	150.0	167.0	18.8	20.9
Westerns	762.0	744.0	95.3	93.0
Warangal and Cocanadas	102.7	90.1	19.5	16.8
Chinnapathi (short staple)	7.5	7.5	9	9

* Includes Karunganni in Coimbatore and Uppam, Karunganni and mixed country cotton in Madura, Ramnad and Tinnevely.

The local cotton trade is not generally active at this time of the year. The average wholesale price of cotton lint per imperial maund of 82½ lbs. as reported from important markets on 2nd December 1940 was about Rs. 16-7-0 for Cocanadas, Rs. 18-6-0 for White Northernns, Rs. 18-7-0 for Red Northernns, Rs. 15-6-0 for Westerns (*mungari* crop), Rs. 18-13-0 for Westerns (*jowari* crop), Rs. 30-15-0 for Coimbatore Cambodia, Rs. 26-8-0 for Southern Cambodia, Rs. 28-8-0 for Coimbatore Karunganni, Rs. 24-7-0 for Tinnevely Karunganni, Rs. 23-15-0 for Tinnevelles and Rs. 22-14-0 for Nadam cotton. When compared with the prices published in the last report, i. e., those which prevailed on 4th November 1940, these prices reveal a rise of about 12 per cent in the case of Southern Cambodia, 7 per cent in the case of Tinnevelles, 6 per cent in the case of White Northernns, Coimbatore Cambodia and Tinnevely Karunganni, 5 per cent in the case of Cocanadas, 3 per cent in the case of Westerns (*mungari*) and 2 per cent in the case of Red Northernns, the prices remaining more or less stationary in the case of Westerns (*jowari*), Coimbatore Karunganni and Nadam.

(From the Director of Industries and Commerce.)

Statistics—Crop—Cotton—1940-41—Intermediate condition report. Pickings of the *Mungari* or early sown crop in parts of the Deccan are nearing completion and the yield is expected to be slightly below normal. In Coimbatore, Madura, Ramnad and Tinnevely, the growth of the young plants is reported to have been affected to some extent by the heavy and continuous rains of November. The condition of the crop is generally satisfactory elsewhere in the Province.

The average wholesale price of cotton lint per imperial maund of 82½ lb (equivalent to 3,200 tolas) as reported from important markets on 6th January 1941 was Rs. 14-0-0 for Cocanadas, Rs. 16-12-0 for white Northernns, Rs. 16-7-0 for red Northernns, Rs. 12-14-0 for Westerns (*Mungari* crop), Rs. 17-12-0 for Westerns (*Jowari* crop), Rs. 30-13-0 for Coimbatore Cambodia, Rs. 27-8-0 for Coimbatore Karunganni, Rs. 24-2-0 for Tinnevely Karunganni, Rs. 20-13-0 for Tinnevelles and Rs. 21-13-0 for Nadam cotton. When compared with the prices published in the last report i. e., those which prevailed on 2nd December 1940, these prices reveal a fall of approximately 16 per cent in the case of Westerns (*Mungari* crop), 15 per cent in the case of Cocanadas, 13 per cent in the case of Tinnevelles, 11 per cent in the case of red Northernns, nine per cent in the case of white northernns, six per cent in the case of Western (*Jowari* crop), five per cent in the case of Nadam, four per cent in the case of Coimbatore Karunganni and one per cent in the case of Tinnevely Karunganni, the price remaining practically stationary in the case of Coimbatore Cambodia.

(From the Director of Industries and Commerce.)

Statistics—Crop—Groundnut—1940—Fourth or final report. The average of the areas under groundnut in the Madras Province during the five years ending 1938-39 has represented 48.6 per cent of the total area under groundnut in India.

The area sown with groundnut in the Province in 1940 is estimated at 3,820,000 acres. When compared with the corresponding estimate of 3,534,200 acres for the previous year and the actual area of 3,617,600 acres according to the Season and Crop Report of the previous year, the present estimate reveals an increase of 8.1 per cent and 5.6 per cent respectively. The estimated area for this year exceeds the average area of 3,415,210 acres by 11.9 per cent.

The increase in area is general outside Vizagapatnam, East Godavari, Kistna (-25,300 acres), Guntur (-28,800 acres), Cuddapah, Nellore, South Arcot (-25,100 acres), North Arcot, Tanjore and Ramnad and is marked in Kurnool (-69,800 acres), Bellary (+73,500 acres) Anantapur (+45,200 acres), Salem (+38,400 acres) and Coimbatore (+39,600 acres). The area in the Deccan rose from 1,226,300 acres in 1939-40 to 1,405,000 acres in the current year i. e., by 14.6 per cent.

The harvesting of the summer and early crop of groundnut had concluded by the end of October. The harvesting of the winter or main crop is proceeding.

The crop was affected to some extent by drought in Anantapur, Chittoor, North Arcot and Salem, by heavy rains in Nellore, South Arcot, Trichinopoly, Tanjore and Madura, and by insect pests in parts of Trichinopoly, Tanjore and Madura. The yield is expected to be above normal in Bellary, normal in East Godavari, West Godavari, Kistna and Guntur, Kurnool and Ramnad and below normal in the other districts. The yield is estimated to be low in South Arcot (70 per cent), North Arcot (75 per cent), Chingleput and Tanjore (77 per cent in each). The seasonal factor for the Province as a whole works out to 92 per cent of the average as against 95 per cent in the previous year according to the Season and Crop Report. On this basis, the yield is expected to be 1,760,800 tons of unshelled nuts as against 1,702,680 tons in the previous year, an increase of 3.4 per cent. The yield in an average year is estimated at 1,707,550 tons.

The wholesale price of groundnut (shelled) per imperial maund of 82½ lb. (equivalent to 3,200 tolas) as reported from important market centres on 6th January 1941 was Rs. 3-7-0 in Tadpatri, Rs. 3-3-0 in Coimbatore, Rs. 3-2-0 in Vizagapatam, Guntur and Hindupur, Rs. 3-0-0 in Vizianagaram and Cuddalore, Rs. 2-13-0 in Salem, Rs. 2-11-0 in Cuddapah and Vellore, Rs. 2-10-0 in Adoni and Bellary, Rs. 2-8-0 in Nandyal and Rs. 2-7-0 in Guntakal. When compared with the prices published in the last report, i. e., those which prevailed on 4th November 1940, these prices reveal a fall of approximately 27 per cent in Nandyal, 22 per cent in Vellore and Guntakal, 20 per cent in Cuddapah, 18 per cent in Adoni and Bellary, 14 per cent in Vizagapatam, Vizianagaram, Guntur and Cuddalore, 12 per cent in Hindupur, 10 per cent in Salem, eight per cent in Tadpatri and four per cent in Coimbatore.

(From the Director of Industries and Commerce.)

Statistics—Crop—Castor—1940—First or final report. The average of the areas under castor in the Madras Province during the five years ending 1938-39 has represented 17.5 per cent of the total area under castor in India.

The area under castor in the Madras Province up to 25th November 1940 is estimated at 2,50,000 acres. When compared with the area of 272,600 acres estimated during the corresponding period of last year, it reveals a decrease of 8.3 per cent. The estimate of last year was greater than the actual area of 266,051 acres by 2.5 per cent.

The crop is mainly grown in Guntur (27,000 acres), the Deccan (121,000 acres), Nellore (38,000 acres) and Salem. As compared with the actual area of last year, a decrease in area is estimated in Guntur, Anantapur (-21,200 acres), and Nellore and an increase in area in the other districts.

The yield is expected to be normal in all districts outside Vizagapatam. The seasonal factor for the Province as a whole is estimated to be 100 per cent of the normal. On this basis, the yield is estimated at 24,900 tons as against 26,800 tons estimated for the corresponding period of last year and 25,630 tons estimated in the season and crop report of last year.

The wholesale price of castor seed per imperial maund of 82½ lb. (equivalent to 3,200 tolas) as reported from important markets on 16th December 1940 was Rs. 6-4-0 in Vizianagaram, Rs. 6-0-0 in Guntur, Rs. 5-15-0 in Nandyal, Rs. 5-12-0 in Vizagapatam, Rs. 5-8-0 in Cuddapah, Rs. 5-0-0 in Bellary and Anantapur and Rs. 4-10-0 in Hindupur. When compared with the prices reported in the previous year, i. e., those which prevailed on 18th December 1939, these prices reveal a rise of approximately 14 per cent in Vizianagaram, 11 per cent in Anantapur, and ten per cent in Cuddapah and a fall of approximately 14 per cent in Nandyal and nine per cent in Bellary, the price remaining stationary in Hindupur.

(From the Director of Industries and Commerce.)

Subject:—Statistics—Ginger—1940—Final Report. The area under ginger in 1940 is estimated at 12,000 acres in Malabar and 800 acres in South Kanara as against the actual area of 11,112 acres in Malabar and 800 acres in South Kanara in the previous year.

The condition of the crop is generally satisfactory except in parts of Malabar where the crop is affected by "soft-rot" to some extent. The seasonal factor is estimated at 95 per cent of the normal in Malabar and 100 per cent in South Kanara. On this basis, the yield is estimated at 4,360 tons of dry ginger (4,070 tons in Malabar and 290 tons in South Kanara) as against 4290 tons the previous year (4,040 tons in Malabar and 250 tons in South Kanara).

(From the Director of Industries and Commerce.)

Statistics—Paddy—1940-41—Intermediate condition report. The main crop of paddy has been or is being harvested in parts of the Circars, the Deccan and the central districts. The yield is reported to be below normal in Vizagapatam and Chittoor; normal in the other districts.

The crop has been affected to some extent by disease in parts of Guntur, by insect pests in parts of Chingleput, North Arcot and Tanjore and by heavy rains in Tanjore. The condition of the crop is generally satisfactory in the other districts.

The wholesale price of paddy, second sort, per imperial maund of 82½ lb. (equivalent to 3,200 tolas) as reported from important markets on 6th January 1941 was Rs. 3-8-0 in Madura, Rs. 3-2-0 in Rajahmundry, Ellore, Bezwada, Guntur and Virudhunagar, Rs. 3-1-0 in Cocanada, Masulipatam and Trichinopoly, Rs. 3-0-0 in Vizianagaram and Tinnevely, Rs. 2-15-0 in Kumbakonam, Rs. 2-14-0 in Hindupur and Chittoor, Rs. 2-13-0 in Nagapatam, Rs. 2-12-0 in Vellore and Anantapur, Rs. 2-7-0 in Cuddalore and Rs. 2-1-0 in Canjeevaram. When compared with the prices published in the last report, i. e., those which prevailed on 9th December 1940, these prices reveal a rise of approximately 19 per cent in Anantapur and two per cent in Kumbakonam and a fall of approximately 18 per cent in Masulipatam, 12 per cent in Ellore and Vellore, 11 per cent in Cocanada, Rajahmundry, Bezwada and Guntur, eight per cent in Trichinopoly, six per cent in Tinnevely, five per cent in Cuddalore and four per cent in Chittoor, the prices remaining stationary in Vizianagaram, Hindupur, Canjeevaram, Nagapatam, Madura and Virudhunagar.

(From the Director of Industries and Commerce.)

Statistics—Pepper—1940—Final Report. The area under pepper in 1940 in Malabar and South Kanara is estimated at 104,000 acres (95,200 acres in Malabar and 8,800 acres in South Kanara) as against the final area of 104,384 acres (95,599 acres in Malabar and 8,795 acres in South Kanara) in the previous year.

The condition of the crop is generally satisfactory. The seasonal factor is estimated at 100 per cent as against 105 per cent in the previous year. On this basis, the yield is estimated at 9,990 tons (9,140 tons in Malabar and 870 tons in South Kanara) estimated in the previous year.

The wholesale price of pepper per imperial maund of 82½ lb. (equivalent to 3,200 tolas) as reported from important markets on 6th January 1941 was Rs. 9-14-0 in Calicut, Rs. 9-2-0 in Tellicherry and Rs. 10-2-0 in Mangalore. When compared with the prices published in the last report, i. e., those which prevailed on 9th September 1940, these prices reveal a fall of approximately 12 per cent at Calicut and seven per cent at Tellicherry the price remaining stationary at Mangalore.

(From the Director of Industries and Commerce.)

Statistics—Crop—Gingelly—1940-41—Third Report. The average of the areas under gingelly in the Madras Province during the five years ending 1938-39 has represented 16.2 per cent of the total area under gingelly in India.

The area sown with gingelly up to 25th December 1940 is estimated at 558,300 acres. When compared with the area of 601,960 acres estimated for the corresponding period of last year, it reveals a decrease of 7.2 per cent. The area estimated for Coimbatore is the highest reported in recent years.

The estimated area is the same as that of last year in West Godavari, Cuddapah and Malabar. An increase in area is estimated in Vizagapatam, East Godavari (plus 13,000 acres), Kurnool, Coimbatore, Trichinopoly and Tinnevely and a decrease in area in the other districts of the Province, especially in Anantapur (-13,000 acres), Chingleput (-14,000 acres) and Salem (-28,000 acres) owing to want of timely sowing rains.

The main crop has been harvested except in the South. The yield is estimated to be above normal in Kurnool, normal in Vizagapatam, East Godavari, Guntur, Cuddapah, Nellore, Salem, Coimbatore, Ramnad and South Kanara and below normal in the other districts, especially in South Arcot (70 per cent) in Chingleput and Tinnevely (75 per cent in each).

The seasonal factor for the Province as a whole works out to 94 per cent of the average as against 87 per cent for the corresponding period of last year. On this basis, the yield is estimated at 70,700 tons as against 69,400 tons for the corresponding period of last year, an increase of 1.9 per cent.

The wholesale price of gingelly per imperial maund of 82½ lb. (equivalent to 3,200 tolas) as reported from important markets on 6th January 1941 was Rs. 6-12-0 in Tinnevely, Rs. 6-8-0 in Cocanada, Rs. 6-7-0 in Cuddalore, Rs. 6-6-0 in Trichinopoly, Rs. 6-5-0 in Tuticorin, Rs. 6-1-0 in Rajamundry, Rs. 6-0-0 in Vizianagaram, Rs. 5-12-0 in Salem, Rs. 5-8-0 in Vizagapatam, and Rs. 5-3-0 in Ellore. When compared with the prices published in the last report, i. e., those which prevailed on 4th November 1940, these prices reveal a rise of approximately 15 per cent in Tuticorin 13 per cent in Cocanada and 7 per cent in Rajamundry and fall of approximately 13 per cent in Ellore, 4 per cent in Vizagapatam, Vizianagaram and Tinnevely and 3 per cent in Trichinopoly, the prices remaining stationary in Cuddalore and Salem.

(From the Director of Industries and Commerce.)

College News and Notes.

Students' Corner :—Students' Club. Under the auspices of the Students' Club Rao Bahadur T. S. Venkataraman, Sugarcane Expert, Imperial Sugarcane Research Station, delivered an inspiring lecture on 23-1-41 on Indian Village and our duty to it, Sri. N. Lakshmanan occupied the chair. The speaker dealt exhaustively on the out-standing problems confronting the present village folk. Recounting some of his personal experiences of the failure of agriculture as a profession the speaker stated that even the best practices of the West should be modified to suit the Indian conditions, if they are to be useful.

On 24-1-41 Mr. A. J. Macdonald, B. Sc., B. Sc. (Agri.) N. D. A., Officer in charge, Poultry Research Section, Imperial Veterinary Research Institute, Izatnagar delivered an educative and thought-provoking lecture on 'Poultry Farming' in India with Sri. K. Unnikrishna Menon in the chair. The speaker deplored the lack of statistics relating to the Poultry Industry in India and gave a short account of the various aspects of poultry work at Izatnagar and the possible lines of development of the Indian Poultry Industry.

College. The College reopened after the Christmas and New year holidays on the 4th January '41. The students of class II proceeded on an educational tour of the Southern districts of the province.

Selection for University Examination. Five students in the first year and one in second year classes were detained from appearance in the University examinations on the score of their unsatisfactory progress revealed in the terminal examinations.

Educational Tour. Starting from Coimbatore on the 3rd of January the second year class reached Tindivanam on the 4th night. The cultivation of oil seed crops and their marketing were the chief subjects of study at this centre. Reaching Palur on the 7th they made studies regarding the cultivation of sugarcane, paddy, groundnut, ragi etc. On the 10th the party visited the sugar factory belonging to Messrs. East India Distilleries and Sugar Factories Ltd. at Nellikuppam. They were shown round the distillery, sugar factory and the confectionery. The party halted at Chidambaram and visited the Annamalai University Colonisation lands. The progress of the much-discussed Colonisation scheme and the enterprise of the graduate Colonists was a matter of great interest to a batch of agricultural graduates in the making. The same night they reached Agricultural Research Station, Aduthurai and in the course of their stay of two days, made a study of the cultivation of paddy, sugarcane, plantains etc. Next day by noon the party paid a visit to a model farm run by Rao Sahib N. S. Kolandaiswami Pillai, retired Deputy Director of Agriculture, Trichinopoly. That evening they went to a village and held enquiries about the cultivation of plantains and turmeric. Some of the students took advantage of the proximity of the famous shrine at Srirangam to pay a visit to the place. The 14th day of January dawned on the gay and happy students in Koilpatti. Koilpatti afforded the students scope for the study of the cultivation of cotton, cumbu, cholam, thenai etc., and the opportunity was availed to make some rural enquiries. On the 16th morning the party left the place and reached the great and humming city of Madura by noon where a visit was arranged to the Municipal sewage farm. The glorious sight of the holy Meenakshi Temple and the Thirumalainayakan palace, made a great impression on the young minds. The party returned to Coimbatore on the 17th afternoon.

The students played a badminton match at Tindivanam and lost in a keen contest. At Koilpatti they played a volley ball match against the local high school team and had an easy victory. The party was led by Sri. K. Unnikrishna Menon, Senior Lecturer in Agriculture and Sri. P. A. Venkateswaran, Senior Teaching Assistant.

Games:—Football. In the inter-tutorial foot-ball match played in connection with the club day activities Mr. K. M. Thomas' wards defeated Mr. P. V. Ramiah's wards by 3 goals to 1 and Sri. C. R. Srinivasa Ayyangar's wards won over Sri. C. N. Narasimha Ayyangar's by 2 to 1.

Hockey. In the inter-class tournament played for the Victory cup the match between 1st and 3rd year classes ended in a draw on the first day, but in the replay the third year class defeated their enthusiastic juniors by 4 goals to 1.

New Year Honours. It is gratifying to note that the title of K. C. S. I. was awarded to Sir. C. P. Ramaswami Ayyar, Dewan of Travancore and a Patron of the Madras Agricultural Students' Union and that the title of Rao Bahadur was conferred on Sri. K. T. Alwa, Headquarters Deputy Director of Agriculture, Madras. We offer our congratulations to these friends of the Union on their well-merited distinctions.

Visitors. Mr. P. H. Rama Reddy, Director of Agriculture, Madras, Mr. T. J. Hurley, Director of Veterinary Services, Madras, Mr. M. J. Narasimham, Deputy Director of Agriculture, Mysore state, Mr. J. B. Polding Esq., Research Officer on contagious abortion, Imperial Veterinary Institute, Mukteswar, and Mr. M. J. Macdonald, Officer in charge, Poultry Research section, Imperial Veterinary Institute, Izatnagar, were among the visitors to the Agricultural College and Research Institute during the month.

Association of Economic Biologists, Coimbatore. At a meeting of the association held on the 20th December 1940, Rao Bahadur M. R. Ramaswami Sivan, Retired Principal, Agricultural College gave a lecture on 'The Anand Institute'. This was followed by a paper by Rao Bahadur G. N. Rangaswami Ayyangar and V. Achuta Warier on "The little Millet, *Panicum miliare*."

First aid Training. The training class for the benefit of the College Estate residents under the auspices of the St. John's Ambulance Association was inaugurated on the 22nd January 1941. The first class was held on the 26th January at which over fifty residents of the estate attended. We are grateful to the Coimbatore branch of the Ambulance Association and Dr. P. R. Kuppuswami for arranging the classes to be held at the College.

New Year's Day War Victory Celebration. The Agricultural College and Research Institute Estate celebrated New Year's Day in three meetings held in three different places—north, centre and south of the Estate area. Meetings were held at the Millets Breeding Station (North), the Central Farm (Centre) and the Paddy Breeding Station (South). All the labourers were assembled along with the villagers of the surrounding hamlets. After due publicity large gatherings were held, and the success over the Italians was explained to them. They were given an insight into the harrowing conditions of the War areas and of the peace and security enjoyed by us, under the protection of the strong arm of Britain. The villagers were made to realise the necessity for making liberal contributions towards this War effort. At the end of the meetings, sweets were distributed to the children.

OBITUARY

We regret to record the sad demise of Sri. P. Parthasarathy, B. Sc. Ag. on the night of 25th December at the early age of 33. He passed out of the Agricultural College, Coimbatore in 1933 and joined the Department first at Kodur. In recognition of his experience in fruit research work he was recently posted to Guntur for developing the newly started orchard at this place. He took ill recently and passed away in spite of the best medical help. He was a capable and enthusiastic young man and in his death the Department loses a very promising officer. He leaves behind him his young widow, aged mother, several relations and a host of friends to bemoan his loss.

Moffussil News.

Agricultural Exhibition at Tiruvannamalai. An Exhibition stall was opened in the Fourth All India Swadeshi Exhibition held at Tiruvannamalai, on the occasion of the *Karthigai Deepam* Festival, from the 5th to the 16th. The Exhibition Committee afforded all facilities for arranging the stall. The exhibits were arranged into eight sections, comprising of seeds and crops, implements, manure preservation, horticulture, cream jaggery making, malt making, insect pests and diseases and bee-keeping. The exhibits in each section were arranged tastefully and supplemented with coloured pictorial and word posters. The Coimbatore strains of paddy and the ground-nut specimens of seeds and plants from Tindivanam Research station elicited many inquiries. The horticultural exhibits included fruit plants from Kodur, fruit research station, fruits from Coonoor, posters illustrating care and management of citrus gardens and preserves which demonstrated the possibility of the utilisation of surplus fruit. Considerable interest was evinced in all these exhibits. Malt making which was demonstrated as a practicable cottage industry was appreciated by many visitors. The section dealing with insect pests and diseases with control measures elicited nearly 64

inquiries from persons coming from different districts. Bee-keeping exhibits effectively kept a crowd of interested people always about them. Honey and wax produced from the Haji Apiaries, Vaniambadi, were sold during the period and there was a large demand for more honey.

The Exhibition was opened by the Revenue Divisional Officer, Tiruvannamalai, Sri. M. P. Narayanan Nair. The committee of judges awarded a Gold medal certificate to the Department.

Denkanikottai Rural Exhibition. A Rural Agricultural Exhibition was organised under the auspices of the Village Uplift Sangham, Denkanikottai, Hosur taluk, Salem. The Exhibition attracted considerable interest from all the surrounding villages. There was keen competition amongst the exhibitors for the prizes. Besides exhibits kept in the stall crops were judged in situ in the fields. The Department exhibited improved strains of seeds and iron implements for various agricultural operations suited to the locality. The exhibits were judged by the Sub-Collector, Hosur and prizes in the form of Seeds, Implements and Manures were awarded to the best exhibits.

Weather Review—DECEMBER 1940.

RAINFALL DATA

Division	Station.	Actual for month	Departure from normal @	Total since January 1st	Division	Station	Actual for month	Departure from normal @	Total since January 1st
Circars	Gopalpore	0.0	-0.7	72.1	South	Negapatam	18.1	+6.7	76.3
	Calingapatam	0.0	-0.7	45.9		Aduthurai *	5.6	-2.5	50.3
	Vizagapatam	0.7	0.0	34.6		Madura	1.9	+0.1	44.6
	Anakapalli *	0.0	0.0	0.0		Pamban	4.1	-3.4	48.4
	Samalkota *					Koilpatti *	5.8	+3.1	44.2
	Maruteru *	0.0	-0.3	40.6		Palamkottah	7.9	+3.9	33.7
	Cocanada	0.0	-0.9	46.0	West Coast	Trivandrum	5.2	0.0	71.8
Ceded Dists.	Masulipatam	0.2	-0.7	41.2		Cochin	2.7	+1.0	132.1
	Guntur *	1.1	+0.2	34.3		Calicut	0.0	-1.1	128.8
	Kurnool	0.0	-0.2	29.1		Pattambi *	1.0	+0.2	99.0
	Nandyal *	1.4	+1.3	24.3		Taliparamba *	0.2	-0.5	149.1
	Hagari *	0.2	+0.1	22.0		Kasargode *	0.7	-0.1	152.5
	Siruguppa *	0.1	0.0	24.5		Nileshwar *	0.5	-0.2	157.3
	Bellary	0.1	0.0	22.1		Mangalore	0.6	+0.1	150.3
	Anantapur	0.0	-0.3	27.3	Mysore and Coorg	Chitaldrug	0.1	-0.2	34.0
	Rentachintala	0.7		27.4		Bangalore	0.6	+0.1	36.5
	Cuddapah	0.0	-0.9	39.9		Mysore	0.9	+0.5	48.4
Carnatic	Anantharajupet *	0.9	-0.7	50.4		Mercara	0.6	-0.1	143.0
	Nellore	1.6	-1.6	55.2	Hills	Kodaikanal	1.6	-2.8	73.4
	Madras	5.6	-0.2	52.2		Coonoor			
	Palur *	10.4	+3.4	69.9		Ootacamund *	1.3	+0.1	53.7
	Tindivanam *	4.2	0.0	47.9	Central	Nanjanad *	1.0	-0.7	52.4
Central	Cuddalore	7.8	+0.6	77.3					
	Vellore	1.4	-1.3	37.2					
	Salem	0.4	-0.6	39.7					
	Coimbatore	3.3	+2.1	40.9					
	Coimbatore								
	A. C. & R. I. *	3.3	+1.8	33.7					
	Trichinopoly	2.4	-0.2	36.7					

* Meteorological Stations of the Madras Agricultural Department.

@ From average rainfall for the month calculated up to 1937 (published in Fort St. George Gazette).

The weather during the first three days of the month was generally dry over the whole area. But from the 4th till the 6th and again between the 11th and 14th the trough of low pressure over the south of the Bay was again active as in the month of November and occasioned widespread and locally heavy rain in the South East of the peninsula. On the 15th of the month conditions were unsettled in the south of the Bay but failed to develop into a depression. Conditions were again unsettled in the South East of the Bay on the 20th, and developing into a depression on the next day moved north east and crossed the North Burma coast by the 23rd. Thereafter conditions were more settled and weather dry with a tendency for falling temperatures over South India.

Rainfall was in large excess locally in the Coromandel Coast, and in the Central Districts and normal or in defect elsewhere.

Other climatic elements were nearly normal. The chief falls reported were:

Panruti	(South Arcot)	8.1" (4th)
Chidambaram	(do.)	6.7" (..)
Tiruchendur	(do.)	6.1" (..)
Vridhachalam	(do.)	6.2" (5th)
Cuddalore	(do.)	5.7" (4th)
Porto Nova	(do.)	5.6" (5th)
Negapatam	(Tanjore)	5.3" (6th)

Note on the total rainfall for 1940. The rainfall for the twelve months of the calendar year 1940 was characterised by its being in very large excess over the whole presidency with the exception of parts of the Ceded districts and the Nilgiri hills.

The first three months of the year were generally dry except for locally heavy rainfall in the northern districts due to the effect of secondary low pressure areas derived from western disturbances traversing the upper part of the peninsula.

During the hot weather months of April and May rainfall was generally in excess owing to thunderstorm activity during April and with the formation of a depression in the Bay which stimulated a temporary advance of the monsoon in May, when heavy rainfall occurred in the Circars.

The monsoon set in over the peninsula early in June and continued strong during July and August. During these months rainfall was in excess in Malabar coast and in the Circars. The rainfall was generally below normal everywhere during September and outside the Ceded districts in October, but in November the weather over the south was continuously unsettled and rainfall in large excess, these unsettled conditions extended till the middle of December and rainfall was again heavy in the south during that period.

The annual rainfall was in large excess over the whole of the presidency outside parts of the Ceded districts and Nilgiris and most markedly so in the Ganjam district of Orissa, and on the Coromandel Coast and in parts of the Central districts, West Coast and Mysore.

Weather Report for the Agricultural College and Research Institute Observatory :
Report No. 12/40.

Absolute maximum in shade	...	89°F
Absolute minimum in shade	...	58.0°F
Mean maximum in shade	...	83.1°F
Departure from normal	...	-0.7°F
Mean minimum in shade	...	66.0°F
Departure from normal	...	+0.9°F
Total rainfall for the month	...	3.29"

Departure from normal	...	+1.8"
Heaviest fall in 24 hours	...	1.40"
Total number of rainy days	...	6
Mean daily wind velocity	...	1.23 m. p. h.
Departure from normal	...	-1.73
Mean humidity at 8 hours	...	78%
Departure from normal	...	-0.8%

Summary. Under the influence of unsettled weather, rainfall was in large excess. Skies were generally moderately clouded. Day and night temperatures were nearly normal.
P. V. R. & R. S.

Departmental Notifications.

Gazette Notification.

Transfers.

Name of officers.	From	To
Sri. U. Vittal Rao,	Offg. Asst. Marketing Officer, Madras,	Offg. Asst. D. A., Tellicherry.
" G. Ganapathi Ayyar,	Asst. in Chemistry, Agri. Res. Inst. Coimbatore,	Offg. Asst. Agri. Chemist in charge of the Agri. Res. Station, Siruguppa.

Leave.

Name of officers.	Period of leave.
Sri. K. T. Alwa, Headquarters Dy. Director of Agriculture, Madras.	L. a. p. for 4 months from 16-2-41 preparatory to retirement.
" C. V. Ramaswami Ayyar, Asst. Agri. Chemist. A. R. S. Siruguppa.	L. a. p. for 2 months and 26 days with effect from the date of relief.

Subordinate Services.

Transfers.

Name of officers.	From	To
Sri. R. Narasimha Ayyar,	A. D. in Mycology St. Thomas Mount,	Mycology section, Coimbatore,
Mr. Herbert Adisheshaiah,	A. D. Vellore,	Sugarcane Resaarch Station, Gudiyattam.
" S. D. S. Albuquerque,	Botanical Asst., A. R. S. Pilicode,	A. R. S. Neleswar III.
" S. V. Kuppuswami,	Asst. in Chemistry, A. R. S. Siruguppa,	Asst. in Chemistry Section, Coimbatore.
Sri. E. Kunhappa Nambiar,	Upper Subordinate (on leave)	F. M. Taliparamba.
Janab Soopi Hajee	Asst. F. M. Taliparamba	Asst. A. D. Manjeri.
Sri. C. Raman Moosad,	A. D., Manjeri,	A. D., Wallajah.
" M. A. Balakrishna Iyer,	A. D. Wallajah,	A. D., Vellore.
" S. P. Fernando,	Asst. A. D. (on leave),	A. D. Ramnad.
" A. B. Adishesha Reddy,	A. D. (on leave),	A. D., Guntakal.
" M. Vaidyanathan,	A. D. Guntakal,	A. D., Rayadrug.

Leave.	
Name of officers.	Period of leave.
Sri. M. S. Kylasam, Asst. to the Entomologist, Coimbatore.	L. a. p. for 30 days from 30-1-41.
„ S. Madhava Rao, F. M. Central Farm, Coimbatore.	Extension of l. a. p. for 28 days from 12-1-1941.
„ A. Shanmugasundaram, A. D., Pattukottai.	L. a. p. for 25 days from 16-1-41.
„ K. Hanumantha Rao, A. R. S., Pattambi.	L. a. p. for 49 days from 20-1-41.
„ D. S. Subramania Ayyar, A. D., (on leave).	Extension of l. a. p. on m. c. for 6 weeks from 24-12-40.
„ M. Vaidyanathan, A. D., Guntakal.	L. a. p. for 3 months and 15 days from 15-1-41.
„ G. Venkataramana, A. D., Bapatla.	Extension of leave on half average pay on m. c. for 1 month from 24-12-40.
„ M. Bhavani Shanker Rao, Asst. Groundnut scheme, A. R. S., Tindivanam.	Earned leave for 30 days from 6-1-41.
„ L. Krishnan, A. D., Tanjore	Extension of l. a. p. on m. c. for 6 weeks from 9-1-41.
„ V. Achyutharamiah, A. D., Jami.	Extension of l. a. p. on m. c. for 6 weeks.
„ K. Sreeraman, Asst. in Chemistry, A. R. S., Siruguppa.	L. a. p. for 29 days from 3-1-41.
„ C. K. Ramachandran, Asst. in Cotton, D. F. S., Hagari.	L. a. p. for 30 days from 3-1-41.
„ N. Ranganathachari, A. D., Dhone.	Extension of l. a. p. for 1 month on m. c. from 19-12-40.
„ V. Ratnajirao, A. D., (on leave).	L. a. p. for 4 months with m. c. from 16-12-40.
„ C. S. Gopalaswami Rao, Mycology Asst. Bellary.	L. a. p. for on m. c. for 2 months from 7-1-41.
„ Raman Moosad, A. D., Manjeri.	L. a. p. for 1 month from the date of relief.

Agricultural College and Research Institute, Coimbatore.

Additions to the Library during the quarter ending 31st December 1940.

A. Books.

1. *Elementary Lessons in Agriculture in Telugu*. Jogi Raju, G. (1940).
2. 'Pandlu' — *Fruits Part I — General in Telugu*. Jogi Raju, G. (1936).
3. 'Pandlu' — *Fruits Part II — Mango in Telugu*. Jogi Raju, G. (1940).
4. *Elementary Agriculture*. Mitra, S. K. (1940).
5. *Savola Kasar: Study of a Deccan Villages in the Famine Zone*. Jagalpure, L. B. & Kale, K. D. (1938).
6. *Price Fixing by Government in Foreign Countries, 1926-1939 — A selected list of references*. Hannay, A. M. (1940).
7. *Agricultural Produce Act, 1937 with rules made prior to 31st August 1940*. Government of India Publication. (1940).
8. *Report on the marketing of Grapes in India*

& Burma - Marketing Series No. 20. Government of India Publication. (1940). 9. *Report on the marketing of Coffee in India & Burma - Marketing Series No. 121*. Government of India Publication. (1940). 10. *Report of the Committee on Co-operation in Madras - 1939-40*. 11. *Planting Directory of Southern India*. U. P. A. S. I. Publication. (1940). 12. *Plant Physiology - A Text Book for College & Universities*. Meyer, B. S. and Anderson, D. B. (1940). 13. *Colorimetric methods of Analysis - Volume II - Organic and Biological*. Snell, F. D. & Snell, C. T. (1937). 14. *Diptera - Volume VI - Family Calliphoridae (Fauna of British India Series)*. White, R. S. & others. (1940). 15. *Records and Research in Engineering and Industrial Science*. Holmstrom, J. E. (1940). 16. *Utilization of Indian Vegetable Oils and Lubricants in Internal Combustion Engines*. Aggarwal, J. S. and Verman, L. C. (1940). 17. *Broadcasting in India - The first report (1938-39) of the office of the Controller of Broadcasting*. Government of India Publication. (1940).

B. Annual Reports of The Agricultural Department etc.

1. *Proceedings of the Association of Economic Biologists, Coimbatore*. (1938).
2. *Administration Report of the Madras Electricity Department for the year 1939-40*.
3. *Annual Report of the Mysore Agricultural Department for 1938-39*.
4. *Report on the Administration of the Meteorological Department of the Government of India in 1939-40*.
5. *Annual Report of the Agricultural Meteorology Section, Indian Meteorological Department for 1938-39*.
6. *Proceedings of the 29th meeting of Advisory Board of the Imperial Council of Agricultural Research*. 1940.
7. *Administration Report of the Baluchistan Agency for 1938-39*.
8. *Annual Administration Report of the Department of Agriculture - United Provinces for 1938-39*.
9. *Annual Report of the Department of Agriculture - Baroda State - for 1938-39*.
10. *Report on the Experimental Farms in the Central Provinces and Berar - for 1938-39*.
11. *Proceedings of the Indian Society of Soil Science*. 1939-40.
12. *Report on the operations of the department of Agriculture, Burma for 1939-40*.
13. *Report of the Agricultural Stations in Burma for 1939-40*.
14. *Annual Report of the Grain Research Laboratory, Winnipeg, Manitoba, Canada for 1939*.
15. *Annual Report of the Department of Agriculture, Malaya for the year 1939*.
16. *Canadian Seed Growers' Association - Annual Report, 1937-33*.
17. *Report of the Department of Agriculture - British Honduras - 1939*.
18. *Cornell University Agricultural Experiment Station - Annual Report for 1939*.
19. *Annual Report of the Iowa Agricultural Experiment Station for 1938-39*.
20. *Annual Report of the Iowa Corn Research Institute for 1938-39*.
21. *Transaction of the Iowa State Horticultural Society for the year 1939*.
22. *Annual Report of the Nebraska Agricultural Experiment Station for 1939*.
23. *Annual Report of the Ohio Agricultural Experiment Station for 1937-33*.

New Periodicals.

Report on the Seaborne Trade of the Province of Madras for the year 1939-40.

52
Jan 21, 1941
NU. 29. 11



The Madras Agricultural Students' Union.

PATRONS

Sri V. Arumugam Pillai, Sundaikamuthur, Coimbatore District.
 Dewan Bahadur C. Arunachala Mudaliar, Zamindar of Chunampet, Chingleput
 Sri Lodd Govindoss, Madras. [District]
 Sri A. M. Kalinga Rayar, Zamindar of Uttukuli, Coimbatore District.
 Raja Sree Krishnachandra Gajapati Narayana Deo, Maharajah of Parlakimidi,
 Ganjam District.
 Raja V. Madhava Rajah, Rajah of Kollengode, Malabar District.
 Sir A. P. Patro, Berhampur.
 Sri T. A. Ramalingam Chettiar, Coimbatore.
 Sir C. P. Ramaswami Ayyar, Trivandrum.
 Dewan Bahadur C. S. Ratnasabapathi Mudaliar, Coimbatore.
 Sri M. Sambanda Mudaliar, Coimbatore.
 „ Veerarayan Thirumalpad, Nilambur, Malabar District.
 „ V. C. Vellingiri Gounder, Coimbatore.
 Dewan Bahadur C. V. Venkataramana Ayyangar, Coimbatore.
 Sri. R. Jeevandas, Varnasi, Parlakimidi.
 G. R. Hilson, Esq., Norwich, England.
 Sri Ramachandra Ananga Bhima Dev, Kesari Gajapathy, Rajah Saheb of
 Bodokhemidi, Berhampore, Ganjam.

OFFICE BEARERS FOR 1940—41.

Rao Bahadur G. N. Rangaswami Ayyangar, *President, (ex-officio).*
 C. R. Srinivasa Ayyangar, *Resident Vice President.*
 K. M. Thomas, *Editor.*
 T. Nataraj, *Secretary.*

COUNCIL (16 members),

President.
 Resident Vice president.
 Editor
 Secretary
Moffussil Vice presidents.
 Sri M. V. Raghava Rao
 „ M. Anandan.
 „ R. Swami Rao.
Moffussil Members.
 Sri M. D. Prabhu
 „ K. Jagannatha Rao.
 Janab Mahommed Ali Sahib.
 Sri K. Kuppumuthu.
Resident Members.
 Rao Sahib V. Muthuswami Ayyar,
 Sri M. S. Kylasam.
 „ C. Balasubramanian.
 „ G. Rama Rao (Student).
 „ N. Baskara Reddy, Secy. Students'
 Club (*ex-officio*).

MANAGING COMMITTEE (10 members).

Resident Vice President.
 Editor.
 Secretary.
 Sri. K. Ramaswami, *Manager.*
 „ A. Mariakulandai, *Treasurer.*
 „ K. Sanjiva Shetty, *Member.*
 „ P. A. Venkateswara Ayyar, „
 „ V. Gomathinayakam Pillai, „
 „ T. Chellappa, (Student.) „
 N. Baskara Reddy, Secretary,
 Students' Club (*ex-officio*)
 EDITORIAL BOARD (8 Members).
 Editor.
 Secretary.
 Manager.
 Sri. M. C. Cherian
 „ M. Kantiraj
 Dr. N. Krishnaswamy
 Sri. R. Narayanamurthy (Student).
 „ N. Baskara Reddy, Secy. Students'
 Club (*ex-officio*)

Editor and Publisher : K. M. Thomas, B.A., M. Sc., D. I. C.
 Printed at The Scholar Press, Palghat.

The SERVICES


Prefer...



*the 100% service
wrist watch...*

ALRESIST

All ranks of the fighting services hail ALRESIST as the ideal watch. Water- and dust - proof, non - magnetic, shock-proof, rustless, fitted with unbreakable glass, and opened when necessary without special tools, ALRESIST is the watch which meets everyone's needs.



3213 Staybrite case Rs. 58
3212 30 microns Rolled gold case, Staybrite back Rs. 74
3216 9ct. gold case, Staybrite back Rs. 100

FAVRE-LEUBA & CO. LTD.
BOMBAY ★ RANGOON ★ CALCUTTA