

No. 1. Read the following letter from J. BROUGHTON, Esq., Government Quinologist, to the Chief Secretary to Government, Fort St. George, dated Ootacamund, 19th December 1872:—

I HAVE the honour to report on a chemical examination I have, during the past eighteen months, been making on the bark of *Azadirachta Indica*, well known throughout India as the Margosa tree, though in the South more generally called the Nim tree.

2. The use of the bark of this well-known tree is quite general throughout the whole of India as a tonic and febrifuge. It is generally employed in the form of a decoction, though sometimes as powder, and has been stated by certain English authorities to have the same febrifuge qualities as Cinchona bark. The taste of the bark is intensely bitter; the inner layer or liber is, however, by far the bitterest. The tree being very handsome, and the wood of value, it is well known in avenues; the plants generally bearing marks of having been robbed of their bark for medicinal use. The pinnate leaves have also an intensely bitter taste, and are employed in native medicine; though not as a febrifuge, being used as a poultice externally, and are said to have a powerful effect in preventing glandular tumours from coming to maturity. They are also employed by natives in their fresh green state to cover the bodies of patients suffering from small-pox, though, I believe, this is done rather from a religious than a medicinal motive. The seeds yield a considerable amount of oil, which has also a strong bitter taste, is used for lamps, and in medicine. The roots are said to have vermifuge properties but I have not examined them. The main virtues of the tree reside in the bark.

3. The investigation has been a matter of much labour and difficulty from the circumstance of the bitter principle having scarcely any definite qualities, it being a neutral resin. It is slightly soluble in water, and hence the decoctions of the bark are intensely bitter, though water will not extract the whole present. To obtain it in purity the bark is exhausted with spirit of about 60 per cent., the filtered tincture precipitated with water, the dried precipitate again exhausted with benzol, again evaporated, again exhausted with carbonic disulphide, then by dry ether, and finally by absolute alcohol. The last exhaustion causes the separation of a small amount of a white transparent substance crystallizing in fine needles, which is quite insoluble in water, has no taste, and consists of carbon, hydrogen, and oxygen. Obtainable in but small quantities it has not been satisfactorily analysed, and has the general characters of a crystalline fat though fusing at 175° C. It is certainly not the effective principle of the bark nor an alkaloid.

4. The resin purified by the successive solvents is a dark brown somewhat soft resin of agreeable smell, resembling that of the bark. It is slightly soluble in water and entirely in all the foregoing solvents, but not in fixed oils. In any dilute solution its bitter taste is strong, but not disagreeable. It is soluble in strong boiling solutions of the fixed alkalies, from which acids again precipitate it apparently altered, likewise in strong sulphuric acid by which it is also altered. It fuses at 92° C.

5. The determination of its true chemical formula has been a matter of great difficulty and labour from the circumstance of its refusal to combine with other substances in a definite manner. It was finally determined by the discovery of a nitro-compound which yielded a silver salt, not however crystalline. From

the analyses of these I ascribe to the resin the formula $C_{28}H_{40}O_{11}$. Its nitro-compound is formed in the usual manner by nitric acid and precipitation with water, is yellow, uncrystallizable, somewhat soluble in water, more readily in alkalis, and having the formula $C_{28}H_{40}(NO_2)_4O_{11}$.

6. The resin is, therefore, not in any respect an alkaloid, as it contains no nitrogen. Its chemical constitution appears to me scarcely to confirm the ascribed anti-periodic febrifuge qualities of the bark, though doubtless the bitter solution of the resin may be a good tonic. If required for medicinal purposes it will be most conveniently employed in alcoholic solution. As the bark varies very considerably in its content of the bitter resin, a direct tincture would be unsuitable, but the impure resin after the application of benzol in the foregoing process of preparation would be fit for solution in alcohol and employment in medicine.

7. It is singular that native physicians never appear to have employed a tincture of the bark instead of an aqueous decoction. From some conversation I had with a native physician in Bengal I believe it is the native religious dislike in that country to the use of alcohol which has prevented its use for this purpose.

8. The leaves contain a small amount of bitter substance of a similar nature, but which is far more soluble in water. The same substance is also contained in the bark in addition to the one described. It has very similar properties to the resin, of which it is a hydrate. No peculiar alkaloid is contained in the leaves. The oil contains a small amount of the bitter substance of the leaves, not the resin of the bark. The powerful smell of the tree is well known, but it is not due to the presence of a sulphuretted oil as has been surmised; indeed, I have not succeeded in obtaining an essential oil from the tree. If the bark is distilled with water, the distillate has the perfume of the tree, but no oil is perceptible, though it has the power of decolourizing potassic permanganate.

9. A condensed account of our present knowledge of the therapeutics and chemistry of *Azadirachta Indica* is to be found in the Indian Pharmacopœia, p.p. 53 & 443, from which, however, I differ in some points. My conclusions are given in the preceding pages.

No. 2. ORDER THEREON, 3rd February 1873, No. 133.

Miscellaneous.

ORDERED to be communicated to
the Inspectors-General, British and
Indian Medical Departments.

2. Copy will be forwarded to the Secretary of State and Government of India.

3. Mr. Broughton is thanked for this report.

(True Extract.)

(Signed) D. F. CARMICHAEL,
Secretary to Government.

To the Inspector-General, Indian Medical Department.
" the " British Medical Service.
" J. Broughton, Esq.

Ed. F. R. Bantleman.

Govt. of 1873 Madras.

REVENUE.

THE INSPECTOR-GENERAL, INDIAN
MEDICAL DEPARTMENT.

Secy. Secy. Secy.

Dated February 3rd.

Recd.

Encl.

G. O. 3rd February 1873, No. 133.

FEBRUARY.

Nos. 1-2.

Azadirachta Indica.—Report on a chemical examination of it by Mr. Broughton ordered to be communicated to the Inspectors-General, British and Indian Medical Departments, and to the Secretary of State; and Mr. Broughton thanked for it.

Proceedings of the Madras Government, Revenue Department,
29th January 1873.

No. . Read the following letter from J. BROUGHTON, Esq., Government
Quinologist, to the Chief Secretary to Government, Fort St. George, dated
Ootacamund, 19th December 1872:—

I HAVE the honour to report on a chemical examination I have, during the
past eighteen months, been making on the bark of *Azadirachta Indica*, well known
throughout India as the Margosa tree, though in the South more generally called
the Nim tree.

2. The use of the bark of this well-known tree is quite general throughout
the whole of India as a tonic and febrifuge. It is generally employed in the
form of a decoction, though sometimes as powder, and has been stated by certain
English authorities to have the same febrifuge qualities as Cinchona bark. The
taste of the bark is intensely bitter; the inner layer or liber is, however, by far the
bitterest. The tree being very handsome, and the wood of value, it is well known
in avenues; the plants generally bearing marks of having been robbed of their
bark for medicinal use. The pinnate leaves have also an intensely bitter taste, and
are employed in native medicine; though not as a febrifuge, being used as a poultice
externally, and are said to have a powerful effect in preventing glandular tumours
from coming to maturity. They are also employed by natives in their fresh green
state to cover the bodies of patients suffering from small-pox, though, I believe, this
is done rather from a religious than a medicinal motive. The seeds yield a consid-
erable amount of oil, which has also a strong bitter taste, is used for lamps, and
in medicine. The roots are said to have vermifuge properties, but I have not
examined them. The main virtues of the tree reside in the bark.

3. The investigation has been a matter of much labour and difficulty from
the circumstance of the bitter principle having scarcely any definite qualities, it
being a neutral resin. It is slightly soluble in water, and hence the decoctions
of the bark are intensely bitter, though water will not extract the whole present.
To obtain it in purity the bark is exhausted with spirit of about 60 per cent., the
filtered tincture precipitated with water, the dried precipitate again exhausted
with benzol, again evaporated, again exhausted with carbonic disulphide, then
by dry ether, and finally by absolute alcohol. The last exhaustion causes the
separation of a small amount of a white transparent substance crystallizing in fine
needles, which is quite insoluble in water, has no taste, and consists of carbon,
hydrogen, and oxygen. Obtainable in but small quantities it has not been satis-
factorily analysed, and has the general characters of a crystalline fat but fusing at
175° C. It is certainly not the effective principle of the bark nor an alkaloid.

4. The resin purified by the successive solvents is a dark brown somewhat
soft resin of agreeable smell, resembling that of the bark. It is slightly soluble in
water and entirely in all the foregoing solvents, but not in fixed oils. In any
dilute solution its bitter taste is strong, but not disagreeable. It is soluble in
strong boiling solutions of the fixed alkalis, from which acids again precipitate it
apparently altered, also in strong sulphuric acid by which it is also altered. It
fuses at 92° C.

5. The determination of its true chemical formula has been a matter of
great difficulty and labour from the circumstance of its refusal to combine with
other substances in a definite manner. It was finally determined by the discovery
of a nitro-compound which yielded a silver salt, not however crystalline. From

the analyses of these I ascribe to the resin the formula $C_{30} H_{40} O_{11}$. Its nitro-compound is formed in the usual manner by nitric acid and precipitation with water, is yellow, uncrystallizable, somewhat soluble in water, more readily in alkalis, and having the formula $C_{30} H_{40} (NO_2)_4 O_{11}$.

6. The resin is, therefore, not in any respect an alkaloid, as it contains no nitrogen. Its chemical constitution appears to me scarcely to confirm the ascribed anti-periodic febrifuge qualities of the bark, though doubtless the bitter solution of the resin may be a good tonic. If required for medicinal purposes it will be most conveniently employed in alcoholic solution. As the bark varies very considerably in its content of the bitter resin, a direct tincture would be unsuitable, but the impure resin after the application of benzol in the foregoing process of preparation would be fit for solution in alcohol and employment in medicine.

7. It is singular that native physicians never appear to have employed a tincture of the bark instead of an aqueous decoction. From some conversation I had with a native physician in Bengal I believe it is the native religious dislike in that country to the use of alcohol which has prevented its use ~~in this case.~~ *for this purpose*

8. The leaves contain a small amount of bitter substance of a similar nature, but which is far more soluble in water. The same substance is also contained in the bark in addition to the one described. It has very similar properties to the resin, of which it is a hydrate. No peculiar alkaloid is contained in the leaves. *2 of* The oil contains a small amount of the bitter substance of the leaves, not the resin of the bark. The powerful smell of the tree is well known, but it is not due to the presence of a sulphuretted oil as has been surmised; indeed, I have not succeeded in obtaining an essential oil from the tree. If the bark is distilled with water, the distillate has the perfume of the tree, but no oil is perceptible, though it has the power of decolourizing potassic permanganate.

9. A condensed account of our present knowledge of the therapeutics and chemistry of *Azadirachta Indica* is to be found in the Indian Pharmacopœia, p.p. 53 & 443, from which, however, I differ in some points. My conclusions are given in the preceding pages.

Revenue 1872 Sept
From, No. 4078.
The Govt. Zoologist
Cotacamund

To, The Chief Secretary
to Government
Fort Saint George

Dated 19th Dec 1872
Recd 23rd

Enclosures — none

Madrashta Indica
Report on a chemical
examination of.

with the

6. 20. 1872
1. 20. 1872
2. 20. 1872
3. 20. 1872
4. 20. 1872
5. 20. 1872
6. 20. 1872
7. 20. 1872
8. 20. 1872
9. 20. 1872
10. 20. 1872
11. 20. 1872
12. 20. 1872
13. 20. 1872
14. 20. 1872
15. 20. 1872
16. 20. 1872
17. 20. 1872
18. 20. 1872
19. 20. 1872
20. 20. 1872
21. 20. 1872
22. 20. 1872
23. 20. 1872
24. 20. 1872
25. 20. 1872
26. 20. 1872
27. 20. 1872
28. 20. 1872
29. 20. 1872
30. 20. 1872
31. 20. 1872
32. 20. 1872
33. 20. 1872
34. 20. 1872
35. 20. 1872
36. 20. 1872
37. 20. 1872
38. 20. 1872
39. 20. 1872
40. 20. 1872
41. 20. 1872
42. 20. 1872
43. 20. 1872
44. 20. 1872
45. 20. 1872
46. 20. 1872
47. 20. 1872
48. 20. 1872
49. 20. 1872
50. 20. 1872
51. 20. 1872
52. 20. 1872
53. 20. 1872
54. 20. 1872
55. 20. 1872
56. 20. 1872
57. 20. 1872
58. 20. 1872
59. 20. 1872
60. 20. 1872
61. 20. 1872
62. 20. 1872
63. 20. 1872
64. 20. 1872
65. 20. 1872
66. 20. 1872
67. 20. 1872
68. 20. 1872
69. 20. 1872
70. 20. 1872
71. 20. 1872
72. 20. 1872
73. 20. 1872
74. 20. 1872
75. 20. 1872
76. 20. 1872
77. 20. 1872
78. 20. 1872
79. 20. 1872
80. 20. 1872
81. 20. 1872
82. 20. 1872
83. 20. 1872
84. 20. 1872
85. 20. 1872
86. 20. 1872
87. 20. 1872
88. 20. 1872
89. 20. 1872
90. 20. 1872
91. 20. 1872
92. 20. 1872
93. 20. 1872
94. 20. 1872
95. 20. 1872
96. 20. 1872
97. 20. 1872
98. 20. 1872
99. 20. 1872
100. 20. 1872