

(1) Number of pages 114(2) Date of repair 20-1-86 21-1-86(3) Initials of the Referencer lvmb**TAMIL NADU ARCHIVES****GOVERNMENT OF TAMIL NADU***Revenue* **DEPARTMENT****G.O. No.**
B.P. No.A79**Press.**
Mis.**Dated**28-1-86**197 .**

Not to be folded or roughly handled

1179
1

GOVERNMENT OF MADRAS.

REVENUE DEPARTMENT.

28th April 1880, No. 479.

No. 249. READ THE FOLLOWING PAPER:—

Extract from the Proceedings of the Government of India in the Home, Revenue, and Agricultural Department (Agriculture and Horticulture), under date Fort William, 13th February, 1880, No. 3—74-84.

READ—

A despatch from the Secretary of State, No. 144, dated the 11th December 1879, forwarding copy of a letter from Sir Philip Cunliffe Owen relative to a collection of the wild silks of India presented by Mr. Thomas Wardle to the Chamber of Commerce of Lyons, and also copy of a lecture by Mr. Wardle on the wild silks of India, principally tasar.

ORDER.—Ordered that a copy of the above despatch, and its enclosures, be forwarded to Local Governments and Administrations, and that attention be invited to Mr. Cobb's remarks, in the discussions printed at the end of Mr. Wardle's lecture, regarding the extensive demand for tasar silk, and to Mr. Clayton's observations about the uses of tasar and other wild silk waste.

Government of Madras.
Do. of Bombay.
Do. of Bengal.
Do. of the North-Western Provinces and Oudh.
Do. of the Punjab.
Chief Commissioner of the Central Provinces.
Do. of British Burma.
Do. of Assam.
Do. of Coorg.
Resident at Hyderabad.

(True Extract.)

(Signed) E. STACK,
Offg. Under Secy. to the Govt. of India.

ENCLOSURE No. 1.

From Her Majesty's Secretary of State for India, to His Excellency the Right Honorable the Governor-General of India in Council, dated India Office, London, 11th December 1879, No. 144 (Statistics and Commerce).

ABSTRACT.—Forwarding a copy of letter from Sir P. C. Owen, relative to a collection of the wild silks of India presented by Mr. Thomas Wardle to the Chamber of Commerce of Lyons, &c.

No. 250. ORDER THEREON, 28th April 1880, No. 479.

Miscellaneous.

Col. Beddome.
Dr. Bidie.
Dr. Short.

ORDERED that the above extract with a copy of the Secretary of State's despatch and its enclosures be communicated to the Board of Revenue, the Chamber of Commerce, the Trades Association and other gentlemen marginally noted, in continuation of G.O., 26th March 1879, No. 729.

(True Extract.)

Secretary to Government.

To the Board of Revenue.
,, the Chairman, Chamber of Commerce.
,, the Chairman, Trades Association.
,, Colonel R. H. Beddome.
,, Surgeon-Major G. Bidie, M.B.
,, Dr. J. Short.

Exd. W. H. French.

60 Copies.

Recd: No. 1753

179
2

To

THE SECRETARY TO THE GOVERNMENT OF INDIA,

DEPARTMENT OF *Home, Revenue*
and Agricultural
Dept.
Calcutta

SIR,

I am directed to request that you will be good enough to furnish this Department with as many number 100 copies of the report on the collection of ^{the} wild silks of India forwarded with your ^{letter} ₇₄₋₈₄, dated 13th Feb^y 1880.

Noted and
17 May 1880.

List from orders
in Council

G. W.

Special

G.O. of 118301 27th June.

REVENUE.

THE BOARD OF REVENUE,

Secy, Secy, Secy.

Dated April 25th.

Printed by M.


G.O. 25th April 1880, No. 479.

A P R I L.

Nos. 249-250.

IT is Sd/- to India—Communicating to the
Board of Revenue the Chamber of Commerce,
No. 25, papers relating to the

2/6/80



Vide cat 3769/80.

Handwritten note:
See serial
11 600

Note on Co. O. 28 April 1880 No 479. A29

*
original
available
serial 11
been put
into folder
When sending up the bundle of ϕ O's to be marked for the Editors' table, I put ~~up~~ a slip on this stating that there was no copy of Mr Wardle's lecture available for that purpose or to send to Mr Garton as directed.

Now the Board have sent in an application for 80 copies of the ϕ O for communication to Comm. The ϕ O containing merely the stereotyped order can be of no use without its envelope, (the lecture referred to), and as the

2/1/80

✓

Letter is of general interest may it
not be printed as suggested by me?

11/15/80

1279/5

INDIA OFFICE;
London, the 11th December 1879.

Statistics & Commerce.
No. 144.

To His Excellency the Right Honourable the GOVERNOR GENERAL
of India in Council.

MY LORD,—I forward herewith, for the information of Your
Excellency, a copy of a letter* from
Sir Philip Cunliffe Owen, relative
to a collection of the wild silks of India presented by Mr. Thomas
Wardle to the Chamber of Commerce of Lyons, and also a copy of a
lecture delivered by the latter gentleman on this subject at the
Society of Arts in May last.

* Dated 18th November 1879.

I have the honour to be,

MY LORD,

Your Lordship's most obedient, humble Servant,

CRANBROOK.

Dated No. 2, The Residences, South Kensington, the 18th November 1879.

From—SIR PHILIP CUNLIFFE OWEN.

To—Her Majesty's UNDER SECRETARY OF STATE for India.

WHEN I had the honour as acting as representative for the Indian section of
the Paris Exhibition of 1878, I received an urgent request from the Chamber
of Commerce of Lyons that I would endeavour to procure a collection of the wild
silks of India. The interesting exhibition prepared for the India Office by Mr.
Thomas Wardle deservedly attracted universal attention. The International
Jury, to show the value of this contribution made by the Government of India,
unanimously awarded to His Excellency the Viceroy the Grand Medal of Honor
for silks as the highest distinction in their power to bestow.

I communicated to Mr. Wardle the request made by the Lyons Chamber
of Commerce, and he begged for the permission to provide the collection at his
own expense. Therefore Mr. Wardle informs me that this collection is now ready
for transmission to Lyons. It contains a complete series of moths, cocoons, raw
silks, thrown silks, woven, dyed, and undyed silks, and either printed from Native
blocks in Native colours, or in gold and silver.

I have therefore the honour to submit, for the consideration of the Secretary
of State for India, that a communication should be transmitted to the Secretary
of State for Foreign Affairs, informing him of Mr. Thomas Wardle's gift, in order
that the same may be brought before the French Government for presentation
to the Lyons Chamber of Commerce.

I take this opportunity of bringing under the notice of the India Office
the lecture delivered by Mr. Wardle at the Society of Arts which has been
reprinted with illustrations, and in doing so I may express a hope that the subject
may be considered of sufficient importance to be brought under the notice of the
Government in India.

One result of the Exhibition in Paris has been that tussier silk was on sale
in the market previous to the Exhibition at 2.6 per lb., and now the averaged
value per lb is 7, and the silk in Europe is getting scarce from the increasing
demands for it, chiefly from France.

ON THE WILD SILKS OF INDIA, PRINCIPALLY TUSSER.

INTRODUCTION.

This interesting and important subject has received so much learned attention during the last fifty years, from both entomologists and sericulturists, that it is impossible now to treat of it in a lecture without saying much that is not absolutely new.

Whilst, therefore, abstract research may have but a narrow field left, I hope that what I have to say may be useful in stimulating a greater utilisation of these products, which are as beautiful as they are curious, and in calling the attention of manufacturers, printers, dyers, and users to some most important improvements and developments in each of their several departments, the result of a lengthened study of the exact nature of the fibres, and of new and improved modes of manufacturing and decorating them.

Amongst the many names of persons in various countries interested in the cultivation of wild silkworms, the utilisation of their products, and in the entomology of the subject, I venture to give the following list, which will be found interesting by many, and will serve to show more forcibly to what a large extent this important subject has been made a matter for study and investigation:

English.

Dr. Roxburgh, Captain Hutton, Mr. Hugon, Dr. Birdwood, Mr. F. Moore, Mr. P. H. Gosse, Captain Mitchell, Mr. Butler, Major Couesmaker, Lady D. Neville, Lady Gilbert, Mr. Culvert, Mr. Geoghegan, Dr. Alex. Wallace, Dr. Mackenzie, and others.

Continental.

M. Guérin Méneville, M. Personnat, M. Robert, M. Camille Mayne, M. W. Reed, M. Braine, M. Maurice Girard, M. Wailly, Wilhelm Carl Berg, M. Costa, M. Matthieu Bonagous, Mich. Judizky, Rob. Van Westmas, N. H. de Graaf, Dr. Chavannes, and many others. My friend M. Rondot, Member of the Chamber of Commerce, Lyons, and President of the Jury Class 34, Silks, at the Paris Exhibition last year, author of several important works on silk and Eastern dyes, is at the present time giving much economic attention to this question, for the purpose of considering supply for the increasing demand of the French silk trade in wild silks.

Amongst those who have interested themselves especially in the utilisation of these wild silks must be mentioned Dr. Birdwood, who was one of the first, if not the first, to call public attention to the importance of tussier silk, in a lengthy report to Government in 1838, which I regret I have not time to read.

Lady Dorothy Neville is well known by her efforts in trying to acclimatise the *attacus* worm, and in utilising its silk during a long series of observations of a practical nature.

Mr. Geoghegan is the author and compiler of a valuable report to the Government of India on the silk industry of India, which is indispensable to a study of the subject.

Major Couesmaker is pursuing with a praiseworthy determination, at Poona, his culture of tussier silkworms, with a view of making tussier silk as systematic an industry as that produced by *Bombyx mori* and the other mulberry silks.

Besides many others, the enthusiastic labours of the French, aided by their Société d'Acclimatation, set us a great example. They have succeeded in domesticating, if not in almost naturalising, several Eastern species of wild silk producers, and in obtaining from them silk of industrial value.

Classification.

The silk-producing *Lepidoptera*, insects are of many species, possessing, as is here shown, very marked structural differences, whilst the variety and quiet beauty of their colours and, with the exception of the mulberry feeders, their large size, contribute greatly to the charm of studying this branch of natural history, and they make a collection, apart from their great usefulness, worthy of being placed in the first rank.

They belong, as I have stated, to the order *Lepidoptera*, and are all members of but two families, *Bombycidae* and *Saturniidae*.

All the *Saturniidae* are silk spinners, but not all the *Bombycidae*.

The British Museum catalogue contains the names of 294 species of *Saturniidae*. Mr. Butler, of the British Museum, informs me there have been 100 more species added since the publication of the catalogue.

The following table shows the position of these two families in the great system of classification of the animal kingdom:

DIVISION III.

Articulata.

SUB-DIVISION II.

Anthropoda (or true articulata).

CLASS VIII.

Insecta.

SUB-CLASS III.

Metabola.

ORDER X.

Lepidoptera.

SUB-ORDER I.

Heterocera (Moths—6 groups or tribes).

GROUP.

Bombycina.

FAMILY 10.

Genera.—*Bombycidae*.

Bombyx,
Theophila,

Ocinara,
Trilocha.

FAMILY 8.

Genera.—*Saturniidae*.

Attacus,
Antheraea,
Actias,
Salassa,
Rinaca,
Rhodis,

Caligula,
Neoris,
Saturnia,
Loepa,
Cricula.

Mr. Frederic Moore, Curator of the India Museum, the acknowledged authority on Indian wild silk-moths, has kindly furnished me with a list of all the silk-producing *Lepidoptera* of India known at the present time.

The India Museum contains a collection of most of the wild silk moths, with specimens of the cocoons and the silk of their larvae, arranged by Mr. Moore, a description of which will be found in the second volume of the catalogue on Indian *Lepidoptera*, published in 1858-59.

To illustrate my lecture, I have here to-night a very interesting collection of the moths of the greatest industrial importance, as well as the cocoons of their larvae, and examples of their silk in most, if not all, the various stages of industrial development.

The following is Mr. Moore's list of all the known species of silk producers in India. It is the most valuable list yet published, and shows how rich India is in silk-producing insects:

MULBERRY-FEEDING SILKWORMS—DOMESTICATED.

- Bombyx mori* (Linnaeus).—The common silkworm, domesticated in China, Bokhara, Afghanistan, Cashmere, Persia, S. Russia, Turkey, Egypt, and Algeria, Italy, France, and Spain, in all which countries it produces but one crop annually, spinning the largest cocoon and the best silk, of a golden yellow, or white.
- Bombyx textor* (Hutton).—The *Boro poolloo* of Bengal, domesticated in S. China and Bengal; an annual only, producing a white (sometimes yellow) cocoon, of a different texture and more flossy than *B. mori*.
- Bombyx sinensis* (Hutton).—The *Sina*; *Cheena*, or small Chinese monthly worm of Bengal, partially domesticated in Bengal, where it was introduced from China: produces several broods in the year; cocoon white and yellow.
- Bombyx crasi* (Hutton).—The *Nistry* or *Madrassee* of Bengal, introduced from China; domesticated in Bengal; yielding seven or eight broods of golden yellow cocoons in the year, of larger size than *B. sinensis*.
- Bombyx fortunatus* (Hutton).—The *Dasee* of Bengal yields several broods annually, spinning the smallest cocoon of a golden yellow colour.
- Bombyx arracénensis* (Hutton).—The Burmese silkworm, domesticated in Arracan, said to have been introduced from China through Burma; yields several broods annually; cocoons larger than the Bengal monthly species.

MULBERRY-FEEDING SILKWORMS—WILD.

- Trilecta Huttoni* (Westwood).—The wild silkworm of the N. W. Himalayas. A wild species, the worms being found abundantly feeding on the indigenous mulberry in the mountain forests of the N. W. Himalayas.
- Theophila sherwilli* (Moore).—The wild silkworm of the S. E. Himalayas.
- Theophila Bengalensis* (Hutton).—The wild silkworm of Lower Bengal. Discovered in the neighbourhood of Calcutta feeding on *Artocarpus lacucha*. Found also at Ranchee in Chota Nagpur.
- Theophila religiosa* (Heller).—The *Jorae* of Assam and *Deo-mooga* of Cachar. Feeds on the bar tree (*Ficus indica*) and the pupal (*T. religiosa*).
- Theophila mandarina* (Moore).—The wild silkworm of Chekiang, N. China. Worms stated to feed on wild mulberry trees, spinning a white cocoon.
- Ocinara lactea* (Hutton).—Mussooree, N. W. Himalayas. Feeds on *Ficus venosa*, spinning a small yellow cocoon, yielding several broods during the summer.
- Ocinara Moorei* (Hutton).—Mussooree, N. W. Himalayas. Also feeds on *Ficus venosa*, as well as on the wild fig, spinning a small white cocoon. It is a multivoltine.
- Ocinara diaphana* (Moore).—Khasia Hills.
- Trilecta varians* (Walker).—N. and S. India.

ATLAS AND ERIA GROUP.

- Atteus atlas* (Linnaeus).—China, Burma, India, Ceylon, Java. This appears to be almost omnivorous, feeding in different districts upon the shrubs and trees peculiar to them. At Mussoorie it is found upon *Bradleya ovata*, *Pulconeria insignis*, and several other trees; at Almora the yellow flowering barberry is said to be its favourite food. In Cachar it feeds on various other trees. Cocoon well stored with a fine silk.

- Allium afflicta* (Heller).—Sikkim.
Allium Eburnia (White).—Sikkim, Cherra, and Khasia Hills.
Allium erythra (Drury).—China. Domesticated in the Provinces of Shantung and Honan. Grows on the varnish tree (*Alseodaphne glandulosa*).
Allium Khasi (Jones).—The *King* of Assam, and *Arindi* of Dinagepur. Domesticated in the Northern parts of Bengal (Bogra, Rangpur, and Dinagepur), in Assam and Cachar, feeding on the castor-oil plant (*Gliricidia sepium*), yielding a species of onion of 1½ inches. Grows somewhat loose and fleshy, orange red, and is eaten raw. The seedling is *Alsatia* *arundinacea* of Europe—the result of a cross between *A. sativa* and *A. arundinacea*.

fertile hybrid between the Chinese and the Bengal species, was produced some years ago in France, by Monsieur Guérin Méneville, and subsequently reared, from whence it was introduced into various parts of the world.

Attacus Canningi (Hutton).—N. W. Himalayas. Common in a wild state, feeding on the leaves of *Coriaria nipalensis* and *Xanthophyllum hostile*. Cocoons hard and compactly woven, rusty orange or grey. An annual.

Attacus lunula (Walker).—Sylhet.

Attacus obscurus (Butler).—Cachar. Not very common. Stated to feed on a plant called *lood*.

Attacus Guerini (Moore).—Eastern Bengal.

ACTIAS GROUP.

Actias Selene (McLeay).—Mussooree, Sikkim, and Khasia Hills; Madras. The worms feed upon *Andromeda ovalifolia*, *Coriaria nipalensis*, wild cherry, and walnut, at Mussooree, and on *Odina nodier* in Madras.

Actias Sinensis (Walker).—N. China.

Actias Leto (Doubleday).—Sikkim and Khasia Hills.

Actias Maenas (Doubleday).—Sikkim and Khasia Hills.

Actias ignescens (Moore).—Andaman Isles.

TUSSEE AND MOONGA GROUP.

Antheraea mylitta (Drury); *Antheraea paphia* of authors; the Tusser, Tussar, or Tussah silkworm.—These well known and valuable insects (of various undetermined species) are widely distributed over India, from East to West and North to South, on the coast, and in the Central Provinces. They feed in a wild state upon the ber (*Zizyphus jujuba*), the asun (*Terminalia alata*), the scemul (*Bombax heptaphyllum*), &c.

Antheraea mezankoorie (Moore); the *Mezankoorie* silkworm of the Assamese.—The worms which produce the mezankoorie silk are stated to feed on the addakoory (? *Tetranthera* sp.), which is abundant in Upper and Lower Assam. The silk is nearly white, its value being fifty per cent. above that of the moonga.

Autheraea nebulosa (Hutton).—This is the *Tusser* of the Sonthal jungles of Colong. It is also found in Singbhum, Chota Nagpur.

Antheraea Perrotteti (Guér. Mén.).—Described as being found in the districts of Pondicherry, feeding upon a species of *Zizyphus*, the jambool (*Syzygium jambolanum*). &c. Stated to produce four broods in a year.

Antheraea Andamana (Moore).—An allied species to the tusser. Inhabits the S. Andamans.

Antheraea Frithi (Moore).—Sikkim Himalayas. A common species, inhabiting the hot sub-tropical valleys below 2,000 ft. Known only as a wild species. The cocoon is stated to be similar to that of the tusser in form, but of finer silk.

Antheraea Helferi (Moore).—Sikkim Himalayas. This is a common species found in the hot valleys of Sikkim.

Anthracæ Assama (Helfer).—The *Moonga* or *Mooga* of the Assamese. The moonga silkworm feeds upon the trees known in Assam as the champa (*Michelia* sp.), the soom, kontoolva, digluttee (*Tetranthera diglottica*), the pattee shoonda (*Laurus obtusifolia*), and the sonhalloo (*Tel. macrophylla*). It is extensively cultivated by the Natives, and can be reared in houses, but is fed and thrives best in the open air and upon the trees. The silk forms an article of export from Assam, and leaves the country generally in the shape of thread.

Antheraea Roylei (Moore).—The oak-feeding silkworm of the N. W. Himalayas. A common species, feeding on the bill oak (*Quercus incana*) of the N. W. Himalayas (Simla, Mussooree, Almora). The cocoon is large and very tough, the silk being pronounced as promising, and worth cultivating. They can be reared easily in the house.

MISCELLANEOUS GROUP.

Salassa Lola (Westwood).—Sikkim Himalayas.

Rinaca Zuleika (Hope).—Sikkim.

Rhodia Newara (Moore). Nepal (Katmandoo). Worms feed upon a species of weeping willow. Spins a brilliant green cocoon, pendant from the twigs.

Caligula Thileta (Westwood).—Mussooree, N. W. Himalayas, 7,000 ft. Common, the worms feeding on *Andromeda ovalifolia*, wild pear, and the cultivated quince, forming a light, open, net-like cocoon.

Caligula Simla (Westwood).—Simla, N. W. Himalayas, 5,000 ft. Feeds on the walnut (*Julia balyonica*), wild pear, &c.; forms an open, net-like cocoon.

Caligula Cachara (Moore).—Cachar.

Neoris Huttoni (Moore).—Mussooree, N. W. Himalayas, 6,500 feet. The worms appear in April, feeding upon a species of wild pear tree; spins a thin silken cocoon.

Neoris Shadulla (Moore).—Yarkund.

Neoris Stoliczka (Felder).—Ladak.

Saturnia Cidesa (Moore).—Hot valleys of the Sikkim Himalayas.

Saturnia Grotei (Moore).—Sikkim Himalayas.

Saturnia Lindia (Moore).—Sikkim Himalayas.

Saturnia Anna (Moore).—Sikkim Himalayas.

Loepa katinka (Westwood).—Sikkim, 5,000 to 7,000 feet. Assam.

Loepa sikkima (Moore).—Hot valleys of Sikkim.

Loepa sivalica (Hutton).—Mussooree, 5,000 feet. Spins a long cocoon, pointed at each end, and of a dark greenish-grey colour.

Loepa miranda (Moore).—Sikkim Himalayas.

Cricula trifenestrata (Heller); the *Hampiltonia* of the Assamese.—Noted as being very common in Assam, the worms feeding on the soon tree, forming an open net-like cocoon of a beautiful yellow colour and of a rich lustre, the silk being spun in the same manner as the Eria cocoon. Occurs also in Moulmein, where the worms are stated to feed upon the cashew-nut tree (*Anacardium orientale*).

Cricula drepanoides (Moore).—Sikkim.

To this number may be added a few others which, although not of India, are well worth the attention of the Government of India for the purpose of acclimatisation there.

Antheraea Pernyi (Guér. Mén).—The oak-feeding silkworm of Mantchouria, N. China. This is described as having been long known to the Mantchour Tartars, very large quantities of the silk being used among the Chinese. The worms feed on various species of oak (*Quercus mongolica*), &c., the cocoon differing from the tusser in form and texture. The silk is represented as strong, but with little lustre. Two crops of silk are produced in the year—a spring and autumn crop.

Antheraea Confucii (Moore).—A species allied to *A. Pernyi*, inhabiting the hills in the neighbourhood of Shanghai, N. China.

Antheraea Yama-mai (Guérin Méneville).—The *Fama-mai* silkworm of Japan.

An oak-feeding species, forming a cocoon of a pale yellowish-green colour. This worm feeds on the oak, and produces excellent silk of considerable commercial value in Japan. I should strongly recommend its introduction into India. It has been acclimatised in Europe, and crossed with *Bombyx Attacus—Pernyi*, is successfully reared in France, the eggs hatching at almost freezing point. The silk is much cultivated and used in Japan. Its fibre is oval, and 950th of an inch thick.

Saturnia pyretorum from South China.—The worm feeds upon the *Liquidamber formosana* in Canton, Amoy, where the silk is stated to be woven into a coarse fabric.

Neoris Shadulla (Moore).—Yarkund.

Theophila mandarina (Moore).—N. China.

HISTORY OF SILK.

A few words on the silk of commerce and its history may form a fit preface in introducing to your notice those wild silks which it is my object to describe to-night.

The name silk is derived from the name of the people of Eastern Asia, whom the ancient Greeks called Σῆρες, Seres, and who, no doubt, were the Chinese, and who were then, as now, celebrated for silken fabrics or seric stuff.

From seres comes the Latin *sericum*, the French *soie*, the German *seiden*, the Anglo-Saxon *seole*, the Icelandic *silki*, and the English *silk*.

We are informed by Hawac-nan-tze, in a Chinese work called the "Silkworm Classic", that Teling-she, the principal queen of Hwang-te, B. C. 2640, was the first to rear silkworms, and the Emperor Hwang-te was induced to invent robes and garments from this circumstance.

The queen and wives of the nobles through successive generations personally attended to the rearing of the silkworms. Alas! that, in these days of expediency, hurry and greed, the faith in machinery should have so disastrously stifled the better faith in manipulation, both in silk handicraft work and the old love for the spinning wheel, to the detriment alike of all classes, from the noble to the low born, who have been robbed of much that in bygone days contributed to the comfort, occupation, and enjoyment, especially of the gentler sex. Surely here is a worthy object of reform for our national art schools. That this silk was of the mulberry-fed kind is evident from a further extract from the same work, which says that afterwards, "when Yu regulated the waters B. C. 2200, mention is made in his work on the tribute of the land adapted for the mulberry tree having been supplied with silkworms, from which time the advantage thereof gradually increased".—Horsfield and Moore's Catalogue, p. 377.

It is not known whether silk was utilised in India at so early a period as this, probably not, but that India learned the art from China is generally understood, although at what period is not known.

For more precise information respecting the Westward spread of silk culture, I would strongly recommend my hearers to read Dr. Birdwood's account in his Handbook to the British India Section of the Paris Exhibition. This learned history of silk will be read with much pleasure by all who can see in silken stuffs something more than a mere commercial value. The account of its utilisation and spread from East to West is described with almost the charm of romance. Its development is traced from its earliest days in the East to its introduction at last into our own country in the reign of Henry VI., and again to those sad French times of the persecution of the religious Huguenots by Louis XIV., which drove their silk-workers by a happy tide to our shores. How that tide has in our own times returned to France, and carried with it not the workers, but the industry, I leave for statesmen—and they have much to answer for—and manufacturers to think over and retrieve.

Europe may be said to have got hold of the silk industry by a fraud; two monks are said to have brought away the eggs from China concealed in their walking canes. A similar account is well related by Dr. A. Wallace of the way the eggs of the prized Yama-mai silkworm were abstracted from Japan by a young Japanese, who obtained them at the instigation of his European tutor at the risk of his life, for this was an offence there punishable by death. It would be but a small return for the benefits we have obtained to ourselves by these frauds if we could teach those weaker peoples the benefits of the better making, and, above all, the more friendly interchanging of stuffs and commodities. It is with this hope that I put the wild silk question before the Society of Arts, that, having India, with its extensive wild silk regions, in our possession, we may, by gentle means, teach the Natives to improve the culture and preliminary stages of its manufacture; so that it may be brought from them in a state fit to be used by us for all the purposes of which, in its improved state, it is really capable, and, to use the words of Sir Louis Mallet in the first letter of instruction I received from the India Office, that "a new and very profitable industry may be founded in India".

The natural history of every kind of silk may be briefly stated to be this: From a small egg laid by the moth, of whatever species, appears in due season a small larva, or caterpillar, or worm, as it is usually called. This worm, after having lived its day, feeding only on the leaves of certain plants characteristic of its species, spins, or rather secretes, a fine thread of silk around itself, for covering and protection during the time it lies dormant in the next stage of its existence. As soon as it has secreted all the silk, it changes into a pupa or chrysalis, and remains inside its silken cell until the time comes for its appearance as an imago, or perfect insect, having four scaly wings, legs, and antennae. When its hibernation is ended, it emits a fluid which softens the end of its cocoon cell, and by means of its wing spines and legs, parts the fibres aside until the opening is large

enough for it to creep out. After a short time, its wings dry and expand, and it has entered into its perfect state. It lives only a few days in this phase of its existence. It is in this stage only that the race is perpetuated, the female laying a number of eggs, and dying soon afterwards.

BOMBYX MORI.

What is generally known as the silk of commerce, in both ancient and modern days, is distinguished from all others by the singular circumstance of the larva which produces it feeding on the leaves of the mulberry tree. The moth is therefore named *Bombyx mori*.

It must not be inferred, however, that there is only one species, as is too generally thought. Mr. F. Moore assures me there is sufficient structural variation in moths, which were at one time understood as *Bombyx mori*, to constitute specific differences. The following species feed on the mulberry, and their silk is in ordinary and promiscuous use along with that of the *mori*;

Bombyx sinensis (Hutton),

Ditto *croesi* (ditto),

Ditto *fortunatus* (ditto),

Ditto *arracanensis* (ditto),

Ditto *textor* (ditto),

and no doubt several others not yet defined, but whose silk varies very much, such as that called Canton, and that called Tsatlee, although both are from China, as well as the silk from Japan, which has a thicker fibre than the *Mori* silk of other countries, being $\frac{1}{16}$ of an inch thick, as against $\frac{1}{100}$ of an inch of the *Mori* silk of Italy. The strength and tension of this and the wild silks, as well as the respective sizes of their cocoons, are given in the table of microscopic measurements which I have carefully revised for my lecture, confirming generally, but in several instances correcting those I made for my monograph for the Paris Exhibition.

I have here some beautifully-prepared Italian silks of *B. mori*, kindly lent to me for my paper by my friend M. H. Meyer, a large silk manufacturer at Milan.

The eggs of *Bombyx mori*, and the other species of *Bombyx*, form a most important article of commerce between Japan and Europe. They are imported from thence into Italy and the South of France annually, to supply the deficiency caused by the silkworm disease known as *pebrine*, and also to infuse, as it were, new blood. The value of the eggs is about 25s. per ounce. Twelve pounds of cocoons are required to produce 1lb. of silk. The value of the raw silk of Bengal in London is now 14s. 6d. per lb. The colour is generally of a rich golden yellow of this sample, but it is also sometimes white. In China and Japan it is all white, with rare exceptions.

WILD SILKS.

As far as I can learn, the only species of cocoons at present utilised in India, besides those of the several species of mulberry-feeding worms of the genus *Bombyx*, are the wild ones of the following species:

Attacus ricini,

Ditto *cynthia*,

Ditto *atlas*,

Ditto *edwardsia*,

Antheraea assama, or Moonga worm,

Ditto *paphia*, or Tusser worm,

Ditto *perotteti*,

Ditto *nebulosa*,

Ditto *roylei*,

Ditto *frithii*,

Ditto *mezankooria*,

Cricula trifenestrata,

and, perhaps,

Attacus selene and *Antheraea helferi*.

The India localities of the four principal ones, *Ricini*, *Atlas*, *Moonga*, and *Tusser*, are shown on these maps by various colourings.

THE ERIA SILKWORM.

We now arrive at the consideration of the *Eria*, *Arindy* or *Arundi* worm, of which there are two species, *Attacus ricini* and *Attacus cynthia*; *Attacus ricini* (Boisd.) is known also under the names *Saturnia ricini* (Boisd.) and *Attacus lunula*, (Walker).

Attacus ricini is a worm which feeds on the *Palma christi*, or castor-oil plant (*Ricinus communis*). It is a native of Assam, and, according to Mr. Geoghegan, is found to the South-West in a track comprising the districts of Nepal, Kumaon, Ladak, Darjeeling, Dinagepur, Rungpur, and, perhaps, parts of Bhagalpur, and Purneah, and in Assam. In 1791 Sir W. Jones drew attention to this silk, and Dr. Roxburgh in 1801.

The leaves of the castor-oil plant, *Ricinus communis*, or *Palma christi*, are the best and commonest food, but it will also feed on the following: *Koosool*, *Hindu* grass, *Murkundal*, *Okonnee*, *Gomaree*, *Litta pakoree*, *Birzonally*, *Xanthoxylon hostile*, *Coriaria nipalensis*, *Ailanthus glandulosa*, *Ailanthus excelsa*.

Attacus ricini is of a larger size than *Attacus cynthia* owing probably to its freedom, and there is no doubt it is a silk-producer of great importance in India. Probably the most complete account of it is by Mr. Geoghegan in his report on the silk industry of India, of which I avail myself to gather some useful particulars.

The *Attacus ricini*, according to Dr. Buchanan, feeds both on *Ricinus communis* and *Ricinus viridis*.

The female moth lays her eggs round a twig, and then dies. These twigs are sold in the markets covered with eggs, the dead moths frequently hanging to them, and presenting a very curious appearance.

The caterpillar moults four times, and when fully grown is about 3½ inches long.

According to Mr. Hagon, the Natives soften the cocoon in potash, and draw the silk off roughly with the finger and thumb, thus making a kind of spun silk. Dr. Buchanan, however, says the silk is wound on a reel in Dinagepur.

M. Guérin Méneville, however, stated in 1860 that it was impossible to reel this cocoon. I am certainly disposed to think it is quite impracticable. Still there is a method of using it at hand, and I know of no silk better adapted for spinning.

Some of the cocoons I received from the India Office for examination I have had carded or dressed, and the result is, as is shown in this sample, a fibre of great fineness and length, which would be eagerly sought after by English and Continental spinners, if the cocoons could be collected and sent over in quantity. Owing to the very small quantity of cocoons in my possession, I am sorry I have not been able to have specimens of yarn or cloth made for my lecture, but I have here Native-spun yarn undyed, bleached and dyed, also Native-woven cloth in the undyed and dyed states, as well as specimens of prints executed on it, for, I believe, the first time in its history or use. The cloth is beautifully soft.

I find no difficulty in making this silk yield to tinctorial influences, either in dyeing or printing, and I have no hesitation in predicting a great future for it as soon as it is subjected to the better spinning appliances of Europe, and I am recommending the Government of India to give all possible encouragement to the importation of *Eria* cocoons into Europe.

Dr. Helfer in 1837 stated in the *Journal of the Asiatic Society of Bengal* that this worm is so productive as to give sometimes twelve broods of silk in the course of the year. The worm grows rapidly, and offers no difficulty whatever for an extensive speculation.

The industry of the natives of India should be stimulated to the gathering in of all kinds of wild silk cocoons, whether windable or not, for there is no doubt that those kinds which cannot be reeled can be most easily spun, and there is at the present moment a request on the part of silk spinners for a much larger supply of Tusser silk cocoons and Tusser silk waste for spinning purposes, and no doubt the other wild silk cocoons would be gladly bought up, especially those of *Attacus ricini*.

Mr. Geoghegan says, with regard to the thread from cocoons of *Attacus ricini*, that a seer of 96 sicca weight ($2\frac{1}{16}$ lbs.) of this thread is worth from annas 12

(one shilling and sixpence) to Re. 1 (two shillings), but it is very seldom sold, and the people who keep the insect in general rear no more than is just sufficient to make cloths for their own families. The cloth lasts very long, owing to which quality it is probable that some use might be found for this material in our manufactures at home.

Mr. Michael Atkinson, of Singapore, describes the cloth made of Eria to be of incredible durability, the life of one person being seldom sufficient to wear out a garment made of it, so that the same piece descends from mother to daughter. The thickness of this fibre is $\frac{1}{16}$ of an inch on the outside of the cocoon, and $\frac{1}{16}$ in the inner part. (For other particulars see Table.)

Mr. Hugon says that "in Assam, the quantity of Eria cloth the merchants formerly used to take away was very considerable, but in the latter years of the Assam Rajah's rule, from the disorganized state of the country, the number of merchants gradually diminished. The quantity the country is capable of exporting under an improved management would be very large, for it forms at present the dress of the poorer classes at all seasons, and is used by the highest for winter wear. So long ago as 1769, vast quantities were being produced in the country around Guraghaut". He estimated the annual production at 1,000 maunds, or 82,000 lbs. "In the district of Darrung, the annual yield of Eria is 1,000 maunds, of which one-third is exported either in the form of cocoons or woven into heavy cloths (Borkapor)." Mr. Hugon stated that, for want of a proper solvent of the gum, the Natives could not reel the cocoons. Mr. Brownlow states that in Cachar the Eria or Ricini worm is trained by the Cacharis, a people living in isolated villages on the hills. They soften the cocoons in a mixture of cowdung and water, and they are then carried off to a spindle by the women of the tribe. This silk is dyed by the Natives with lac, mungeet, and indigo.

Table of the diameter, strength and tension of a single fibre and dimensions of cocoon of the chief mulberry and Indian wild silks. (continued)

[illegible]

10

ATTACUS CYNTHIA.

- Attacus cynthia* (Drury).
Phalaena attacus cynthia (Drury).
Phalaena cynthia (Roxburgh).
Bombyx cynthia (Oliver).
Samia cynthia (Hübner).
Saturnia cynthia (Westwood).
Saturnia arrandi (Royle).
Eria of Assam (Hugon).

I have mentioned the close relationship of *Attacus cynthia* to *Attacus ricini*.

Mr. Geoghegan says *Attacus cynthia* occurs in a wild state throughout a great part of the Himalaya, in the Dehra Dun, and in Assam and Cachar.

In China, *Attacus cynthia* feeds on *Ailanthus glandulosa*. During the last 20 years much study has been bestowed on the species in Europe, and it has been successfully domesticated in France, and later on in England, so much so, that this study rejoices in the name ailanthiculture, and the breeding-houses of the insect are called ailantheries.

It is to M. Guérin Méneville, in France, and to Lady Dorothy Neville, in England, that we are indebted for leading the way to the successful rearing of the *Attacus cynthia*. A description of their labours will be found in a memoir by Dr. Wallace, published in 1866, in the *Journal of the Entomological Society*. Besides giving full particulars of all that relates to ailanthiculture, Dr. Wallace describes with great minuteness his own experience in producing this silk. He received the prize offered by the Council of the Society in 1865 for an essay on "Economic Entomology".

The diameter of all the fibres in the cocoon is tolerably uniform, and about $\frac{1}{100}$ inch. The strength of the outer fibres is, however, only $2\frac{1}{2}$ drams, that of the inner being $2\frac{1}{2}$ drams. The tension of the outer fibres is one inch to the foot, and of the inner $1\frac{1}{2}$ inches. All the fibres are flat and longitudinally striated, and united in pairs by their edges. The cocoon is about the size of a thrush's egg, $1\frac{1}{2}$ to $2\frac{1}{2}$ inches long, and $\frac{3}{4}$ of an inch in diameter, and consists of very fine silk, rather loosely, but closely laid together. It offers great difficulty in winding, and I am not aware that this has yet been successfully accomplished, I mean commercially.

In 1864 M. Forgemol informed the Imperial Society of Acclimatisation, in France, of the particulars of a mode he had invented to reel this silk. Dr. Wallace has quoted this report, describing the method, in his memoir. I fear, however, it is not now in practical use. But whether the reeling has ever been done, or ever will be, or not, there is no doubt that a great future remains for this silk, now that spinning machinery has been so perfected. All the cocoons that could be bred would be easily brought up by the silk spinners. The thickness of a single inner cocoon fibre is about $\frac{1}{100}$ of an inch, and an outer fibre about $\frac{1}{100}$ of an inch. Throughout the cocoons are a few pairs of fibres $\frac{1}{100}$ of an inch in diameter. The breaking strength of a single fibre is 75 grains, and the tension is about two inches for every twelve inches, i. e., a fibre twelve inches long stretches to fourteen inches before it breaks.

Attacus cynthia came originally to India from China. In 1856 cocoons were sent by Abbe Fantoni from the Province of Shan Tung in the North of China.

Successful breeding took place from these cocoons in Turin in 1857, and from these spring all those which have been since cultivated in France, England and other European countries. Mr. Moore was the first to rear them in England in 1859.

Dr. Wallace gives a lengthened account of the tree on which *Attacus cynthia* best thrives, *Ailanthus glandulosa*, from which I extract the following interesting remarks:

"The genus *Ailanthus* belongs to the family *Rutaceae*, and to the tribe *Xanthoxyleae*, or yellow woods. It is a large and beautiful tree, reminding us some-

what of the walnut. The leaf stalks are of great length; some I have measured were three feet and a half to four feet long, pendant with a graceful curve at the lower third of their length; the leaflets are large, some measuring eight and a half inches in length by three in breadth, from thirteen to seventeen on a leaf-stalk; near the base of the leaf the margin is toothed, and at the extremity of the tooth is a little gland, emitting a strong resinous odour, whence the name *glandulosa*; of these teeth and glands there are generally two, three or more on each leaflet. The same odour seems to be communicated more or less to the leaf, also to the larva, and in a less degree to the imago of *Attacus cynthia*."

The pupae of this insect remain in the cocoon through the winter, except in those cocoons which are formed in the autumn, from which they emerge in three or four weeks. The remaining moths emerge in April, and pair in about 12 hours afterwards, and the female begins to lay her eggs almost immediately. The moths do not live long after their exit from the cocoon. About eight days is the duration of their existence in this last and perfect stage. The eggs of one female vary from 150 to 350—30 of them weigh a grain. They hatch in 12 to 18 days, according to temperature. In seven days the tiny caterpillars complete their first stage by moulting, their second in five or six days after, being, up to this time, gregarious; their third moult occurs in another six days, and their fourth and last in six days more. By this time the larvæ are four inches long, and begin to spin their cocoons. *Attacus cynthia* has two broods in the year.

ATTACUS ATLAS.

The next silkworm to which I wish to draw your attention is the *Attacus atlas* of Linnaeus, which is also known by the following names:

- Phalaena-Attacus atlas* (Linn).
Bombyx atlas (Fabricius).
Saturnia silhetica (Helfer).

The imago of this species is the largest of all the silk moths. As you will see in the case, it is a grand insect. It is called in France "Le Géant des Papillons". The largest specimens measure upwards of ten inches in expanse of wing. According to Horsfield, it feeds on the melokka (*Phyllanthus emblica*), a plant known also in India as *Kupu gaga*. Mr. Geoghegan says this silkworm is found in Mussooree on the *Falconeria insignia*, *Bradleya ovata*, and other plants; in Kumaon, on the barberry, where it is abundant, and also Eastward to Cachar. Mr. Moore describes it as almost omnivorous. Mr. P. H. Gosse has just written a most interesting life-history of this species, describing all that is known of it. His paper, from which I quote a few pertinent facts, will be found in the *Entomologist* of February and March last. He says:

"It is a widespread species, ranging over the South and East half of Asia, continental and insular; common on the slopes of the Himalayas, and all through India to the points of both peninsulas; abundant in China; scattered over the isles of the Archipelago, from Java to the Moluccas, to Borneo and the Philippines, a range of 35° of latitude and 55° of longitude."

His efforts to breed the Atlas worm in England are described, and although he has not been very successful, owing to sickness and death in the larval stage, his account is very interesting. Captain Laudy, of Surbiton, has succeeded in obtaining fifteen good cocoons from twenty-four eggs, the larvæ having been fed on the common barberry. Mr. Gosse has fed his on the willow, others on the plum and apple-tree leaves. He states that all the family of *Saturniidae* are very polyphagous.

It is interesting to observe the micaceous or window-like spot on each of the four wings. These, although characteristic of the *Saturniidae*, are almost more largely developed in *Attacus atlas* than in any other species. The French call these vitreous membranes *porte miroirs*.

There are several varieties of *Attacus atlas* which show their divergence from the type, by the variations of these fusiform window-like ornamentations. In some they are single, in others double.

The diameter of the external fibres of the cocoon is very variable, averaging about $\frac{1}{100}$ inch, whilst that of the internal fibres is more uniform, and about $\frac{1}{1000}$ inch. The outer fibres are capable of supporting an average weight of $2\frac{1}{2}$ drams, and the inner $2\frac{1}{4}$ drams. The tension of the outer fibres is one inch to the foot, and the inner $1\frac{1}{4}$ inches. The fibres are flat and longitudinally striated, and united in pairs by their edges.

The egg of *Attacus atlas* is $\frac{1}{100}$ of an inch long. The larvæ moult six times. The cocoon of *Attacus atlas* is from two to three inches long, and about one inch wide at the widest part, and weighs two grammes, or the $\frac{1}{4}$ th of an ounce.

Mr. Gosse says "the colour of the cocoon is a light amber, or drab; its surface (independently of the impress of leaves) roughly granular, scarcely at all silky or floccose, except at the mouth; its substance thin, parchmenty, very firm; the interior very smooth, and even sub-glossy. The upper extremity forms a natural orifice for the exit of the moth, made by the conveyance of a great number of silk fibres, which are left ungummed, and are thus soft and fleshy, the gummed, stiff silk passing up on one side, and contracting into the cord. Thus the cocoon is not closed, like those of *Bombyx mori*, of *Telea*, of the *Antheraea*, but open, like those of *A. cyathia*, of the *Samia*, of the *Saturnia*. As a result of this structure, the exit of the imago leaves no disturbance behind, no witness, no disarrangement of these soft fibres, such as is the case with Yama-mai, Pernyi, and Mylitta".

Mr. Geoghegan states the silk to be difficult to reel, though it reels partially if boiled with vinegar. Captain Hutton says the silk is decidedly good, and Dr. Chavannes, of Lausanne, states it to be desirable of introduction into France. He says the worm is the Fagara of China, where it has long been cultivated.

ATTACUS EDWARDSIA.

In the India Museum there are two specimens of an atlas moth, from Darjeeling, of an intensely dark colour, and possessing sufficient structural divergence from *Attacus atlas* to warrant Mr. Moore's affirming it to be a distinct species. It has been named *Attacus edwardsia*. Mr. Gosse has the following interesting note respecting it:

"As Darjeeling is 7,000 feet above the sea, and has a climate in which rain and snow are abundant in winter, and humidity is constant, it surely would not be difficult to acclimatise this noble form (be it variety or species) in the British Isles."*

THE MOONGA, MOOGA, OR MUGA SILK.

I have here a silk produced from the worm known as the Moonga or Mooga, *Antheraea assama* (Helfer), and *Saturnia assama* (Westwood). It is found in Assam, and also sparingly in the Dehra Dun, and is the next in importance to tussar. Mr. Geoghegan's description of this silk occupies three pages of the Blue Book on the Silk Industry of India, from which I abstract the following particulars:

"The worm that gives the common fawn-coloured moonga silk, when fed on the most common plants, gives a whitish silk when fed on the leaves of other silk. The plants it feeds on are named and estimated as follows:

"No. 1. *Champa* (*Michelia*).—The silk produced from the worm feeding on this plant gives the finest and whitest silk, used only by the rajah and great people, and is called champa patten moonga. The thread is sold at from Rs. 11 to Rs. 12 a seer (11s. to 12s. per lb.)

"No. 2. *Mozankoorie* (called also addakoor).—The old trees are cut down and the jungle about burnt, and the worms are fed upon the tender leaves of the offshoots for one year, when the leaves become too old and hard for the worms. Silk is sold at Rs. 6 to Rs. 7 per seer (6s. to 7s. per lb.)

"No. 3. *Soom*.—This is the common tree of the vicinity; the silk from the worms fed on this gives the finest sort of fawn-coloured moonga. Silk is sold at Rs. 3½ to Rs. 4 per seer (3s. 6d. to 4s. per lb.)

"No. 4. *Sonhalloo* *Tetrathera macrophylla*.—This is also a brown silk of inferior quality. This plant is most common in Dharuppar and about Russa Chokey.

* Gosse's "Life History of *Attacus atlas*," p. 6.

"No. 5. *Digluttee* *Tetrathera digluttea*.—This is also brown silk of inferior quality, but the worms fed on the leaves of this tree increase much in size.

"No. 6. *Pattees* *hoonda* *Laurus obtusifolia*.

"The moonga worm gives broods five times a year, and the cocoon is very large, but thin. I could only obtain silk, the produce of worms feeding on Nos. 3 and 4, and manufactured into cheap cloths for the lower classes.

"In its natural fawn-colour it stands washing much better than ordinary silk, keeping gloss and colour till the last."*

Mr. Geoghegan, on page 114 of his "Silk Industry of India", says—

"The cycle of the insect is thus given:

From emergence from the egg to commencement of cocoon...	30 days.
In the cocoon	20 do.
As a moth	6 do.
In the egg	10 do.
Total	66 do."

In 1873 Colonel Hopkinson, the Commissioner of Assam, gave more modern figures:

"It thence appears that the soom forests (on which the worm is chiefly fed) cover an area of about 34,000 acres, of which about 18,000 are assessed, yielding a revenue of nearly Rs. 28,000 (£2,800). By far the greater portion of the assessed area lies in the district of Sibsagar. The production of the silk is said to employ some 48,000 persons, but it is not their sole calling. The outturn of silk is estimated at upwards of 100,000 lbs. But as it is admitted that the greater part of the silk is reserved for home manufacture, this estimate cannot be regarded as absolutely trustworthy. The price of the yarn per seer (2lbs.) varies from Rs. 5 (10s.) to Rs. 9 (18s.) in the several districts. The small portion exported goes to Calcutta and Dacca. From the former place it is said to find its way, to some extent, to Bhagalpur and Bombay.

"The Silk Committee of the Agri-Horticultural Society reported favourably on some moonga silk sent down by Captain Jenkins in 1839, and expressed their opinion that the article was calculated to become of extensive and valuable use to our home manufactures."*

One acre of land yields 50,000 muga silk cocoons, which yield upwards of twelve seers (24 lbs.) of silk, price Rs. 5 per seer, or 5s. per lb.

From Mr. Hugon's description of the mode of reeling it is evident it is of the rudest kind, and points to a remedy in the improved continental reeling appliances.

The following particulars by Mr. Hugon in 1834 are interesting: The muga silk industry is confined to Assam and some Tipperah villages. The quantity of land planted with food for the muga was 5,000 acres, capable of yielding 1,500 maunds (123,000 lbs.) of silk. This silk formed one of the principal exports of Assam. The average quantity was 257 maunds (21,070 lbs.) valued at Rs. 66,054 (£5,605), leaving the country principally in the shape of thread. He advocates the use of the moonga silk in coloured fabrics, it being easily dyed.

Having only a very small quantity of coarse reeled muga silk at my disposal, my experiments with it have been limited. I find it bleaches well, is very lustrous, and takes the dye freely, better than tussar. Here are specimens of it in the raw, boiled-off, bleached, and dyed states, and also some waste from cocoons for spinning.

The diameter of the fibres of muga silk taken from the external part of the cocoon averages $\frac{1}{100}$ inch, but the external fibres are very variable. The diameter of the inner and less variable fibres is $\frac{1}{1000}$ inch. The outer fibres will break with a weight of $2\frac{1}{2}$ drams on the average, but the inner will support three drams. The tension of the outer fibres averages one inch to the foot, and of the inner $1\frac{1}{4}$ inches. All the fibres are, like tussar, flat and striated, and united in pairs by their edges.

* "Geoghegan, Silk Industry of India," p. 25.

The following table is given by Mr. Hugon showing the nature and prices of the various kinds of cloth made from moonga silk :

Name of cloth.	Size of yards and inches.	Weights.	Price of thread.	Cost of weaving.	Total.	REMARKS.
		lbs. oz.	s. d.	s. d.	s. d.	
Goovias	4 x 50 in.	0 12	3 9	0 4½	4 1½	Dhoties.
Ditto	9 x 1-4 in.	2 0	10 0	1 0	11 0	
Mekla	2½ x 1	0 8	2 6	0 3	2 9	Petticoats.
Rhia	6½ x 1½	1 0	5 0	0 6	5 6	Scarfs.
Gaursha	4½ x 20 in.	0 4	1 3	0 1½	1 4½	Worn as turbans, or round the waist.
Jeonta Bor Capper	6½ x 1½	2 0	4 0	0 9	4 9	Made of the fleece, and worn in winter.

ANTHEREA PEROTTETI.

This is described by Mr. Geoghegan as producing a strong, wiry, and brilliant silk, but requires carding. The larvæ feed, in captivity, on the *Odina rodier* (Roxb.). "Undergoes four moults, and yields four crops a year." Fishing lines of this silk are said to be made in Dinapur.

ANTHEREA ROYLEI.

Is described by the same author as being found at Darjeeling and in the Himalayas from Kumaon to the Punjab, and feeds on *Quercus incana*. The silk is good, but not abundant, and the insect can be domesticated. Yields two or three crops a year.

TUSSER SILK.

I now come to the principal subject of my paper, the tasar silk, called also tusser, tusseh, tussah, tussore. It is the product of the larvæ of the moth *Antheraea paphia*, of Linnaeus. It is known by the following synonyms:*

- Phalæna, Atticus paphia (Linnaeus).
- Bombyx paphia (Fabricius).
- Phalæna paphia (Roxburgh).
- Saturnia paphia (Helfer).
- Phalæna, Attacus mylitta (Drury).
- Bombyx mylitta (Fabricius).
- Antheraea mylitta (Hübner).
- Attacus mylitta (Blanchard).
- Saturnia mylitta (Westwood).
- Bughy silkworm, moth of the Burbhoo hills.
- Kolisurra silkworm, moth of the Mahrattas.

There is but little doubt that this silk has been utilised for many centuries, both in India and China, but I have not been able to find any account of its ancient history.

* Horsfield and Moore's "Catalogue of Lepidopterous Insects", p. 335.

For the history and other particulars of tusser silk I am glad to acknowledge my indebtedness to the report of Mr. Geoghegan. He attributes the derivation of the word tusser to *tusuru*, the Hindustani for shuttle, and states this caterpillar to be the most widely distributed, as well as the most important, of the wild silk-producers of India.

One of the earliest notices of this insect, or of a species very nearly related to it, is given by the venerable Rumphius, who was born at Hanau in 1657, in his "Herbarium Amboinense" (dedicated by him to the East India Company), Vol. III., p. 113. pl. 75. He discovered the larvæ in Amboina feeding on the *Mangium caseolari rubrum* (*Rizophora caseolaris*, Linn.), a plant of the order of *Terbintaceæ*.* He says:

"When I had kept the cocoons for three weeks, a moth came out quite perfect, which was the most beautiful and largest I had ever seen, which biting away (the silk), showed its head, and at the same time drew out with it a little flock of yellow silk; this the moth performed at night. Its body, like all other moths, is a dirty yellow colour, and in length two joints of a finger; it has two downy horns on its head, of a golden hue, and four large wings, of which the two largest are about an inch long and of a golden colour, but a purple line runs through them transversely, and every wing has, as it were, in its middle a window-like eye, which is surrounded by a purple circle, and, as it were, of the transparency of glass."

The larvæ, when fully grown, are about four inches in length; they have twelve joints or articulations besides their extremities; their colour is green, resembling the leaves on which they feed; and they are marked with reddish spots, and a reddish-yellow band running lengthways. They feed on several plants:

- Rhizophora calcularis* (Linn).
- Terminalia alata glabra* (Assam tree).
- Terminalia tomentosa* (the saj tree).
- Terminalia catappa* (country almond tree).
- Tectona grandis* (teak tree).
- Zizyphus jujuba* (ber tree).
- Shorea robusta* (sal tree).
- Bombax heptaphyllum* (Semul).
- Careya sphaerica*.
- Pentaptera tomentosa*.
- Pentaptera glabra*.
- Ricinus communis* (castor-oil plant).
- Cassia lanceolata*.

In six weeks from the time they are hatched they begin to spin their cocoons, which they most curiously suspend from the branches of the trees, by constructing a thick, hard, cord or pedicle of silky matter, which is made to grasp the branches, as seen in these specimens.

Tusser silk is found, as you will see by the map, over nearly the whole of India.

In the Central Provinces, Mr. Geoghegan says, tusser is utilised in Raipur, Bilaspur, Sambalpur, the Upper Godavari, Chanda, Bhandara, Nagpur, Balaghat, Seoni, Chhindwara, Betul, and Narsinghpur. Sambalpur is said to yield 3,500 seers (7,000 lbs.) of silk, Raipur 60,000 (12,000 lbs.), Bilaspur 900 (1,800 lbs.), and Chanda, 22,500 (45,000 lbs.). The silk is woven and used in the Provinces in mixed fabrics of cotton wool and tusser web. But, at any rate, in some districts, muktahs, garments worn by Brahmins after bathing, cholees, women's bodices, and doputtas and dorwas, seem to be made of pure tusser silk.

Captain Brooke says: †

"In Seoni a regularly organised and thoroughly understood industry, from the rearing of the insects to the weaving of silk into cloth, with all its accompanying machinery of trade guilds, money-lenders, &c. This state of things is, in my opinion, no disadvantage, for, in place of having to contend with the difficulties

* Horsfield and Moore's "Catalogue of Lepidopterous Insects," pp. 336-7.

† "Geoghegan, Silk Industry of India," p. 109.

which in India always surround the introduction of anything new or unknown, the demand is all that is necessary to stimulate production to any extent required. Nor is this a figure of speech, for the natural food of the tussar worm is the leaves of the saj, lendeyā and dhowra trees, all of which are found in every part of this district, and are, I believe, common to the whole of Gondwana. These trees are besides propagated with facility, and, as far as the requirement of the silk insect goes, are of rapid growth; hence, if the silk became more known and better valued, and the profits sufficiently attractive, we might witness a development of the culture similar in kind to that which has of late years taken place in the case of cotton. Supposing then a demand to spring up, I am of opinion that the supply would in a very short time amply meet it. The nucleus of no inconsiderable trade now exists, and only awaits the stimulus of high prices. The primary question, whether the product is, or may become, of such value as to occasion a large demand, is one perhaps that more nearly concerns traders than Indian administrations; still, so convinced am I of the value and beauty of the fabric that can be woven from well-reeled tasar, that I would venture to strongly urge Government action in introducing it to the markets of Europe."

The worms require protection from birds and ants, which are their greatest enemies. The first cocoons are made in August, and are sold, after the moth has escaped, to the silk-dealers at 4 to 8 pice (1½d. to 3d.) the hundred. The unpierced cocoons are only sold to rearers as seed, at Rs. 1-8 to Rs. 2 (3s. to 4s.) per hundred.

Captain Brooke says: "In Chanda and Bilaspur, Central Provinces, the rearing of the worms is attended by many ceremonial observances, which begin when the insect leaves the egg, and are not discontinued until the cocoons are gathered and taken to the rearer's house. During the feeding of the worm the Dheemurs lead lives of the strictest abstinence. None of the sex are allowed within a considerable distance of the trees upon which the worms are feeding, and if by chance a woman or impure man passes near the feeding grounds, the trees and worms are sprinkled in the name of Iogni (an incantation of the god Mahade, whom the worms are supposed to represent) with water taken, if procurable, from a running stream, and in which tulsi leaves have been steeped. During the same period the Dheemurs carefully abstain from flesh, fish, or haki as their food, nor do they cut their hair or shave, and carefully deny themselves all ablution. When the cocoons are formed, they collect into a heap, and a goat, pig, or fowl is sacrificed to Mahadeo in his form Iogni, the blood is sprinkled over the cocoons, and, after a bout of liquor, are taken home. On the third day following the Dheemurs shave and resume their normal condition."*

The caterpillars moult five times, at intervals of from five to eight days. When first hatched they weigh but one-fifth of a grain, and are about ½ inch long; but at the end of their larval existence, which is from forty to forty-five days, they have attained a size of seven inches long, one inch in diameter, and weigh about 370 grains. They then begin to spin their cocoons, which are, as you see, of an egg shape, and silvery drab in colour. The silk is all regularly deposited in a compact manner, resembling in thickness and substance the shell of an egg.

The cocoons vary much in size. The largest I have seen are from Sambalpur, and are two inches long and 1½ inches in diameter, whilst smaller ones are not more than 1½ inches long and ¾ inch in diameter. The weight of the large cocoons is, without the pupa and supporting pedicle, 28 grains; the smaller ones 8 grains. I have here an unbroken double thread reeled for me, from one cocoon, by Mr. H. Meyer, of Milan. It weighs 12 grains, and measures 1,332 yards, or a little more than three-quarters of a mile.

Mr. Coussmaker remarks:†

"As a rule, there are certainly two crops in the year; the moths of the first batch come out in about four or six weeks after the first lot of worms (which come out at the commencement of the rains) have spun; those of the second batch remain quiescent until the rains begin again, that is to say, until May. As this entails the chrysalis remaining in the cocoon as long as eight months, exposed to the hottest sun and occasional thunderstorms, the cocoon had need to be made of a hard impenetrable material; so indestructible is it that Rheels and other tribes, which live in the jungles, use the cocoon as an extinguisher to the bamboo tube in which they keep the "falita" or cotton-rope tinder, used by them for lighting their tobacco, and the slow matches of their matchlocks. The cocoon is also cut into a

* Captain Brooke, as quoted in "Geoghegan, Silk Industry of India," p. 110.

† The Tussar Silkworm, p. 9, 1873.

long spiral band, and used for binding the barrel of the matchlock to the stock, being, as the Natives say, unaffected either by water or fire. The cocoon consists of two kinds of silk; what it first spins is reddish, and of this the pedicle and outside network is made. This silk consists of threads of different lengths, but the rest is generally unbroken from beginning to end. . . . After the caterpillar has spun a layer of silk thick enough to conceal itself, it discharges some kind of gum or cement, thick and white, like plaster of Paris, and then, with its muscular action, it causes the gum to thoroughly permeate the whole cocoon and solidify the wall. In this manner it goes on spinning layer after layer of loops, and cementing them all together until the whole of its silk is exhausted, and the wall of the cocoon becomes so hard that it requires a sharp penknife to cut through it. The ring at the end of the pedicle, which has been spun round the twig, is a most necessary provision of nature, for it often happens that either the caterpillar has been unable to attach its cocoon to a leaf, or that, during the long time the cocoon remains unburst in the tree, the leaf or leaves to which the cocoon was at first attached become separated from it, and then the cocoon hangs suspended to the twig like a berry."

After eight or nine months in the pupa state, a moist spot is observed at one end of the cocoon. The moth is now about to emerge both from its pupa shell and from the cocoon. It secretes an acid fluid, which softens the cement of the cocoon, and enables it to separate the fibres sufficiently to allow its creeping out, it being, no doubt, assisted in this by its short pointed spines. The head of the moth first appearing with its antennæ, broad in the male, and narrow in the female, thus enabling the observer to note the sex, and to put them in pairs. The male moth generally flies away the night of his exit from the cocoon after his wings become extended and dried. The female rarely does, but during the first three days of her existence she lays her eggs, which hatch in about twelve days afterwards. The new life of the moth does not extend to more than eleven days. As you will see from the specimens in the case, the moth is a fine and handsome insect, measuring across the wings about six inches in the male, and about five inches in the female. You will notice the similar vitreous and transparent wing spots to those of the atlas moth. These spots are regarded with superstitious reverence by the Natives, who see in them a resemblance to the chakra or discus of the god Vishnu, and are therefore induced to consider the moth a sacred insect.*

The latest report on tussar culture is one from Major Coussmaker, dated Jangaon, Ahmednagar, on the 10th of January of the present year.

At Poona he has established a breeding establishment, and has planted many young trees for feeding the worm. He says it thrives well on *Lagerstramia indica*, an ornamental shrub fairly abundant in the cantonment of Poona. He has changed his plan of feeding them on cut twigs brought to them in captivity, and allows them, carefully watched, to feed in the open air on the growing shrubs. This was an evident change for the better; the larvae moulted in four to five days instead of five to eight days; they spun their cocoons in 28 to 35 days instead of from 40 to 50 days, and the moths emerged from the cocoons in 27 to 30 days, their eggs proving more fertile than under the former plan. He fed them also on *Carissa carandus*, and they thrive still better on this; but the *Lagerstramia*, after having all its leaves eaten off, being cut off and repotted, was in thick leaf again in a fortnight, and was ever sprouting, whilst the *carissa* did not sprout again after having been once denuded.

For the next season's experiments Major Coussmaker has planted between 700 and 800 plants of the following species—*Lagerstramia indica*, *Lagerstramia parviflora*, *Conocarpus latifolia*, *Carissa carandus*, *Zizyphus jujuba*, and *Pentapetia tomentosa*, so that we may look forward with a large degree of interest to the result of so practical an experiment to domesticate this interesting silk-producer in the Deccan.

Captain Brooke says:†

"In Chanda and Sambalpur, Central Provinces, when the cocoon crop is gathered, Keshias, a weaving caste, visit the villages and buy them from the rears. They are then, as soon as practicable, boiled in a lye made from the ashes of jungni stalks, a plant grown for the oil expressed from its seed. This process effectually kills the chrysalis; at the same time dissolving the mucilage of the cocoon. The cocoons are then stored for use. The method of reeling is primitive in the extreme, and to its imperfections I solely attribute the scant attention this valuable and

* Geoghegan.

† "Geoghegan, Silk Industry of India," p. 111.

very beautiful silk has hitherto received. A description of the process is as follows: The spinner, always a woman, sits on the ground; on her left is an earthen vessel, with a thickish rim, about six inches in diameter and three inches deep. The saucer is three parts filled with a mixture of potash and ashes, patted down to a level surface, and kept damp with water. Upon this the cocoons to be spun are placed, the outer portion of inferior and nearly useless silk having been first removed. The thread in ordinary use amongst the weavers is spun from seven cocoons; these are all placed at the same time in the earthen saucer, a filament is then taken up from each cocoon, and, being brought together, are rolled between the hand and left thigh of the spinner, which are kept damp by an acid solution of tamarind and water."

In Bengal the cocoons are put into boiling water to kill the pupa; in some districts, when intended for sale, they are put in boiling water and dried in the sun. In the Nizam's country the cocoons are loaded with dhobee's earth and alkaline ashes to make the reel. In the Midnapur District they are boiled in cow-dung and reeled by hand.

Captain Brooke says that in Seoni the pierced cocoons are wound, and that no Koshta rejects a cocoon simply because the moth has eaten its way through it. He has fallen into an error as to the moth's mode of exit from its cocoon. It separates the fibres with its legs and wing spine, and so creeps out. It has neither teeth nor mouth proper.

Each species of silkworm has two stores of silk, one on each side of the alimentary canal, and below its mouth it has two so-called spinnarets or orifices, through which the silk issues simultaneously in fine parallel filaments. As the silk is drawn out of the stores, the worm coats it with a varnish technically called gum, which contains a brownish-yellow colouring matter.

The tussar worm, in spinning his cocoon, takes short sweeps of his head from side to side, depositing the silk very closely in parallel fibres as he does so. It has been thought that the worm twists the silk as it exudes it, but this is not the case. Besides the gum which coats the silk, the worm secretes at intervals a cementing fluid, which it kneads by an expanding motion of its body through the whole cocoon to consolidate and harden it. This cement gives to the cocoon its drab colour.

There is a striking peculiarity about the fibre of tussar silk. I have carefully and thoroughly examined it many times under the microscope, and find undoubtedly that the fibre is flat and not round, like mulberry silk.

There is no doubt that it is to this property that tussar silk owes its glossy or vitreous look, reflecting a little glare of light from the angle of incidence on its flat surface, whilst the mulberry silk fibre, being round, reflects the light equally in all directions.

By some this property is considered a drawback, but by the time the fibre has become modified, and the flatness diffused in the loom, I think the lustre of the cloth is enhanced by it.

This tape-like appearance gives the fibre this disadvantage, that it is less homogeneous than the round fibre of the mulberry silk, and I find an undoubted tendency in it to split up into smaller fibrils, thus causing the silk to swell out when subjected to severe dyeing processes, particularly the bleaching one of recent date, thus giving a substantial and important reason why its coloured cements should be removed by gentle action.

The fibrils have a distinct structure, upwards of twenty in number, and seem compactly laid together, showing the striated longitudinal appearance of the fibre under the microscope. I dare say it is this fibrous compound structure, absent, as you see, in the mulberry fibre, which is an element in its dye-resisting power. I found permanganate of potash to be the best agent to separate these fibrils.

The diameter, from edge to edge, of a single flat fibre of tussar silk from the outer part of the cocoon, averages $\frac{1}{10}$ part of an inch, and from deeper in the substance of the cocoon $\frac{1}{15}$ of an inch, but the external fibres are much more variable than the internal. The thickness from side to side is $\frac{1}{1000}$ of an inch. The outside fibres are capable of supporting, without breaking, an average weight of seven drams, and the inner eight drams, whilst the usual amount of tension in all the fibres is one inch to the foot. The fibres, like all other silk fibres, are laid in the cocoon by the silkworm in pairs, united by their edges, and not by their flat surfaces.

All the *Saturniidae* fibres I have examined are more or less fibrous and flat, except the English species *Saturnia carpi*, or Emperor moth, which, in North Staffordshire, spins a beautiful cocoon in the heather of our moorlands. My son

has drawn for me an enlarged microscopic appearance of this silk, which shows its transparent and fibreless nature; and also, what is very curious, that the fibres are round, except where they come in contact, when they become flat, no doubt, from pressure. This apparent exception to the *Saturniidae* depositing flat fibres, and the *Bombycidae* round ones, which I had established, possibly points to the secretion of the sericine in the different species varying in fluidity, that of the *Saturniidae* being excreted in a more fluid state than that of the *Bombycidae*. There may also be a difference in the structure of their seripositors; but this I have not had an opportunity of investigating.

I am glad to state the other diagrams were prepared for me by Mr. Rider, a pupil of the Leek Art Class. The maps, showing the wild silk districts, were done by Mr. Alfred Moore, of Leek, and I have been much helped in the microscopic work by my assistant, Mr. Rigby, and I must not omit to mention the assistance rendered me by my printing staff.

It is a fallacy, held by some entomologists, that the worm twists the two threads together as it forms them at the orifice of its spinnarets, in all species, both *Bombycidae* and *Saturniidae*. The two threads are simply laid side by side, as you see in the diagrams. It would be impossible to twist the two threads without the worm itself revolving continuously with the omission of the silk, or for it to have spinning-wheels at the secreting orifice. I propose, therefore, to change the word spinnaret, which conveys an inaccurate impression, and substitute for it seripositor.

Leaving now the more beaten track of the natural history side of the question, I come to speak of its merchantable and art side. Finding many years ago that tussar silk opposed a resistance in no ordinary degree to tinctorial matter, I took an interest in the subject with a view of overcoming this resistance. In its small affinity, ordinarily speaking, for colouring matter, it ranks with the vegetable fibres of cotton and flax, and whilst in many processes it would come out scarcely tinted, the mulberry-bred silk would be found to have seized the colour with avidity. It, however, takes the aniline dyes, under certain conditions, moderately well. At that time, and for some years previously, little tussar silk had passed through the dye-time, and an attempt was made to introduce it in Macclesfield houses. About forty years ago an account of its irregular way of taking the dye, it for sewing silk for black, but, on account of its irregular way of taking the dye, it was abandoned. Mr. David Clarke, of Macclesfield, at that time, with his father, Mr. Jeremiah Clarke, were much interested in bringing it to the front; but, as Mr. D. Clarke informs me, the second parcel not coming from the dyer in a saleable state, a costly trial took place at Chester to determine whether the blame lay with the dyer or the silk. I believe it was decided in favour of the manufacturer, and against the dyer, who, unfortunately for him, had succeeded in dyeing the sample parcel successfully. However, the result was, that little or no tussar silk has been used for sewing purposes from that day to this. About twelve years ago I made many experiments in dyeing this silk, and had the satisfaction of seeing my way to further utilisation and improvement. In 1873 the firm of which I am senior partner, consisting of my brother and myself, exhibited at the International Exhibition at South Kensington, the result of progress up to that time, in a series of black and coloured silks, which were in advance of any similar effort, either English or Continental, as far as my observations or knowledge extended, and they attracted a good deal of attention and led to a further utilisation of tussar silk, then a drag in the market, except for dress silks for women and girls in the undyed and pleasing shade natural to it, which is fawn-colour.

The development up to that time had been that this silk could be dyed into any middle or dark shade of drab, slate, brown, green, violet, or dark red, whilst, to pale shades of blues, pinks, cerise, scarlet and others the dark natural ground colour of the silk interposed an insuperable barrier, as sulphur, or any then known bleaching agent, could not reduce the silk to a whiter state. The desideratum of pale shades led our quick French neighbours to study the composition of the brown colouring matter and to find a solvent for it. The credit of this achievement must be awarded to M. Tessié du Motay, who was led to try permanganate of potash, which was at that time attracting much attention on account of its great oxidising power on organic matter. He found the brown colourant yielded to this agent. Unfortunately, the oxidising action being too violent, the fibre of the silk as well as its colouration was affected, and the silk was tendered by the time it became white enough for dyeing into pale colours, so much as to render it useless. However, a secret had been discovered, and it was this, that oxygen, under certain combining conditions, united with the colouring matter, which then became separated from the silk. The object now was to apply the oxygen under gentler conditions. This M. Tessié du Motay again succeeded in doing, and in a very ingenious way. He brought into contact with the silk an insoluble body, which

on contact, should yield up an atom of oxygen in the nascent form which should gently unite with the fawn-coloured matter of the silk without attacking the fibre. This, although a rough method, solved the difficulty, and the silk, originally of the colour of this sample, can now be bleached in this way to that of the sample I show you, which is of sufficiently pale a ground to admit of its being dyed into any pale colour except white. The substance he found to comply with the required condition is benoxide of barium. Unfortunately, the process is too expensive, and prevents an extensive utilisation of tussier silk, but there is a probability of the principle being shortly applied by other methods, which will be at the same time cheaper and more within the legitimate sphere of dye-house technical operation than that of M. Tessié du Motay, I mean, whereby the nascent oxygen shall be presented to the silk in the vat from a solution, instead of from a solid as at present.

In 1874 I had the honour to receive a communication from Sir Louis Mallet, Under Secretary of State for India, asking me to communicate, "for the information of the Government of India, any details I might be in a position to furnish on the subject of dyeing the wild silk produced by the tussier worm".

On my report being received I was requested to make a full investigation of the subject, which divided itself naturally into two heads—a consideration of the silk, and of tinctorial matters. For the first, I found the raw silk as it comes into this country to be prepared by the natives of India and China in such a rude and filthy state as to interpose unnecessary obstacles to its taking the dyes. I felt sure that cleaner and more skillful methods of reeling and preparing the silk for the market would be accompanied by less resistance to tinctorial matter, as well as furnishing a greatly improved quality. At my request orders were issued for the collection in the different Provinces of India of a complete assortment of Native dye-stuffs as well as a supply of tussier silk.

In recommending the Government of India to have the Natives taught the dyeing of their wild silks with dye-stuffs indigenous to India, I had two motives—one to prevent the Native art of India being tampered with by the introduction of European fugitive dyes and crude colours, and another that they could be made to utilise, what their country has ever been so rich in, the remarkable variety of Native-grown dye-stuffs which, in other than wild silk fabrics, they have known probably for thousands of years so well how to use. To take dye-stuffs to India must surely be carrying coals to Newcastle. I have since received an extensive and most interesting series of India dye-stuffs and tanning materials, which I have at present under examination. I also received a quantity of tussier cocoons, and, not being able to have them reeled in England, I was authorised to go to Italy to see if I could have them reeled there, and effect my hope for improvement in the manufacture. By the introduction of a friend I obtained permission to visit one of the filatures in Piedmont, that of Messrs. Gaddum and Company. On arriving there I found an extensive mulberry silk reeling and throwing establishment, situated in a most beautiful valley, in one of the southern spurs of the Alps, about three hours' journey north of Turin.

On explaining my mission, and showing the wild cocoons, I was told there was not much chance of success, for they had several times tried them, and had found them difficult to soften, and impracticable to work; but knowing too well how natural is the tendency of mankind, in any new idea, to suggest objections rather than the means, I asked for permission to be allowed to try myself. The permission being generously granted, and every assistance kindly afforded me, I was taken to the reeling room, where about 100 young women were at work, with well-trained fingers, reeling the small Piedmont cocoons of *Bombyx mori*. The operation was interesting in the extreme, heightened as it was by their strange singing of old French songs, in a dialect not even understood by the Italians, a strange and all but forgotten tongue, which has to be learned by the mill-overlookers before they can communicate their instructions to them. I was told these girls were the descendants of Huguenot refugees, escaped probably from Provence to the Italian side of the Alps, at the Revocation of the Edict of Nantes, and that they still retained their patois and their folk lore; they worked hard for the few months of cocoon-reeling, from five in the morning until eight at night, for a franc a day; after work dancing and singing for the hour before bed time in the most joyous way. Apartments are provided for them at the factory, and when the reeling season is over they separate and return to their Alpine villages, to wait for the next season's work.

The operation of unwinding silk from the cocoon is as follows: A number of cocoons are immersed in an iron pan, in water nearly boiling, with a little alkali to soften them. A semi-rotating brush is placed over them, which quickly catches

the exterior fibres of each cocoon, and the more readily enables the reeler to find the windable thread. They are then taken out and transferred to the reeler, who sits leaning over an iron pan of about 12 inches in diameter, in which she has a few cocoons in hot water, the found-ends of several being in one hand. Four or six cocoons, as the case may be, are being simultaneously reeled into a single thread by the reel at her back which draws off over her head the cocoon threads; they dancing and turning in the water. When a thread breaks, or the cocoon is reeled, another is quickly presented from the lot in the other hand, the manipulation being one of great dexterity. Several years are required to attain proficiency, and it is not until the fifth to the seventh year that a reeler is entrusted with the most delicate reeling; the keeping of the size of the thread regular and free from rough places being the most important care.

It is this branch of the manufacture that in tussier silk is so defectively treated in India, the reeling being done in some instances round the naked knee-cap, but generally with this hand-reel.

I took some of my wild cocoons, and, with much difficulty and patience, after several trials, succeeded in softening them by the aid of long-continued boiling in water, to which was added soap, potash, and glycerine. When soft enough, one of the most skilled girls was told off to reel them for me, and, after ridding the cocoons of the outer and coarser threads, she reeled the thread of four cocoons into one, almost without a break, much to her own delight and to the surprise of my friends and myself.

The next day, the resulting tussier raw silk was taken to the throwing mill, and there made into organzine and tram, of such fineness as to surprise my friends, who said that they had no idea that tussier silk could be made so fine, and that they should think seriously about sending a person to India to collect tussier cocoons that their work people might wind them after their mulberry crop had been finished.

The usual size, that is, thickness, of thread of tussier raw silk of commerce is 152 to 255 deniers, that is, skeins of 1,000 yards long, weighing 9 to 15 drams. From some of the finer raw silk a size of six to seven drams is obtained, but it is generally coarser. From the cocoons, the reeling of which I superintended, I obtained a size of 51 deniers, or three drams, per 1,000 yards, a sample of which I have the pleasure to show you.

Here are samples of tussier raw and thrown silk I have received from M. H. Meyer, of Milan, obtained from cocoons I sent him. The size of the raw is 23-27 deniers, or 1½ drams per 1,000 yards; the organzine and tram are 50-55 deniers, or 3 drams. He found some of my cocoons very difficult to reel, no doubt owing to their age, and to not having been reeled before weather exposure. Fourteen pounds and a half yielded one pound of raw silk. He informs me that some cocoons he has just purchased in Marseilles are larger in size than those I sent him; they are darker in colour, but reel much better. He is obtaining from ten pounds of them one pound of raw silk.

Even a finer thread might be obtained, but as the fibre is only the 1/10th part of an inch, or three times as thick as ordinary silk, I think 51 deniers is a good and practicable limit when Native reelers can have proper appliances and be taught to be as handy as the reelers of Italy or the South of France. I daresay some of my hearers may remember the improvements which took place in reeling the mulberry silks of Bengal and Bruttia, when superior skill and machinery were introduced, a good many years ago. Before that time Bengal silks were held in very low estimation, and were very difficult to work, but after the introduction of better appliances Bengal silk was shown to be as capable of refinement, as any other; and Bruttia silk now commands, by its superior quality, the highest price in the market; and I have no doubt that, in degree, equal success lies waiting for the tussier silk industry.

I trust I may point to this manufacturing development and great improvement with pardonable pride, more especially as I am not a manufacturer, and could scarcely expect to find untrodden ground in a domain distinct from, although allied to, my own.

The new reeled silk is much lighter in colour, as you see, than Native reeled, and has very much more lustre; in fact, it is the most lustrous in the undyed state of all silks, and possesses greater strength and tension. I found, what I expected to find, that the silk thus reeled dyes much more easily; more shades and lighter ones, can be dyed upon it than Native reeled, it has no disagreeable smell, and only loses two ounces per pound in being cleaned for dyeing, where Native reeled tussier loses in some cases as much as six to seven ounces per pound, and never less than four to five ounces. It is as clean, to use a technical term which

means free from slubs and irregularities of thread, as ordinary silk. The cost of reeling new and good cocoons, and manufacturing them into organzine and tram, is about seven shillings per pound, and it is certain to make its way in many fabrics where extreme fineness is not required, and for a variety of purposes in passmenterie, trimmings, braids, scarves, broad and narrow goods. It is beginning to be largely used for these purposes in France. Its price has lately risen, whilst that of other silks has either remained stationary or actually depreciated.

I have urged on the Government of India the importance of introducing to the natives of India the European modes of reeling cocoons, and some time ago drew their attention to an invention which simplifies and economises this operation.

Mr. Mackenzie, Engineer, of Milan, has introduced a Milan House of filatures, who have invented and patented another mode of reeling, by which skilled labour is dispensed with. If this machine is pronounced by experts to be a success, there is no reason why cocoon-reeling should not be carried on in any village home as flax-spinning was formerly.

M. David, the largest ribbon manufacturer in St. Etienne, seeing this improved manufacture and dyeing in the Indian Section of the Paris Exhibition, where they were first displayed, offered to buy all the cocoons produced in India if the price would not be more than one franc per kilogramme—a price which Dr. Birdwood assures me is reasonable. He has applied to the India Government for 2,000 kilogrammes of cocoons for experiments at his own cost. It would be a very good thing for a trade to spring up in tussier cocoons. The Natives could easily be encouraged to breed a larger supply, whilst improvements in reeling would require time, and would meet with obstacles of race, religion, and habit difficult to overcome; also the enterprise in this direction would have to be purely private and mercantile, as I think the Government of India would not enter into commercial undertakings, but would probably, and certainly ought to, give most strenuous encouragement and help to stimulate the further spread of this most interesting industry.

I was requested by Dr. Birdwood last year to exhibit the developments of which tussier silk was capable in the Indian Section of the Paris Exhibition. Sir P. Cunliffe Owen entered most warmly into the idea, and took the greatest interest in it throughout, giving me all the encouragement and help required to make it worthy of being represented side by side with the beautiful objects in which India had determined to assert herself. In no exhibition before had India been shown in a manner so worthy of the gorgeous East. It was the India of the artist which asserted itself at Paris—the old historic land, from which are manufactures in brocades, printed calicoes, jewellery, ivory carving, and pottery may still draw their highest aspirations. Neither had ever before been so much done for the promotion of Indian commerce. These results are entirely due to Sir P. Cunliffe Owen, and to one who so ably assisted him, Mr. P. C. Clarke.

In the wild silk exhibits which I was entrusted to bring together in this Section, not only were the improvements shown in manufacturing and dyeing, to which I have alluded, but another and more decorative phase, and one developed, so far as I can gather, for the first time in the history of either the East or the West—that of printing. It had struck me that fabrics made of tussier silk, either of Native or home manufacture, would be susceptible of much enrichment if they could be printed upon. After many fruitless attempts I at last succeeded, and since that time I have had the satisfaction of succeeding, in applying and fixing a much wider range of colours.

Thinking that designs of an Eastern type were naturally the most applicable to cloths of this wild silk, I have obtained, by the courtesy of Dr. Forbes Watson, the loan of a large series of wood printing blocks, of Native design and workmanship, from the India Museum. I have used them for printing nearly all the illustrations of my lecture, and have placed a few on the table to show how beautifully they are cut. In England the finer details would be in copper, but in these the hardest wood has been chosen and most skillfully cut. To complete the consistency, I have adhered to the use of Native Indian colours and colouring matters. You will notice the deep rich red of the India print in madder or munjeet, the good-toned and permanent indigo shades, as well as a variety of other well known Native dyes. The designs on these blocks are extremely interesting, and if I had time to exhibit the whole series of the impressions I have taken from them, I am sure you would agree with me how they abound in originality and

beautiful drawing. If from this we are led to think generally of the Native art of India, we may justly feel some sorrow and regret that our influence there does not tend to perpetuate it, and regret with Dr. Birdwood, as he so well describes in his hand-book, that it daily deteriorates. If it were not that Sir Cunliffe Owen is sitting so near to me, I might be led to suggest whether we could not try the experiment of turning the tables somewhat, and that, if we must send our art masters to India, we might at least import some from there to try to bring us into better ways.

Truly our credentials to teach an artistic people are in a sad condition; if we take but architecture for example, we are but copyists. Imitation seems to be the evil genius of our time, and even what little originality we have to be thankful for in decorative or structural art, is no sooner born than it is preyed upon by an unprincipled selfishness and rascality, which, in the greed for gain in such matters, sees no distinction between *meum* and *lum*; whilst, on the other hand, so strong is the existing jealousy to protect that which should benefit all, that we may scarcely with safety look over our neighbour's palings to see how green his lawn is.

I would call attention to the sweetness with which the colours repose on the natural and unbleached ground of the cloth, as well as the greater sharpness and depth of those printed on bleached grounds.

Many of my examples are painted in print colours, on outline printed designs, an ancient and most interesting mode of decorating cloth, which I have revived.

I think you will agree with me that the material so decorated is beautifully suited to wall hangings, curtains, coverlits, and all kinds of furniture work, and whilst not having quite the brilliancy of the mulberry silk in its printed state, it has a richer and softer surface than those of cretonnes or challis, whilst its lasting qualities are superior to those of any other material.

Messrs. Durant, of London, have kindly informed me that tussier raw silk comes from China, and they believe the large shipments of two years ago were principally owing to the famine in the districts of productions. Scarcely a bale, they say, has come forward during the present season, nor do they expect any at the present prices. The present price of tussier raw silk in London is 4s. 6d. per lb. The stock in London is China tissore. It is collected in the district of Chefoo and shipped from Shanghai. The price of Indian tissore cloth is about 2s. per yard, 34 inches wide.

There is a very large quantity of China tussier cloth exported from London to the colonies. An immense trade would be developed in India if better qualities were woven there.

TUSSIER.

The following table shows the state of the London market in tussier silk for the last few years:

YEAR.	Stock, January 1st.	Imported.	Consumed.
	Bales.	Bales.	Bales.
1874	662	None.	168
1875	494	None.	319
1876	175	427	174
1877	423	1,037	281
1878	1,181	837	736
1879	1,252		145

The stock, February 1st, was 1,137 bales, with a consumption for January of 145 bales. Should the demand continue at this rate, the supply would be insufficient for the year if no more comes in.

The average consumption for the four years ending 1877 was 235 bales, whilst the consumption for 1878, the year that attention was drawn to it by the Paris exhibits, more than trebled itself, the purchases being 736 bales for that year.

The following table shows present prices of silks in the London market (April 18th, 1879):

		s.	d.
China raw Tsatsee, No. 4, per lb.	...	15	6
Canton ditto No. 1, do.	...	13	6
Ditto ditto No. 4, do.	...	11	6
Japan raw Marbush, No 2½, per lb.	...	16	0
Italian Organzine	...	25	0
Bengal raw	...	14	6
Brutia do.	...	23	0
Tusser do.	...	4	6

The great improvements made in this country and on the continent of late years in carding and spinning machinery have enabled manufacturers to utilise all the silk that could not be reeled, such as pierced cocoons and all kinds of waste silk.

Mr. Clayton has been kind enough to lend me, for my lecture, illustrations of this useful phase of manufacture, specimens of spun tusser silk in each stage of manufacture, from the first carding or dressing operations to such perfected fabrics as you see before you in cloths of varied design and substance, yarns for weaving and sewing, and shawls in plain and printed states.

There is now a great demand for tusser and other wild silk waste, and England possesses more than sufficient machinery to spin all that can be imported.

Another form for the use of tusser silk is the manufacture of embroidery silks, and their application to cloth by the needle. I have had manufactured a few silks, which have been arranged by my wife for illustration this evening. She has also worked a few pieces of tusser cloth in these silks in various designs. The larger piece of embroidery, which is unfinished, is a trial piece. Mrs. Wardle began to work first with untwisted silk, which, as may be observed at the left corner of the work, has a fluffy appearance; therefore I recommend for embroidery purposes a slightly twisted silk, which I think will be an improvement on crewels, and possibly filozelles, and so prove a useful industry.

DISCUSSION.

A meeting of the Indian Section of the Society of Arts was held on Friday, May 2nd; Sir P. Cunliffe Owen, C. B., K. C. M. G., in the chair, when Mr. Thomas Wardle, F. C. S., F. C. S., read a paper on "The Wild Silks of India, principally Tusser".

The Chairman, in introducing Mr. Wardle, said this evening's subject was one of the most practical results of the Paris Exhibition. There was nothing which more interested the French people than Mr. Wardle's collection, and India was specially honoured by the International Jury in receiving the Grand Medal of Honor for silks, as well as for tea. The manufacturers from Lyons were greatly interested in the Indian collection, notwithstanding it was contained in a very unpretending set of cases; and he must confess that when this matter was first brought under his notice by Dr. Birdwood, he did not think anybody could have had an idea of the practical results which would be achieved by Mr. Wardle's efforts. Mr. Wardle had been for some years employed by the India Office in making inquiries as to the development of this industry, and particularly as to the application of dyes from the plants to be found in the neighbourhood in which the silks were obtained. One could always see the mistakes which had been made after a thing was over, and he certainly thought those in charge of the Indian collection (of which he himself was one) might have taken more trouble in developing it. Mr. Wardle had great difficulties to contend with, but, notwithstanding, he would repeat that the specimens of silk now exhibited were the most practical results of any International Exhibition, and Englishmen might be proud to think that the Empire of India had come out with such glorious results; but he was afraid many of them hardly realised what Mr. Wardle had done. The Lyons

manufacturers, however, saw the importance of it; they were now in correspondence with him, and were anxious that his paper should be read before the Lyons Chamber of Commerce.

Mr. Wardle then read his paper, as printed in the preceding pages, after which there was the following discussion:

Dr. Birdwood said: Ptolemy was the first to use the word *serice* for China, or, rather, the Northern part of it, known later as Cathay; and the word is derived from the Chinese name of the silkworm *ser*. In Coran *ser*, whence the Greek *Ser*, the silkworm; *Seris*, the people furnishing silk. The Latin *sericeus* had been traced direct to the Mongol *serikah*; and the *serikah* of Isaiah, xix. 9, has been supposed to be silk. If the latter identification is correct, the trade in silk between the East and West went back to the remotest antiquity. But there is no undoubted proof of a knowledge of silk in Europe until the time of Alexander. Aristotle certainly describes the silkworm, and the Greeks of Alexander's expedition must have seen both Chinese and Indian silks in India, and tusser cocoons. When Chinese silks and the manufacture of tusser silk were first introduced into India is also an obscure question. In the *Ramayana* silks are mentioned among the presents made to Sita, the bride of Rama; but whether these were of Chinese or Native manufacture cannot be determined. In the *Mahabharata* it is said that the "Chinas, Hunas, Kashas, and Cauchas, who lived in the mountains", brought as tribute to Yudhishtra "silk and silkworms". If the "Chinas" here really were Chinese, the question would be settled. But I believe Colonel Yule has somewhere stated that the "Chinas" of the *Mahabharata* were some tribe of the North-Western Himalayas, near to the Dards; and so everything is left in obscurity as to the first introduction of Chinese silk into India. It is not even known whether the Arabs, on their first arrival in India, found the silk manufacture already going on there, or introduced it themselves. Pliny's account of the silkworm is taken from Aristotle, but he knew Chinese silk well, and he also tells us that in his time the wild silk of the Tigris Valley was woven into gauze. Pausanias, about 190 years later, also describes the silkworm; and the allusions to Chinese silk by the Roman poets from the time of Augustus are too numerous to cite; but it is clear that before the time of Julius Caesar silk was but little known in Europe. Procopius, about A. D. 550, tells the story of the introduction of the silkworms' eggs into Constantinople in the reign of Justinian, and how the culture of silkworms and the manufacture of silk spread from that time all over Southern Europe is a three-fold tale. Mr. Wardle has been so good as to refer to my report of 1854 on tusser cocoons in Bombay was the late Mr. Edwin Heycock. A meeting was held in Bombay in 1857 to found the Victoria and Albert Museum, and Mr. Heycock, in pointing out how useful it would be in bringing jungle products into public notice, instanced the waste of tusser cocoons in the Bombay Presidency, simply through the ignorance of merchants in Bombay of their existence. I reported this matter to Government, and a large quantity of cocoons were at once sent to this country, and inquiries made all over the Presidency, which elicited the fact that tusser cocoons were as abundant in Bombay as in Bengal. Afterwards, Major Consmaker and Dr. McKenzie paid great attention to the matter, when at last the task of practically investigating the comparative commercial value of tusser silk was fortunately taken up by Mr. Wardle. His paper has scarcely done justice to his own labours in the matter. It is perhaps too encyclopaedic, and at the same time says too little of his own experiments and their practical results; and as I know more of Mr. Wardle's work than perhaps any one else in the room, I doubly regret that I am under the necessity of having to leave at once. Briefly, Mr. Wardle has at last, after years of patient research, succeeded in spinning tusser silk as fine as the finest Chinese silk, and in giving to it every fineness which Chinese silk will receive. He has thus made it merchantable wherever silk is reeled, spun, woven, dyed, printed, or used for personal or household adornment. Here is the prospect of a great gain for India, and Mr. Wardle deserves for it the grateful acknowledgment of every one interested in the welfare of India. Sir Philip Cunliffe Owen said he regretted not having put Mr. Wardle's exhibits more forward at Paris last year; but I am an eye-witness to the fact that Sir Philip did everything in his power to promote the deep interest of India in Mr. Wardle's work. Universal attention was drawn to it in Paris, and if I heard of tusser as much due to Sir Philip Cunliffe Owen as to Mr. Wardle. I must add that I was perfectly astonished on entering the room and seeing the samples of printed tusser—in Indian dyes, and stamped with Indian blocks—exhibited here to-night by Mr. Wardle. I could not have believed it possible for him to have made such

progress in the short time since I saw his last tentative efforts; but while I congratulate him on the taste and ability in the dyer's art these beautiful fabrics show him to possess, I cannot altogether wish him success in his imitation of Indian designs. Imitation is no more good for us than the natives of India, nor is it possible to welcome these efforts to take a manufacture out of their lands simply because, being rulers in India, we are in a position to force their hands.

Mr. Francis Cobb said there was no doubt that the demand for this silk had increased recently, partly in consequence of a change in fashion. There had been a run on it by the French; and, owing to the change in the method of dyeing, almost every colour, even of pale rose, could now be dyed. How much was worked up in the fabrics which were supposed to be all genuine China silks, he believed, it was impossible to say, especially those subject to heavy dyes. During the last two years the demand in China for this tussier silk for France had been so great, as to seriously interfere with the supply required for the country, which, as it was the cheap silk of the country, was very large. The French had now pretty well exhausted China, and hence one great cause of the anxiety which was noticed at the Exhibition; they were all aware that they were fast exhausting China, and turned with the greatest delight to the specimens received from India, showing that there was a new field open to them. He trusted, however, that the whole of the industry was not destined to find its way to France. Although they had made some rapid strides there, we had also been leading the way in this country, and but for the fact that they beat us in the cost of labour, he had no doubt that the tussier industry would have developed itself to a greater extent here than in France. He might say that the *Attacus ricini* was a worm which was a good colonist—it might be taken to almost any of the colonies and it would thrive. It fed on the castor-oil plant, which grew wild in most of our colonies. For instance in South Africa, when land was cleared, a castor-oil plant would immediately spring up. Mr. Wardle had shown how valuable that particular silk was, and it might be grown to a very large extent in the colonies. He did not believe that these tussier silks, now that attention had been called to them, would ever go out of demand, and all that India or our colonies could send would find a ready sale for many years to come. The *ailanthus*, of which they had heard so much seven or eight years ago, and to which Lady Dorothy Neville devoted a great deal of attention, seemed to have failed, and the reason was that the labour expended upon it did not produce a result which would pay its cost. He doubted whether in such a country as this we could ever rear the *ailanthus* worm to a profit. When he was giving a great deal of attention to this matter some time ago, he remembered being informed that there were thousands and thousands of acres of trees in Assam which were infested, as it was termed, with these wild silkworms, and that they only required gathering. He presumed that was the *Moonga* which Mr. Wardle had spoken of. In the North of Ceylon, again, there was a large forest district of which very little was known, but there was a cocoon found which was open at the end, and the silk came up there and formed a loop. When that cocoon was treated with caustic soda, or potash, it became softened, so that it could be reeled pretty readily, but he had grave doubts whether reeling was ever worth the trouble and cost for any of these tussiers. Although Mr. Wardle had succeeded in producing very fine reeled silk in Italy, when that silk was completed and reeled it was not equal to China or Italian silk in the same condition, and he doubted whether it was a kind from which we should ever get any very fine results. He would rather adopt Dr. Birdwood's suggestions, and take the coarser manufactures of this silk, and trust to the play of light and shade to produce that rich appearance which he spoke of. Mr. Wardle said the tussier silk which came from China appeared to be better reeled, but the reason was that the Chinese reeled the cocoons at once: they did not wait until the chrysalis was killed by heat, and at that time it was more flexible and had greater elasticity and brilliancy. This was a thing to be noted, and to be introduced into India, to secure the cocoons whilst the chrysalis was still in them, and thus to produce them in a condition in which the silk might be spun. No doubt Mr. David would be very glad to get a great deal of this silk for one franc per kilogramme, which Dr. Birdwood thought a reasonable offer; but he thought there would be no difficulty in selling it in this country for two shillings per kilogramme. He did not think that it was the price which stopped the supply, but rather that it was the French demand which had pretty well cleared all the available silk from China, and that they would not send any more, except at a very enhanced price. With regard to the difficulty in dyeing, he believed that had been overcome by Mr. Wardle. He had been as yet unable to find out how it was done in France, and the last answer he got to his pertinacity was that he was not likely to find out, as it was a trade secret. Still he had hopes that Mr. Wardle, in his persevering manner, would

eventually find this out for himself, and would be able to compete with the French even in the most delicate light colours. The exhibit he had shown was the most remarkable he had ever seen, and the dyes were such as, a few years ago, he should have thought were impossible.

Mr. Davenport said Mr. Clayton, who had exhibited some of this tussier silk at the Paris Exhibition, was present, and he hoped he would give them some information upon it.

Mr. Wardle said he wished to supply an accidental omission in his lecture, and that was with regard to the adaptability of tussier silk for spinning. He had there some beautiful manufactures of spun tussier silk, but unfortunately they had been so delayed by the railway that they only arrived at the last moment after his lecture had gone to press. They had been kindly lent by Messrs. Clayton; Mr. Clayton was present, and he hoped would explain them.

Mr. Clayton said that some four or five years ago, a large shipment of tussier waste came over here, and was lying about a long time in Lower Thames Street. From thence it was taken to Manchester, where it also lay for two or three years, as no one would look at it. He accidentally came across some samples, and determined to try some experiments. The first trial was so promising that he got further samples, and the result was that they showed some of the sliver to the Firm of Sir Titus Salt and Company with the idea that this silk could be mixed with brown mohair, which that firm were spinning at that time. They at once asked him not to mention it to any one else, but to get a few skeins of sliver prepared, and to bespeak the whole parcel of this material. They got an order for 1,000 weight, but in working it up Messrs. Salt discovered some hitch or defect in the spinning, and had to discontinue it for the purpose originally suggested. His firm had by this time the whole parcel on their hands, and therefore they had to endeavour to utilize it themselves. After repeated experiments they discovered the means of giving it spinning qualities, and they introduced it into the market in the shape of threads for various purposes. The trade had gradually grown, and twelve months ago they had got through the whole parcel of raw tussier waste, which they found afterwards was all that was brought from China. After those threads and cloths were shown in the market and sold, they found the raw material came over very slowly, and at present there did not seem any prospect of further supplies. This led him to investigate the question of developing the trade from India. They had no longer any of this raw material coming from China. There was not a bale in London, and, as far as he could learn, not one on the way. The importance of this branch of tussier industry would be apparent when it was remembered, as was stated, that, in the reeling off the cocoon, to obtain one pound of raw silk ten pounds of waste were made. According to Mr. Wardle's statement the larger portion of the tussier cloths came from India, and he was rather at a loss to know what became of the enormous quantity of waste produced in making them, inasmuch as, to his knowledge, none had ever come to this country, except a few pounds. It was, therefore, of great importance to those interested in Indian produce to get at the reason why this waste was not collected and brought to this country. Unless some was brought from India or elsewhere, the industry would die out for want of raw material.

Mr. A. Rogers said he had seen this very waste being spun in Messrs. Allen's mill in Bombay. The cocoons were brought in large bales from the jungle just squeezed together, and flattened like a cotton bale. This waste they crushed, and teased it out into very nearly as fine sliver as was shown on the table; and he was perfectly astonished at the cloth made from this waste. He brought home some specimens, and gave them to the Manchester Chamber of Commerce, where no doubt they could be seen. Another curious fact which came under his notice was this: There was a man in Bombay, a barber, who made a most remarkable discovery with regard to tussier silk. He showed him the produce of silk which had been spun by a hybrid worm, produced between the common Indian tussier worm, and what he said was the Japanese *yama-mai*. Whether that was the case he could not say, but he certainly showed him the two worms, which were about the size of his little finger, and he informed him that the silk produced was much freer from tannin than the indigenous tussier found in the jungle. He showed him some specimens, and, so far as he could judge, the hybrid silk was far superior to the other. Mr. Morris assured him that this hybrid would feed on the commonest trees in the country, particularly all trees of the fig tribe. If that were the case, and a superior kind of silk could be produced by hybridisation, one could see what an immense future would be open for production, and it would especially afford a means of employment for the mass of the people, especially women. In the higher

classes the women were never allowed out of doors, and it would be an inestimable boon if they could be induced to amuse themselves by rearing worms and producing silk. Even in the years of famine the trees on which the worms fed never failed, so that in such seasons of disaster the existence of this industry would be an immense benefit to the people.

Mr. R. F. Mackenzie, being called upon to describe his machine for reeling silk, which is now at the India Museum, said he had great confidence that it would work tussler silk admirably. They tried a small experiment with some cocoons, and it had failed, but the failure was not due to any fault in the machinery, or in the softening process, but he discovered from Mr. Moore that the cocoons were 12 years old; consequently the chrysalides inside had got decomposed, the air and water got inside the cocoons, and they would not float. Of course, if the cocoon was fresh it would not imbibe the water, and he had no doubt the silk could be reeled very finely, because one cocoon of tussler would yield a thread which could be worked, and two or three threads would be equal to ten of ordinary Bengal silk. His machine was certainly more applicable to a Bengal cocoon than to any other, and the chief purpose in inventing it was to work these cocoons in large numbers. Of course it would work every other kind, but the advantage would not be so great. When there were ten cocoons going into one thread, as hand labour was depended upon to feed all those ten threads, it became a serious matter, because it took five or six years to become an expert reeler. In India he did not think they were more expert or intelligent. The machine could be seen at work on Mondays, Tuesdays, or Wednesdays, at the India Museum, on Indian silk. Hitherto it had been applied to Italian silk, and had worked admirably. He must say that he was much indebted to the authorities at the India Museum, and to the Indian Government, for the assistance they had given him; and he thought they were all much indebted to the Government for attending to these matters, which, he thought, were quite as important as sending armies to conquer a country. To show how easily it could be worked, he might say that he asked Dr. Forbes Watson if he could procure an intelligent boy or girl to work it, and they were fortunate in getting an intelligent girl of 14. They gave her three hours' instruction, and then set her down to reel. She began with one thread and reeled two threads, which was sometimes equal to twenty cocoons. They were doing Italian cocoons, which were much more easily reeled, and about a week ago she reeled three threads with six cocoons each, and two threads of the Bengal cocoon with ten cocoons each, and he could safely say that the produce would compare with anything ever shown in the London market. They were now making arrangements to send out a staff to India and China to do the thing on an extensive scale. He believed that the tussler cocoon could be reeled as fine as anything, and it required but few threads. It was true that the threads were flat, and that was a point which made it a little difficult. The only way he could account for that, if he might venture to do so, was from the fact that there was a great amount of woody matter in the tussler cocoon, and very much less gluten, while there was a great deal of moisture, and no doubt the silk moth used the threads in a very wet state, and in consequence of the great heat it became very hard, and that must flatten the thread to a very great extent. Some time ago there was an idea of introducing cocoons and spinning and reeling them in this country, but it had not been carried out. In Asia Minor, however, and all over France and Italy, it was carried on largely, and he did not see why they should not be reared in England, or why the Indians should not be put in a position to reel from them and to send the orgauzine and tram to the English market.

Mr. A. Rogers said the introduction of this machine into India would depend a great deal upon the price at which they could be supplied, and a great deal would depend also upon whether the chrysalides required to be killed for the purpose of reeling. A great many of the Natives, particularly the followers of Vishnu, were averse to the taking of life in any shape whatever, and though they would spin or do anything with the cocoons gathered in the jungle from which the moths had emerged, they would not do anything with cocoons in which there was a live chrysalis.

Mr. Wardle, in reply, said some of the observations made would be found replied to in parts of the paper which he had not read. Dr. Birdwood did not intend to speak technically when he said that tussler silk could be made as fine as China, because he did not think that could be done. With regard to the danger he feared of doing injustice to the Native people, he could assure them that he shared his feeling, but if the increased cultivation and improvement of this undeveloped industry were worth anything, they must tend to give the natives of India more

interest in their work, employ a much larger number, and result in the production of a vastly increased quantity of loudly called-for silk, as the discussion that evening had so satisfactorily shown. He also shared Dr. Birdwood's desire for originality in decoration, but the inventive faculty was an uncommon gift; and there could be no imitation in using for a portion of his lecture illustrations the actual printing blocks the Natives had themselves made and used; it was using the actual ornament, and not a simulation of it. With regard to the suggestion of first dyeing this silk one colour and then another, and thus making two vivid aniline dyes produce a better tone, they had a much quicker and more satisfactory mode of doing it by simply putting two or more aniline dyes in the vat together. There was no doubt that any artistic colour could be matched with aniline dyes; the only difficulty was that mordants had not yet been discovered for that tinctorial matter, so as to allow the silk to be dyed in permanent colours. Hitherto, they were all more or less fugitive. With regard to Mr. Cobb's interesting speech, he could allay his anxiety with regard to tussler silk being dyed in weighted blacks in France, and then used to adulterate the best silks. The tussler worms seemed to share his aversion to heavy blacks, and secreted their silken fibres in such a way that they resisted the weighting processes more than they resisted tinctorial matter. He had known it only increase a few ounces to the pound in an attempt to dye it a heavy black, when China silk had increased 32 ounces; but with regard to its being mixed with better silk for various articles in colours, it was so used in France last year, and in England at present. With regard to fashion having brought in these silks, he thought it was just the other way, and that it was the fact of tussler having been made amenable, by improved dyeing, to general requirements, which had led to manufacturers using it. He did not think the French were one whit in advance of what could be done in England in dyeing this silk, both by his own firm and several other principal dyers. The secret which Mr. Cobb could not discover from the French was this, and this only, for he was quite *accourant* with the development of French dyeing (but the air of mystery imparts a charm), it was the *bleaching* of the silk before dyeing, for unless this were done, the pale colours could not be dyed; this effected, the rest was easy. He could have brought all shades of colour, even much paler than what he had shown. He was very glad to hear Mr. Cobb say tussler cocoons were worth more than M. David estimated them at; so much the better for India. With regard to the late supply of tussler waste, he was afraid distress in China had something to do with it; but with regard to the waste in India, he did not think there was so much as 12lbs. to one of silk. The Indians were very economical themselves; in one part of India they used the outer part of the cocoon for making armlets, and even brooches of; but still, if the people could be stimulated to grow more and export it, there would be plenty. With regard to the hybrid worm mentioned by Mr. Rogers, he doubted if he were correct, because the Yama-mai was a Japanese species, which was an oak-feeder, whereas the tussler worm fed on very dissimilar food. Still it was very probable there had been a hybrid formed between the tussler and some other kinds. The tussler cocoon contains no tannin or woody matter. Mr. Wardle concluded by exhibiting a cocoon from South Africa of an immense size, being upwards of a foot long, the inside of which was filled with a number of smaller cocoons, which, he said, was the most curious thing he had ever seen. Still, there was something analogous to be found in some parts of England, minus the matted outer covering.

The Chairman proposed a vote of thanks to Mr. Wardle, not only for the work he had done, but for the modest way in which he spoke of his labours. He knew nothing about the subject himself, but he had taken a great interest in it from the few words he had with Mr. Wardle, about eighteen months ago, when he pointed out the advantages which would accrue from the encouragement of this industry. He would take the opportunity of laying this paper before Sir Louis Mallet and before the Colonial Office, and he hoped some solid advantages would result from the evening's discussion.

The vote of thanks was unanimously passed to Mr. Wardle, and the meeting adjourned.

The Indian silks, diagrams, entomological specimens, &c., shown by Mr. Wardle, to illustrate his paper, were borrowed by Messrs. Liberty and Co., of Regent Street, who devoted a special room for their exhibition.

A 79
21

No. 77-84

Extract from the Proceedings of the Government of India in the Home, Revenue and Agricultural Department (Agriculture and Horticulture),—under date Fort William, the 13 February 1880.

READ—

A despatch from the Secretary of State No. 144, dated the 11th December 1879, forwarding copy of a letter from Sir Philip Cunliffe Owen relative to a collection of the wild silks of India presented by Mr. Thomas Wardle to the Chamber of Commerce of Lyons, and also copy of a lecture by Mr. Wardle on the wild silks of India, principally tasar.

ORDER.—Ordered, that a copy of the above despatch, and its enclosures, be forwarded to Local Governments and Administrations, and that attention be invited to Mr. Cobb's remarks, in the discussions printed at the end of Mr. Wardle's lecture, regarding the extensive demand for tasar silk, and to Mr. Clayton's observations about the uses of tasar and other wild silk waste.

(True Extract.)

Etchell

Offg. Under Secretary to the Govt. of India.

To
The Secretary to the Government of Madras

S. C. D.—Home, Rev. & Agric. Dept. 1880—12-230-231

*Please
no enclosure*

Recd 15/2/80

*I enclose a copy
of letter from
Sir Philip Cunliffe Owen
relative to the
collection of the wild
silks of India
presented by
Mr. Thomas Wardle
to the Chamber of
Commerce of Lyons*

James

No 1154.

Yours Govt of India
A. R. and apri:

103
74-8a

Ordered that the above Extract, with a copy of the Secretary's States Despatch and its enclosures be communicated to the B^d of Res., the Chamber of Commerce, the Trades Association and others.

Latd 13^e Sep-80
 Recd 23rd Feb 1880

Συνολικά

Col. Beddow
Dr. Beebe
Dr. Short.

of L. C. 26: March 1879
N^o 729.

PRINT
1. 5. 80

No 479 - 2009/2010

Forwarding a
copy of Secretary of
States Despatch on
letter from Sir Philip
Cairliffe Owen relative
to the collection of the
wild silks in India.

With Order

20 Feb. Boarding Pass for wife and child:
1. The Chamberlain, Chamber of Commerce
1. do Trade Office and on
1. _____ East of the Boarding
1. _____ George and Company of Dr. One and
1. _____ D. J. Street