

Register No 204
Sheet No 229

III. The area of the village of Rangani corresponds to that given in the account of 1721 & 1724. The present village found no fault with the former.

III Wet lands depend upon one ^{large} tank. It is fed by Jungle streams. In fact there are no waste lands in Slt. Rose plain as waste are fallow the occurrence of
Wet lands are fallowed 2 times to some extent. Dry lands are of long waste. The average wet fields of the village are 232 that there for since the same amount is
used in the cultivation of the village of 7 of the Reporters. Soil is not of a superior quality.

The conference was held
 at Pleasant Lake
 on 21/9/55

St. Louis and Rome
1882

Extract granted to
Amhalassona Radam
Lauise on the 28th inst 1800

Maximum Minimum
 39.9 - 1.7
 10.3 - 5.9
 Average rate of Great Salage per acre
 10.5 - 6
 10.5 - 6

Details of average in course of 15 years of the Gumbur on wet lands
 in the Taluk of Ramnath in Ramnath

From 1856	Cultivated	Grass culture	of which	Average	Value of the	Rates	
To 1870	area in		Gumbur	prices	quantity of	per	
	acres		shown at 100	of grain	grains given	acre	
			percent		in foot		
1856-1870	A C	R In	R In	R In	R In	R In	
35,981.2	58,869.50	2,859.4	2	91417.5	2.10	239970.8	6.5

Any rate prevailing in Gumbur Taluk of Ramnath in Ramnath
 Maximum Minimum
 1.11 - 1.07

Ramnath